

Journal of Family Strengths

Volume 17

Issue 1 *Innovative Practices to Eliminate Health Disparities*

Article 3

5-1-2017

Stigma and the Reluctance to Address Mental Health Issues in Minority Communities

Gary L. Kreps

George Mason University, gkreps@gmu.edu

Follow this and additional works at: <https://digitalcommons.library.tmc.edu/jfs>

Recommended Citation

Kreps, Gary L. (2017) "Stigma and the Reluctance to Address Mental Health Issues in Minority Communities," *Journal of Family Strengths*: Vol. 17 : Iss. 1 , Article 3.

Available at: <https://digitalcommons.library.tmc.edu/jfs/vol17/iss1/3>

The *Journal of Family Strengths* is brought to you for free and open access by CHILDREN AT RISK at DigitalCommons@The Texas Medical Center. It has a "cc by-nc-nd" Creative Commons license" (Attribution Non-Commercial No Derivatives) For more information, please contact digitalcommons@exch.library.tmc.edu



Stigma and the Reluctance to Address Mental Health Issues in Minority Communities

Acknowledgements

My thanks to Dr. Lovell Jones who invited me to speak about mental health disparities, stigma, and communication at the National Health Disparities in America Workshop that encouraged me to work on this important topic.

Introduction

For the members of ethnic and racial minority groups in the United States, health disparities lead not only to inequities in outcomes from serious physical health problems, such as cancer, heart disease, HIV/AIDS, and diabetes, but also to poorer outcomes from serious mental health conditions, including stress-related, manic-depressive, and psychotic disorders, as well as suicidal tendencies (Allen, Balfour, Bell, & Marmot, 2014; Bostwick et al., 2014; Jimenez, Cook, Bartels, & Alegría, 2013; Le Cook et al., 2014; Williams & Williams-Morris, 2000). Although current epidemiological evidence suggests that the actual incidence rates of mental health problems do not differ significantly across various racial and ethnic groups in the United States, research shows that many minority groups, especially African Americans and Latino Americans, seek and receive treatments for mental health problems at a much lower rate than the general population does, resulting in many unmet needs for mental health care and poorer mental health outcomes (Alegria et al., 2008; Cook et al., 2014; Fiscella, Franks, Doescher, & Saver, 2002; Jimenez et al., 2013; Lo, Cheng, & Howell, 2014; Neighbors et al., 2007; Obasi & Leong, 2009).

According to Aggarwal et al. (2016), numerous serious communication factors exist that lead to the reluctance of many minority members to seek treatment for their mental health concerns; this reluctance has led to serious disparities in mental health outcomes for members of minority communities in the United States. One of the biggest communication factors preventing treatment relates to social stigma concerning mental illness in these minority communities, which discourages individuals from disclosing concerns about mental health problems and from seeking mental health care (Clement et al., 2015; Corrigan, Druss, & Perlick, 2014; Robinson, 2012, 2013). Social stigma is a problem that reduces the perceived status of the people who are stigmatized, including those confronting mental health problems (Goffman, 1963). Stigma concerning mental health problems involves widespread pervasive beliefs portraying those who experience mental health problems as dangerous, evil, weak, or purposely uncooperative, beliefs that often lead to the avoidance and disparagement of the mentally ill in the United States (Robinson, 2012, 2013; Thoits, 2011).

Social stigma is a primary factor inhibiting discussion about mental health concerns and leads directly to reluctance to seek mental health treatment in many minority communities. This stigma discourages individuals from disclosing their concerns about mental health problems, assisting others facing mental health challenges, and seeking mental health treatment for themselves or others (Hatzenbuehler, Phelan, & Link, 2013;

Parcesepe & Cabassa, 2013; Pescosolido, Boyer, & Medina, 2013; Pescosolido, Medina, Martin, & Long, 2013; Robinson, 2012, 2013). However, because of the importance of communication in the prevention, diagnosis, and treatment of serious health problems, it is critical to address these complex communication barriers that derive from the social stigma related to mental health problems to help increase the use of effective communication for reducing mental health risk and incidence, and to improve outcomes from mental health care, especially within minority communities (Neumann et al., 2010; Turner, 2016). Access to relevant health information can enable individuals to make good decisions about mental health, identify their best strategies for improving mental health outcomes, and reduce uncertainty about helping others with mental health problems (Grohol, Slimowicz, & Granda, 2014; Kreps, 1988; Schomerus et al., 2016).

Communication, Stigma, and Mental Health Disparities

The serious health communication barriers that make discussing mental health concerns problematic have been closely related to growing disparities in mental health outcomes (Corrigan et al., 2014; Neely, 2015). Evidence shows that minority group members are less likely than other segments of society in the United States to disclose mental health diagnoses, receive adequate support from significant others about such diagnoses, seek care for mental health problems, and receive and continue mental health care treatments (Anglin, Alberti, Link, & Phelan, 2008; Corrigan et al., 2010; Robinson, 2012; Whaley, 2001; Williams, 2009). Lê Cook, Doksum, Chen, Carle, and Alegría (2013) indicate that the number of mental health care providers within minority communities is often limited, making access to high-quality mental health care difficult for residents of these communities. It is clear that there are numerous serious problems confronting members of minority communities in the United States that are likely to reduce access to mental health care services and treatments, so that mental health problems are exacerbated (Clement et al., 2015; Corrigan et al., 2014; Robinson, 2012, 2013).

In addition to the aforementioned barriers to receiving mental health care, research has shown that access to relevant and accurate information about mental health issues is often limited among the members of minority communities (Grohol et al., 2014; Obasi & Leong, 2009; Snowden, 2012). Furthermore, lack of information about available mental health services has been found to reduce the understanding of mental health problems that is needed to diagnose them accurately, seek the best treatments, and reduce the burden of mental health problems in minority communities (Grohol et

al., 2014; Obasi & Leong, 2009; Snowden, 2012). Even when the symptoms of mental health problems are detected by the individuals experiencing them, such people tend to self-treat their mental health disorders in ways that often make them worse (Lipsky, Kernic, Qiu, & Hasin, 2016; Lo, Tenorio, & Cheng, 2012; Turner, 2016). In many cases, members of minority groups choose self-treatment for mental health problems, typically with overuse of alcohol and illicit drugs (Lo et al., 2012; Turner, 2016). Although a self-medication approach may appear to these individuals to be an easier route than seeking professional treatments for coping with mental health problems, masking symptoms with substance abuse rarely solves their underlying mental health problems; instead, this common and dangerous strategy often leads to additional problems, such as addiction (Clement et al., 2015; Corrigan et al., 2014; Hatzenbuehler et al., 2013; Satcher, 2001).

Unfortunately, popular media, such as movies and television programs, have often promoted the stigma of mental health problems with the widespread dissemination of messages that help to establish and reinforce prejudice against those with mental health problems, portraying these sufferers as violent, dangerous, and out of control (Klin & Lemish, 2008; Neely, 2015; Parcesepe & Cabassa, 2013; Sirey, Franklin, McKenzie, Ghosh, & Raue, 2014; Stout, Villegas, & Jennings, 2004). Mental health stigma is rooted in feelings of shame, fear, embarrassment, discomfort, guilt, uncertainty, and lack of control (Corrigan et al., 2014; Klin & Lemish, 2008; Neely, 2015). The stigma-related discomfort associated with communicating about mental health issues makes it difficult for those who have such issues to acknowledge their problems and seek help in confronting them, even from members of their own families (Robinson, 2012, 2013). The groundbreaking research of Robinson (2012) on African Americans living with mental health problems suggests that their family members are often unreceptive to hearing about mental health problems and do not provide needed social support to those who have them. It is important to raise consciousness and awareness within minority communities that mental health issues are legitimate health concerns and that as significant health problems they require professional attention and treatment to be remedied.

Discomfort concerning communication about mental health issues is closely related to the stigmatization of those within minority communities who experience them; they are often categorized as crazy, dangerous, violent, hostile, and out of control (*loco*). In Latino communities, it is not uncommon for people with mental health problems to be perceived as possessed by the devil and in need of exorcism (Moodley & Palmer, 2014;

Sperry, 2015). People confronting mental health problems are often perceived as weak, undisciplined, and untrustworthy. Therefore, it is common for individuals with mental health problems to try to hide them, treat themselves for their mental health disorders, and refrain from seeking professional assistance. Lack of therapeutic communication about mental health issues inevitably leads to an increase in mental health problems and poor health outcomes for many members of minority communities.

Moreover, mental health problems are complex, often difficult to diagnose, and challenging to manage (Pescosolido, Boyer, & Medina, 2013). Serious mental health problems often require long-term treatments with a variety of therapies (Nathan & Gorman, 2015). Those who have mental health problems may not recognize their need for help, and others may interpret their symptoms as bad or antisocial behavior. The stigma surrounding mental health problems often leads to stereotyping and limited knowledge about the best ways to interact with and help persons with mental health problems. Inappropriate responses to those with mental health problems can often make the problems worse by frustrating, frightening, or provoking those who are experiencing them (Rogers & Pilgrim, 2014; Wood & Watson, 2016). It can be difficult to reason with people who have mental health problems. Training and educational programs can help increase understanding about the best strategies for recognizing, responding to, and providing assistance for those with mental health problems.

Suicide Prevention and Communication

The many stressors associated with living as a member of a racial or ethnic minority in the United States can intensify mental health problems (Cokley, McClain, Enciso, & Martinez, 2013; Sirin, Ryce, Gupta, & Rogers-Sirin, 2013). Prejudicial public stereotypes about competence, intelligence, responsibility, and intentions often lead to negative views and expectations of members of minority groups (Sirey et al., 2014; Sirin et al., 2013). Similarly, public assumptions about antisocial behaviors, including violence, substance abuse, and criminality, can increase the sensitivity of minority group members to public scrutiny and suspicion, leading to feelings of limited self-worth and a poor public image (Turner, 2016; Whaley, 2001; Williams & Williams-Morris, 2000; Williams, 2009). Living with the stress of prejudice and marginalization can promote depressive symptoms among members of minority groups that can increase the risk for suicide (Bostwick et al., 2014). Suicide has become a major mental health–related problem in the United States that is closely related to a lack of access to sensitive, caring, and supportive communication (Kleiman & Liu, 2013; Lai, Maniam,

Chan, & Ravindran, 2014; Silk, Perrault, Nazione, Pace, & Collins-Eaglin, 2017).

The communication problems encountered in addressing stress disorders and depression are especially problematic in efforts to reduce the risk for suicide in minority communities, where individuals can experience high levels of stress due to social bias and prejudice (Cokley et al., 2013; Sirin et al., 2013). It is difficult for members of minority communities to let others know about the stresses that are confronting them, and it is very difficult to diagnose and treat serious stress disorders. Suicidal tendencies can grow from unaddressed feelings of stress. Communication programs need to be developed to help recognize stress-related problems in minority communities; provide needed health services, such as counseling, therapy, and medications, to individuals confronting stress; and reduce the progression from stress disorders to depression and suicidal attempts (Kleiman & Liu, 2013; Lai et al., 2014; Silk et al., 2017).

Communication Interventions for Addressing Mental Health Problems

New communication strategies, programs, and policies must be proposed to help address the stigma of mental health problems in minority communities and to encourage open discussion about mental health issues, promoting the provision of needed interventions for effectively addressing these serious health problems. Strategic communication interventions have the potential to help address many of the inequities in the management of mental health problems that are prevalent in minority communities in the United States. For example, training for mental health care providers in communicating in culturally sensitive ways with members of minority groups who have mental health issues has been found to improve health outcomes (Copeland, 2006; Pearson et al., 2015). Similarly, culturally sensitive multimedia health education campaigns have been designed to help reduce the significant discomfort that members of many minority communities feel when discussing mental health issues (Corrigan et al., 2014; Silk et al., 2017; Thornicroft et al., 2016). Schomerus et al. (2016) found that an online internet-based educational intervention that provided information about viewing mental illness as a normal part of a continuum of mental health helped to combat the stigma of mental illness by influencing attitudes toward people with mental illness.

Ngo et al. (2016) found that community engagement programs can promote improvements in key mental health functional outcomes, reduce barriers to care, and increase participation in mental health care services. Ali, Farrer, Gulliver, & Griffiths (2015) found that the use of online peer-to-peer social support showed promise as a communication intervention for

addressing mental health problems. African American churches have been found to hold great potential for providing needed social support and treatment referrals for members experiencing mental health problems (Blank, Mahmood, Fox, & Guterbock, 2002; Bryant et al., 2014; Hankerson & Weissman, 2012). Bridges et al. (2014) found preliminary evidence indicating that the use of integrated mental health services, which seek to reduce stigma and barriers to service utilization by embedding mental health professionals into primary care teams, can help reduce mental health disparities for Latinos. In addition, programs that promote the expression of social support have been found to reduce risks for suicide by encouraging resilience, combatting loneliness, and promoting feelings of self-worth (Farrell, Bolland, & Cockerham, 2015; Silk et al., 2017).

Communication research can help identify the best strategies for reducing the stigmatization of mental health problems in minority communities. Basic research studies are needed to examine how key communication processes, such as relational coordination, information sharing, social influence, social support, and intercultural communication sensitivity, are related to providing help to individuals confronting mental health problems. We also need surveillance research programs to track how societal communication practices that stigmatize mental health issues can be reduced to promote a greater understanding of mental health issues and support for those living with mental health challenges. Most of all, we need to continue to develop and test evidence-based communication intervention programs for educating key audiences about mental health, as well as for training professional and lay caregivers to interact effectively with people seeking care for mental health concerns. There is great promise for improving mental health outcomes among members of minority communities if we can develop key communication strategies to reduce mental health stigma, increase access to relevant mental health information, and introduce communication programs that make high-quality mental health care easily available to all people who need such care. Increasing access, understanding, and the exchange of relevant mental health information can help to reduce disparities in mental health outcomes.

References

- Aggarwal, N. K., Pieh, M. C., Dixon, L., Guarnaccia, P., Alegría, M., & Lewis-Fernández, R. (2016). Clinician descriptions of communication strategies to improve treatment engagement by racial/ethnic minorities in mental health services: A systematic review. *Patient Education and Counseling, 99*(2), 198–209.
- Alegria, M., Chatterji, P., Wells, K., Cao, Z., Chen, C. N., Takeuchi, D., ... Meng, X. L. (2008). Disparity in depression treatment among racial and ethnic minority populations in the United States. *Psychiatric Services, 59*(11), 1264–1272.
- Ali, K., Farrer, L., Gulliver, A., & Griffiths, K. M. (2015). Online peer-to-peer support for young people with mental health problems: a systematic review. *JMIR Mental Health, 2*(2), e19.
- Allen, J., Balfour, R., Bell, R., & Marmot, M. (2014). Social determinants of mental health. *International Review of Psychiatry, 26*(4), 392–407.
- Anglin, D. M., Alberti, P. M., Link, B. G., & Phelan, J. C. (2008). Racial differences in beliefs about the effectiveness and necessity of mental health treatment. *American Journal of Community Psychology, 42*(1–2), 17–24.
- Blank, M. B., Mahmood, M., Fox, J. C., & Guterbock, T. (2002). Alternative mental health services: The role of the black church in the South. *American Journal of Public Health, 92*(10), 1668–1672.
- Bostwick, W. B., Meyer, I., Aranda, F., Russell, S., Hughes, T., Birkett, M., & Mustanski, B. (2014). Mental health and suicidality among racially/ethnically diverse sexual minority youths. *American Journal of Public Health, 104*(6), 1129–1136.
- Bridges, A. J., Andrews A. R. III, Villalobos, B. T., Pastrana, F. A., Cavell, T. A., & Gomez, D. (2014). Does integrated behavioral health care reduce mental health disparities for Latinos? Initial findings. *Journal of Latina/o Psychology, 2*(1), 37.
- Bryant, K., Haynes, T., Yeary, K., Hye-cheon, K., Greer-Williams, N., & Hartwig, M. (2014). A rural African American faith community's solutions to depression disparities. *Public Health Nursing, 31*(3), 262–271.
- Clement, S., Schauman, O., Graham, T., Maggioni, F., Evans-Lacko, S., Bezborodovs, N., ... Thornicroft, G. (2015). What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychological Medicine, 45*(01), 11–27.
- Cokley, K., McClain, S., Enciso, A., & Martinez, M. (2013). An examination of the impact of minority status stress and impostor feelings on the mental health of diverse ethnic minority college students. *Journal of Multicultural Counseling and Development, 41*(2), 82–95.

- Cook, B. L., Zuvekas, S. H., Carson, N., Wayne, G. F., Vesper, A., & McGuire, T. G. (2014). Assessing racial/ethnic disparities in treatment across episodes of mental health care. *Health Services Research, 49*(1), 206–229.
- Copeland, V. (2006). Disparities in mental health service utilization among low-income African American adolescents: Closing the gap by enhancing practitioner's competence. *Child and Adolescent Social Work Journal, 23*(4), 407–431. doi:10.1007/s10560-006-0061
- Corrigan, P. W., Druss, B. G., & Perlick, D. A. (2014). The impact of mental illness stigma on seeking and participating in mental health care. *Psychological Science in the Public Interest, 15*(2), 37–70.
- Corrigan, P. W., Morris, S., Larson, J., Rafacz, J., Wassel, A., Michaels, P., ... Rüsck, N. (2010). Self-stigma and coming out about one's mental illness. *Journal of Community Psychology, 38*(3), 259–275.
- Farrell, C. T., Bolland, J. M., & Cockerham, W. C. (2015). The role of social support and social context on the incidence of attempted suicide among adolescents living in extremely impoverished communities. *Journal of Adolescent Health, 56*(1), 59–65.
- Fiscella, K., Franks, P., Doescher, M. P., & Saver, B. G. (2002). Disparities in health care by race, ethnicity, and language among the insured: findings from a national sample. *Medical care, 40*(1), 52-59.
- Goffman, E. (1963). *Stigma: Notes on management of a spoiled identity*. Engelwood Cliffs, NJ: Prentice Hall.
- Grohol, J. M., Slimowicz, J., & Granda, R. (2014). The quality of mental health information commonly searched for on the Internet. *Cyberpsychology, Behavior, and Social Networking, 17*(4), 216–221.
- Hankerson, S.H., & Weissman, M. M. (2012). Church-based health programs for mental disorders among African Americans: A review. *Psychiatric Services, 63*(3), 243–249.
- Hatzenbuehler, M. L., Phelan, J. C., & Link, B. G. (2013). Stigma as a fundamental cause of population health inequalities. *American Journal of Public Health, 103*(5), 813–821.
- Jimenez, D. E., Cook, B., Bartels, S. J., & Alegría, M. (2013). Disparities in mental health service use of racial and ethnic minority elderly adults. *Journal of the American Geriatrics Society, 61*(1), 18–25.
- Kleiman, E. M., & Liu, R. T. (2013). Social support as a protective factor in suicide: Findings from two nationally representative samples. *Journal of Affective Disorders, 150*(2), 540–545.
- Klin, A., & Lemish, D. (2008). Mental disorders stigma in the media: Review of studies on production, content, and influences. *Journal of Health Communication, 13*(5),434–444.

- Kreps, G. L. (1988). The pervasive role of information in health and health care: Implications for health communication policy. *Annals of the International Communication Association*, 11(1), 2318–276.
doi:10.1080/23808985.1988. 11678988690
- Lai, M. H., Maniam, T., Chan, L. F., & Ravindran, A. V. (2014). Caught in the web: A review of web-based suicide prevention. *Journal of Medical Internet Research*, 16(1), e30.
- Lê Cook, B., Doksum, T., Chen, C. N., Carle, A., & Alegría, M. (2013). The role of provider supply and organization in reducing racial/ethnic disparities in mental health care in the US. *Social Science & Medicine*, 84, 102–109.
- Le Cook, B., Zuvekas, S. H., Carson, N., Wayne, G. F., Vesper, A., & McGuire, T. G. (2014). Assessing racial/ethnic disparities in treatment across episodes of mental health care. *Health Services Research*, 49(1), 206–229.
- Lipsky, S., Kernic, M. A., Qiu, Q., & Hasin, D. S. (2016). Posttraumatic stress disorder and alcohol misuse among women: effects of ethnic minority stressors. *Social Psychiatry and Psychiatric Epidemiology*, 51(3), 407–419.
- Lo, C., Cheng, T., & Howell, R. (2014). Access to and utilization of health services as pathway to racial disparities in serious mental illness. *Community Mental Health Journal*, 50(3), 251–257. doi:10.1007/s10597-013-9593-7
- Lo, C., Tenorio, K., & Cheng, T. (2012). Racial differences in co-occurring substance use and serious psychological distress: The roles of marriage and religiosity. *Substance Use & Misuse*, 47(6), 734–744. doi:10.3109/108226084.2012.666312
- Moodley, R., & Palmer, S. (2014). *Race, culture and psychotherapy: Critical perspectives in multicultural practice*. Abingdon, UK: Routledge.
- Nathan, P. E., & Gorman, J. M. (Eds.). (2015). *A guide to treatments that work*. New York, NY: Oxford University Press.
- Neely, S. Y. (2015). *Mental illness knowledge, stigma, help seeking behaviors, and perceptions of media portrayals of Black culture: The role of spirituality and the African American church* (Doctoral dissertation). University of Cincinnati, Cincinnati, Ohio.
- Neighbors, H. W., Caldwell, C., Williams, D. R., Nesse, R., Taylor, R. J., Bullard, K. M., ... Jackson, J. S. (2007). Race, ethnicity, and the use of services for mental disorders: Results from the National Survey of American Life. *Archives of General Psychiatry*, 64(4), 485.
- Neumann, M., Edelhäuser, F., Kreps, G. L., Scheffer, C., Lutz, G., Tauschel, D., & Visser, A. (2010). Can patient-provider interaction increase the effectiveness of medical treatments or even substitute them? A methodological reflection of the specific effect of the provider. *Patient Education and Counseling*, 80, 307–314.

- Ngo, V. K., Sherbourne, C., Chung, B., Tang, L., Wright, A. L., Whittington, Y., ... Miranda, J. (2016). Community engagement compared with technical assistance to disseminate depression care among low-income, minority women: A randomized controlled effectiveness study. *American Journal of Public Health, 106*(10), 1833–1841.
- Obasi, E. M., & Leong, F. T. L. (2009). Psychological distress, acculturation, and mental health-seeking attitudes among people of African descent in the United States: A preliminary investigation. *Journal of Counseling Psychology, 56*(2), 227–238. doi:10.1037/a0014865
- Parcesepe, A. M., & Cabassa, L. J. (2013). Public stigma of mental illness in the United States: A systematic literature review. *Administration and Policy in Mental Health and Mental Health Services Research, 40*(5), 384–399.
- Pearson, G. S., Hines-Martin, V. P., Evans, L. K., York, J. A., Kane, C. F., & Yearwood, E. L. (2015). Addressing gaps in mental health needs of diverse, at-risk, underserved, and disenfranchised populations: A call for nursing action. *Archives of Psychiatric Nursing, 29*(1), 14–18.
- Pescosolido, B. A., Boyer, C. A., & Medina, T. R. (2013). The social dynamics of responding to mental health problems. In C. Aneshensel, J. C. Phelan, & A. Bierman (Eds.), *Handbook of the sociology of mental health* (pp. 505–524). Dordrecht, the Netherlands: Springer.
- Pescosolido, B. A., Medina, T. R., Martin, J. K., & Long, J. S. (2013). The “backbone” of stigma: Identifying the global core of public prejudice associated with mental illness. *American Journal of Public Health, 103*(5), 853–860.
- Robinson, N. M. (2012). *To tell or not to tell: Factors in self-disclosing mental illness in our everyday relationships* (Doctoral dissertation). Fairfax, VA: George Mason University.
- Robinson, N. (2013). Addressing the stigma of mental illness: Promoting greater receptivity of mental health disclosures in co-cultural group members. In G. L. Kreps & S. Kodish (Eds.), *DC Health Communication (DCHC) Conference Proceedings* (p. 92). Fairfax, VA: Center for Health and Risk Communication.
- Rogers, A., & Pilgrim, D. (2014). *A sociology of mental health and illness*. London, UK: McGraw-Hill Education UK.
- Satcher, D. (2001). *Mental health: Culture, race, and ethnicity. A supplement to mental health: A report of the Surgeon General. 2001*. Washington, DC: US Department of Health and Human Services
- Schomerus, G., Angermeyer, M. C., Baumeister, S. E., Stolzenburg, S., Link, B. G., & Phelan, J. C. (2016). An online intervention using information on the mental health-mental illness continuum to reduce stigma. *European Psychiatry, 32*, 21–27.

- Silk, K. J., Perrault, E. K., Nazione, S. A., Pace, K., & Collins-Eaglin, J. (2017). Evaluation of a social norms approach to a suicide prevention campaign. *Journal of Health Communication, 22*(2), 1–8. Retrieved from <http://dx.doi.org/10.1080/10810730.2016.1258742>
- Sirey, J. A., Franklin, A. J., McKenzie, S., Ghosh, S., & Raue, P. (2014). Race, stigma and mental health recommendations among depressed older persons in aging services. *Psychiatric Services (Washington, D.C.), 65*(4), 537.
- Sirin, S. R., Ryce, P., Gupta, T., & Rogers-Sirin, L. (2013). The role of acculturative stress on mental health symptoms for immigrant adolescents: A longitudinal investigation. *Developmental Psychology, 49*(4), 736.
- Snowden, L. R. (2012). Health and mental health policies' role in better understanding and closing African American-White American disparities in treatment access and quality of care. *American Psychologist, 67*(7), 524–531. doi:10.1037/a0030054
- Sperry, L. (2015). Effective spiritually oriented psychotherapy practice is clinically, ethically, and culturally sensitive practice. *Spirituality in Clinical Practice, 2*(2), 103.
- Stout, P. A., Villegas, J., & Jennings, N. A. (2004). Images of mental illness in the media: Identifying gaps in the research. *Schizophrenia Bulletin, 30*(3), 543–561.
- Thoits, P. A. (2011). Resisting the stigma of mental illness. *Social Psychology Quarterly, 74*(1), 6–28.
- Thornicroft, G., Mehta, N., Clement, S., Evans-Lacko, S., Doherty, M., Rose, D., ... Henderson, C. (2016). Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *The Lancet, 387*(10023), 1123–1132.
- Turner, T. (2016). *Substance abuse and mental disorders among African Americans*. (Doctoral dissertation). Fort Lauderdale, FL: Kaplan University.
- Whaley, A. L. (2001). Cultural mistrust and mental health services for African Americans: A review and meta-analysis. *The Counseling Psychologist, 29*(4), 513–531.
- Williams, D. R., & Williams-Morris, R. (2000). Racism and mental health: The African American experience. *Ethnicity and Health, 5*(3/4), 243–268.
- Williams, T. M. (2009). *Black pain: It just looks like we're not hurting*. New York, NY: Simon & Schuster.
- Wood, J. D., & Watson, A. C. (2016). Improving police interventions during mental health-related encounters: Past, present and future. *Policing and Society, 1*–11.