

12-2011

## DYSREGULATION OF MEOX2 FOLLOWING WT1 MUTATION IN KIDNEY DEVELOPMENT AND WILMS TUMORIGENESIS

LaGina M. Nosavanh

LaGina Merie Nosavanh

Follow this and additional works at: [https://digitalcommons.library.tmc.edu/utgsbs\\_dissertations](https://digitalcommons.library.tmc.edu/utgsbs_dissertations)



Part of the [Medicine and Health Sciences Commons](#)

### Recommended Citation

Nosavanh, LaGina M. and Nosavanh, LaGina Merie, "DYSREGULATION OF MEOX2 FOLLOWING WT1 MUTATION IN KIDNEY DEVELOPMENT AND WILMS TUMORIGENESIS" (2011). *The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences Dissertations and Theses (Open Access)*. 203.

[https://digitalcommons.library.tmc.edu/utgsbs\\_dissertations/203](https://digitalcommons.library.tmc.edu/utgsbs_dissertations/203)

This Thesis (MS) is brought to you for free and open access by the The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences at DigitalCommons@TMC. It has been accepted for inclusion in The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences Dissertations and Theses (Open Access) by an authorized administrator of DigitalCommons@TMC. For more information, please contact [digitalcommons@library.tmc.edu](mailto:digitalcommons@library.tmc.edu).

DYSREGULATION OF *MEOX2* FOLLOWING *WT1* MUTATION IN KIDNEY  
DEVELOPMENT AND WILMS TUMORIGENESIS

by

LaGina Merie Nosavanh, B.S.

APPROVED:

---

[Advisor, Vicki Huff, Ph.D.]

---

[Michelle Barton, Ph.D.]

---

[Joseph Alcorn, Ph.D.]

---

[Mong-Hong Lee, Ph.D.]

---

[Thomas Goka, Ph.D.]

APPROVED:

---

Dean, The University of Texas  
Graduate School of Biomedical Sciences at Houston

**DYSREGULATION OF *MEOX2* FOLLOWING *WT1* MUTATION IN KIDNEY  
DEVELOPMENT AND WILMS TUMORIGENESIS**

A

THESIS

Presented to the Faculty of  
The University of Texas  
Health Science Center at Houston  
and  
The University of Texas  
M. D. Anderson Cancer Center  
Graduate School of Biomedical Sciences

in Partial Fulfillment  
of the Requirements  
for the Degree of  
MASTER OF SCIENCE

by

LaGina Merie Nosavanh, B.S.

Houston, Texas

December 2011

## **Dedication**

To my mother, Sylvia Ann Dozier-Rayne, for the love and support she provided me throughout my entire life. She quoted, “Knowledge is power”, and to always do my best in what I do in my life. She inspired me to pursue the highest degree I can obtain and live life to the fullest. Furthermore, I dedicate my thesis to my father, Boualone Nosavanh, my grandmother, Christine Taylor, and to my siblings, Barry, Jessica, Joseph, Margie and Michael. They have provided me with the utmost love and encouragement. My younger siblings tell me that they look up to me as their role model, which makes me feel as if I have accomplished more than I ever imagined. To my friends, I am so fortunate to have you as a part of my life and express great appreciation for your support, encouragement, and positive energy throughout my graduate career.

## **Acknowledgments**

I would like to express my deepest gratitude to Dr. Fatma Helmy, who provided me an exciting start into my passion of biomedical research. She is the director of the MARC program at Delaware State University and was actively involved in my career choices. My first experience in an actual laboratory was stimulated by the MARC program in Dr. Jeffery A. Frost lab at the University of Texas – Health Science Center at Houston. My passion for research intensified after my first research experience. These two individuals had a significant impact on my maturity as a scientist.

I would like to acknowledge Dr. Vicki Huff for this amazing and exciting project. She provided me with scientific mentorship and the utmost encouragement in the accomplishment of my project. I would like to thank the members of my advisory, supervisory, and examining committees, Dr. Pierre McCrea, Dr. Michael Galko, Dr. Timothy McDonnell, Dr. Kevin Coombes, Dr. Michelle Barton, Dr. Joseph Alcorn, Dr. Mong-Hong Lee, Dr. Thomas Goka, Dr. Stephanie Watowich, and Dr. Elsa Flores for their involvement in my graduate career. I want to give a special thanks to Dr. Joseph Alcorn and Dr. Mong-Hong Lee, because they believed in me throughout my entire candidacy exam and gave me the confidence and strength to complete my Masters degree.

My special thanks to the members of the Huff lab. Dr. Sharada Mokkalapati and Dr. Paramahansa Maturu who constantly shared their knowledge about molecular biology, genetics, cancer biology/genetics, and life in general. My fellow graduate student, Le Huang, has provided support, feedback, and an amusing environment in the lab every day – it has been a delight to have her as a colleague and a wonderful friend.

I would like to acknowledge my funding source, Research Supplements to Promote Diversity in Health-Related Research (5 P01 CA349936-22).

## DYSREGULATION OF *MEOX2* FOLLOWING *WT1* MUTATION IN KIDNEY DEVELOPMENT AND WILMS TUMORIGENESIS

Publication No. \_\_\_\_\_

LaGina Merie Nosavanh, B.S.

Supervisory Professor: Vicki Huff, Ph.D.

Wilms tumor (WT) is a childhood tumor of the kidney and a productive model for understanding the role of genetic alteration and interactions in tumorigenesis. The Wilms tumor gene 1 (*WT1*) is a transcriptional factor and one of the few genes known to have genetic alterations in WT and has been shown to be inactivated in 20% of WTs. However, the mechanisms of how *WT1* mutations lead to Wilms tumorigenesis and its influence on downstream genes are unknown. Since it has been established that *WT1* is a transcriptional regulator, it has been hypothesized that the loss of *WT1* leads to the dysregulation of downstream genes, in turn result in the formation of WTs. To identify the dysregulated downstream genes following *WT1* mutations, an Affymetrix GeneChip Human Genome Array was previously conducted to assess the differentially expressed genes in the *WT1*-wildtype human and *WT1*-mutant human WTs. Approximately 700 genes were identified as being significantly dysregulated. These genes were further prioritized based on their statistical significance, fold change, chromosomal region, spatial pattern of gene expression and known or putative cellular functions. Mesenchyme homeobox 2 (*MEOX2*) was one of the most significantly upregulated genes in *WT1*-mutant WT. *MEOX2* is known to play a role in cell proliferation, apoptosis, and differentiation. In addition to its biological roles, it is expressed during early kidney development in the

condensed mesenchyme similar to *WT1*. Furthermore, the use of the Match® web-based tool from the BIOBASE Biological Data base identified a significant predicted *WT1* binding site within the first intron of *MEOX2*. The similarity in spatial gene expression in the developing kidney and the significant predicted *WT1* binding site found in the first intron of *MEOX2* lead to the development of my hypothesis that *MEOX2* is upregulated via a *WT1*-dependent manner.

Here as a part of my master's work, I have validated the Affymetrix GeneChip Human Genome Array data using an independent set of Wilms tumors. *MEOX2* remained upregulated in the mutant *WT1* Wilms tumor by 41-fold. *Wt1* and *Meox2* gene expression were assessed in murine newborn kidney; both *Wt1* and *Meox2* were expressed in the condensed, undifferentiated metanephric mesenchyme. I have shown that the *in vivo* ablation of *Wt1* during embryonic development at embryonic day (E) 13.5 resulted in the slight increase of *Meox2* gene expression by two fold. In order to functionally demonstrate the effect of the loss of *Wt1* on *Meox2* gene expression in undifferentiated metanephric mesenchyme, I have generated a kidney mesenchymal cell line to genetically ablate *Wt1* *in vitro* by adenoviral infection. The ablation of *Wt1* in the kidney mesenchymal cell line resulted in the upregulation of *Meox2* by 61-fold. Moreover, the upregulation of *Meox2* resulted in the significant induction of *p21* and *Itgb5*. In addition to the dysregulation of these genes the ablation of *Wt1* in the kidney mesenchymal cells resulted in decrease in cell growth and loss of cellular adherence. However, it is uncertain whether the upregulation of *Meox2* caused this particular cellular phenotype. Overall, I have demonstrated that the upregulation of *Meox2* is *Wt1*-dependent during early kidney development.

## Table of Contents

Approval Sheet.....	i
Title page.....	ii
Dedication.....	iii
Acknowledgements.....	iv
Abstract.....	v
Table of Contents.....	vii
List of Illustrations .....	ix
List of Tables.....	xi
Abbreviations.....	xii
Chapter 1 – Introduction.....	1
I. Wilms Tumor.....	1
II. Kidney Development.....	2
III. Genetic Alterations in Wilms Tumor.....	6
IV. Wilms Tumor Suppressor Gene 1.....	8
a. <i>Wt1</i> in Kidney Development.....	10
b. <i>WT1</i> in Wilms Tumor.....	12
c. Genes Regulated by WT1.....	14
V. Rationale.....	15
Chapter 2 – Upregulation of <i>MEOX2</i> in <i>WT1</i> -mutant tumors:	
Validation of WT1 Putative Transcriptional Targets by qPCR.....	17
I. Material and Methods.....	22
II. Results.....	25
III. Discussion.....	27



<b>Chapter 3 – Generation of a Kidney Mesenchymal Cell Line.....</b>	<b>30</b>
<b>I. Material and Methods.....</b>	<b>31</b>
<b>II. Results .....</b>	<b>39</b>
<b>III. Discussion.....</b>	<b>48</b>
<b>Chapter 4 – Mutant Wt1 Regulates the Gene Expression of <i>Meox2</i>.....</b>	<b>49</b>
<b>I. Material and Methods.....</b>	<b>54</b>
<b>II. Results .....</b>	<b>57</b>
<b>III. Discussion.....</b>	<b>67</b>
<b>Chapter 5 – Conclusions and Future Directions.....</b>	<b>69</b>
<b>I. Conclusions.....</b>	<b>69</b>
<b>II. Future Directions.....</b>	<b>71</b>
<b>Appendix.....</b>	<b>74</b>
<b>Bibliography.....</b>	<b>93</b>
<b>Vitae.....</b>	<b>115</b>

## List of Illustrations

**Figure 1.1** Kidney Development

**Figure 1.2** Hematoxylin and Eosin Staining of Newborn Mouse Kidney and WTs

**Figure 1.3** WT1 Gene and Protein

**Figure 1.4** Wt1 Protein Expression during Murine Kidney Development

**Figure 1.5** WT1 mutations detected in WTs

**Figure 2.1** Validation of the Dysregulated Genes in the *WT1*-mutant WTs

**Figure 3.1.** Schematic Diagram of the Cre-Lox System

**Figure 3.2** PCR Analyses of the MMCs using *Wt1-F/PGK*, *1.55/1.75*, *ZP3 se/ZP3 as*, and *Immo I/Immo II* primer sets

**Figure 3.3** Ablation of *Wt1* by Tamoxifen Treatment

**Figure 3.4** Viral Infection of Adeno-Cre-GFP in the F12055 *Wt1*<sup>-flox</sup>; *Immorto* MMC cell line

**Figure 3.5** PCR Analyses of the Viral Ablation of *Wt1* in the F12055 *Wt1*<sup>-flox</sup>; *Immorto* MMC cell line

**Figure 3.6** IF Images of the Generated MMCs from E13.5 Kidney Rudiments

**Figure 3.7** Biological Marker for Condensed Cap Mesenchyme

**Figure 4.1** Representation of Meox2 Gene and Protein Sequence

**Figure 4.2** Meox2 Gene Expression in E14.5 Mouse

**Figure 4.3** PCR Analyses of the E13.5 Kidney Rudiments after the *in vivo* Ablation of *Wt1*

**Figure 4.4** Meox2 Gene Expression after the *in vivo* Ablation of *Wt1*

**Figure 4.5** Meox2 is Upregulated following the *in vitro* Ablation of *Wt1*

**Figure 4.6** The Effect of the Loss of *Wt1* on *Tgf-β1* and *Smad3* Gene Expression

**Figure 4.7** Induction of *p21* and the Repression of *Itgb5*

**Figure 4.8** The Effect of *Wt1* Ablation in *Wt1*<sup>-flox</sup>; Immorto MMCs: Visual Observation in Decrease Cell Number and Loss of Cell Adhesion

**Figure 4.9** The effect of *Wt1* Ablation in *Wt1*<sup>-flox</sup>; *Immorto* MMCs: Cell Number and Cell Adhesion

**Appendix Figure 1** IF Images of the Subcloned Kidney MMCs

**Appendix Figure 2** IF images of the MMCs with the *CreER*<sup>TM</sup> Allele

## List of Tables

**Table 2.1.** Selected Differentially Expressed Genes from the Affymetrix GeneChip Array

**Table 2.2.** Biological and Cellular Description of the Selected Genes

**Table 2.3.** Primer Sets designed by Primer Express v.3 Software for the qPCR Validation

**Table 3.1.** PCR Conditions for Animal Genotyping

**Table 3.2** Primer Sequences for Animal Genotyping

**Table 3.3** MMCs Cell Lines Derived from E13.5 Kidney Rudiments

**Table 4.1.** Primer Sets designed by Primer Express v.3 Software for the qPCR analysis using cDNA from E13.5 kidney and MMCs

**Appendix Table 1** Putative Targets of WT1

**Appendix Table 2.** Calculating the Relative Gene Expression of Validated Genes (n=12)

**Appendix Table 3.** Calculating the relative gene expression of *the Wt1*, *Tgf- $\beta$ 1*, *Smad3*, *Meox2*, *p21* and *Itgb5* after adenoviral infection

## **Abbreviations**

$\Delta$ , Ablated

$\Delta$ Ct, Delta Ct

4-OH-Tamoxifen, 4-hydroxyl-Tamoxifen

A, Adenine

A1, Allele 1

ABC, Avidin-biotin-peroxidase

BLAST, Basic local alignment search tool

BSA, Bovine serum albumin

C, Carboxyl

Cy, Cytosine

CGH, Comparative genomic hybridization

ChIP, Chromatin immunoprecipitation

Chr., Chromosome

Cited1, Cbp/p300-interacting transactivator 1

CKO, Conditional knockout

CMV, cytomegalovirus

CXXC4, CXXC-type zinc finger protein 4

DDR1, Discoidin domain receptor family, member 1

DDS, Denys-Drash syndrome

Del, Deletion

dH<sub>2</sub>O, distilled water

DIC, Differential interference contrast

DMEM, Dulbeccos's modified Eagle's medium

E, Embryonic day

eGFP, enhanced green fluorescent protein

Ex, Exon

FBS, Fetal bovine serum

FDR, False Discovery Rate

FS, Frame Shift

FST, Follistatin

FWT1/FWT2, Familial predisposition genes of WT on chromosome regions 17q12-q21 and 19q13.4

G, Guanine

GAPDH, Glyceraldehyde phosphate dehydrogenase

GFP-FITC, GFP fluorescein isothiocyanate

H19, Imprinted maternally expressed transcript (non-protein coding) H19 allele

HIPK2, Homeodomain-interacting protein kinase 2

IF, Immunofluorescence

IFN- $\gamma$ , Interferon gamma

IGF2, Insulin-like growth factor

IHC, Immunohistochemistry

Ins, Insertion

Itgb5, Integrin Beta 5

IRES, Internal ribosome entry site

KTS, Lysine, threonine, and serine

LOH, Loss of heterozygosity

LOI, Loss of imprinting

MEOX2, Mesenchyme homeobox gene 2

MET, Mesenchymal-to-epithelial transition

MLLT3, Myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, *Drosophila*);  
translocated to, 3

MMC, Metanephric mesenchymal cells

MOI, Multiplicity of infection

N, Amino

NF- $\kappa$ -B, Nuclear Factor  $\kappa$ -B

NLS, Nuclear localization signaling

NTC, Non-treated cells

P1, Post natal day 1

PAX6, Paired box gene 6

PBS, phosphate buffered saline

PCR, Polymerase chain reaction

PRRX1, Paired related homeobox 1

qPCR, Quantitative PCR

rpm, Revolutions per minute

SD, Standard deviation

T, Thymine

TA<sub>g</sub>, T-Antigen

Term, Termination of transcription

TGF- $\beta$ , Transforming growth factor  $\beta$

VSMCs, Vascular smooth muscle cells

WAGR, Wilms tumor, aniridia, genitourinary malformation, and mental retardation

Wilms tumor, WT

WT1, Wilms tumor suppressor gene 1

WTX, Wilms tumor on the X located at Xq11.1

X, Fold change



## **Chapter 1: Introduction**

### **Wilms Tumor**


Wilms tumor (WT) is an embryonal malignancy of the kidney that represents approximately 95% of all pediatric kidney tumors and accounts for 6% of all pediatric cancers (1-4). The overall annual incidence of Wilms tumor in North America is 1 in 10,000 children, resulting in 600-700 new cases each year (3, 4). Approximately 75% of these WTs occur in children under the age of 5 years, with a peak occurrence between 3-4 years of age (5). WTs can be classified into three clinical categories based on its occurrence: 1) the sporadic category, which describes children with no unusual physical features or family history of WT; 2) the familial category, which describes children with family history of WT; and 3) the syndromic category, which describes children with congenital anomalies, such as Wilms tumor, aniridia, genitourinary malformation, and mental retardation (WAGR) syndrome, Denys-Drash syndrome (DDS), and Beckwith–Wiedemann syndrome (BWS). Between 98-99% WT cases are sporadic, while only 1-2% are familial (1, 3, 6). The majority of WT cases (90-95%) are unilateral (occurring in one kidney). However, bilateral cases (occurring in both kidneys) are also reported, particularly in patients with congenital anomalies as well as a family history of WTs (1, 7). Because WT is thought to occur during early kidney development, it serves as an excellent model to study the relationship of cancer to development by understanding normal kidney development and how a disruption during this process results in Wilms tumorigenesis.

## Kidney Development

The mammalian kidney arises during embryogenesis from the interaction between the differentiated epithelial cells of the ureteric bud and undifferentiated mesenchymal cells (8, 9). In mice, the ureteric bud epithelium invades the uncondensed, undifferentiated metanephric mesenchyme, which causes the metanephric mesenchymal cells to condense around the ureteric bud (**Figure 1.1 A-C**). Reciprocal interactions occur between the ureteric bud and the metanephric mesenchyme. The ureteric bud induces the metanephric mesenchyme to simultaneously give rise to stromal mesenchyme, which is rich in growth factors and corresponding receptors for the adjacent epithelial ureteric bud. Concurrently, the metanephric mesenchyme further condenses around the ureteric bud to undergo a mesenchymal-to-epithelial transition (MET) (8-11). Simultaneously, the induced mesenchyme forms into the comma and S-shaped bodies as the ureteric bud continues to branch out and proliferate (**Figure 1.1 D-E**) (8-10). The S- and comma-shaped bodies of the mesenchyme completely differentiates into epithelial cells to make up the majority of the cells of the nephron, such as the tubules and glomeruli (**Figure 1.1 F-G**) (8-10). The extensive branching of the ureteric bud results in the generation of the collecting ducts of the mature kidney, and the kidney continues to develop into the mature kidney after birth in mice (8-10). The stages of kidney development in humans occur similar to that in mice.


WT is thought to arise from undifferentiated metanephric mesenchyme. The classic histology of WTs, in both humans and mice, is triphasic consisting of mesenchymal, stromal, and epithelial cells that suggest disruption(s) during the cellular differentiation in the cells of the maturing nephron (**Figure 1.2**). Some

individual WTs consist of two of the three cell types; one cell type can be predominant over the other cell types present in the tumor. Because of the classic triphasic histology of WT, it is thought that there was a disruption during early kidney development, particularly in the cellular differentiation. This disruption may inhibit the cells to differentiate into the components of the nephron and cause them to overproliferate, in turn resulting in WT.



**Figure 1.1. Kidney Development**

Kidney development in mice starts at embryonic day (E) E10.5-11.5. In the beginning, the uncondensed mesenchyme **(A)** is invaded by the ureteric bud **(B)**. At E12.5, the mesenchyme becomes condensed **(C)**. At E13.5, comma-shaped bodies and S-shaped bodies **(D and E)** start to appear during kidney development. At E14.5, the tubular component of the nephron continues to develop into the functionally distinct cells of the proximal convoluted tubule, descending and ascending tubule **(F)** and the mesenchymal cells transition to become epithelized and differentiate into specialized cells of the glomerulus and tubules of the mature nephron **(G)**. In humans, kidney development occurs similar to that in mice (10, 12). (Adapted from (12))



**Figure 1.2. Hematoxylin and Eosin Staining of Newborn Mouse Kidney and WTs**

The triphasic histology of WTs in both human and mouse illustrates the early stages of kidney development (B- Blastemal, S- Stromal, and E- Epithelial cells). (Provided by Gao, F. and Hu, Q.)

## Genetic Alterations in Wilms Tumor

WT was the second tumor to meet the criteria of the Knudson's two hit model of cancer, in which there are two rate-limiting events, or 'hits', in turn resulting in cancer formation (13). The two rate-limiting events are a first 'hit', (which can be either a germline or somatic mutation), which may be followed by a second 'hit', always a somatic mutation (13). Besides the two rate-limiting events of the classic Knudson's two hit model, it has been suggested that there could be multiple genetic alterations and/or multiple mutated genes involved in an individual WT. Cytogenetic, genetic linkage, loss of heterozygosity (LOH), loss of imprinting (LOI), and DNA sequencing studies provided data that identified alterations that occur during Wilms tumorigenesis.

Cytogenetic studies, such as G-Banding, fluorescent in situ hybridization (FISH) and comparative genomic hybridization (CGH), revealed chromosomal alterations, including the gain and loss of chromosomes and chromosomal regions in WTs. The chromosomal gains in WTs were observed in chromosomes 6, 7, 12, 13, 17, 18, and 20 and the chromosomal losses noted in chromosomal regions 1p, 2q, 3, 7p, 11p, 11q, 14q, 16q, 17p, and 22q (14-17). These chromosomal regions may harbor genes that contribute to Wilms tumorigenesis.

As of now, genetic linkage studies mapped chromosomal region 17q12-q21 as familial Wilms Tumor gene 1 (*FWT1*) and chromosomal region 19q13.4 as familial Wilms tumor gene 2 (*FWT2*) as familial predisposition regions in large WT families. These regions may contain genes that predispose to WT (18, 19). However, no genes have yet been identified within these regions that contribute to the familial

predisposition of WT. The mapping of additional chromosomal regions in WT families is still an ongoing process.

Additional to cytogenetic analyses, LOH studies identified the loss of chromosomal region 11p in tumors versus constitutional DNA from the same individual. The demonstration of LOH at loci 11p in ~40% of WTs suggested that the inactivation of the gene within this particular region is essential to Wilms tumorigenesis (20-22). In addition to the LOH of 11p chromosomal region, WAGR patients carried constitutional deletions encompassing band 11p13 (17). Kaneko *et al.* reported that subsequent observation of 11p13 deletions in tumor DNA from patients who did not carry germline 11p13 deletions indicated somatic mutations were also essential during tumorigenesis (23). Collectively, these analyses of 11p13 deletions led investigators to localize and clone the first WT gene, Wilms tumor gene 1 (*WT1*) (24). This particular gene is mutated in ~20% of WTs (1).


DNA sequencing analysis data identified additional genetic alterations in identifiable genes in WTs. The tumor suppressor protein p53 (*TP53*), located at chromosome region 17p13, is mutated in 5% of WTs (25, 26). The tumors that carry *p53* mutations contain abnormal cells that are poorly developed with enlarged distorted nuclei (25, 26). Germline and somatic mutations of *TP53* include either missense mutations that occur in the exons 5, 6, and 10, and lead to the inactivation of the protein (25). Catenin (cadherin-associated protein), beta 1 (*CTNNB1*), located at chromosomal region 3p21, is mutated in 15% of sporadic WTs (27, 28). Somatic mutations of *CTNNB1* include protein-stabilizing mutations in exon 3 (27, 28). Additional mutations occur in exons 7 and 8 of *CTNNB1* (29, 30). Recently, Wilms tumor gene located at Xq11.1 (*WTX*) was discovered to be mutated in ~20% of WTs

(31-34). Somatic mutations of *WTX* include either whole or partial deletion of *WTX* or nonsense mutations resulting in termination codons (31-34). The discovery of all the genetic alterations and identifiable mutated genes involved during Wilms tumorigenesis is still in progress. *WT1* is one of the genes that is commonly mutated in WTs and essential during kidney development as later described. It is important to further study the loss of *WT1* in order to determine and establish its fundamental role during Wilms tumorigenesis.

### **Wilms Tumor Suppressor Gene 1 (*WT1*)**

Located at 11p13 in humans, the *WT1* gene consists of ten exons and encodes a zinc finger transcriptional factor which contains a proline glutamine-rich protein-interaction domain and four carboxy-terminal zinc finger domains (24, 35) (**Figure 1.3A**). Four isoforms of the WT1 protein result from two different alternative splice sites. Alternative splice I inserts the entire exon 5, which is made up of 51 nucleotides and encodes 17 amino acids (**Figure 1.3B**) (36). Alternative splice II inserts 9 nucleotides in exon 9 that encodes an additional three amino acids, lysine [K], threonine [T], and serine [S] (**Figure 1.3B**) (36).





**Figure 1.3. WT1 Gene and Protein**


**1.3A.** Exons are indicated by yellow and the alternative splice sites are located within exons 5 and 9 that result in four different isoforms of the WT1 protein (36).

**1.3B.** Proline and glutamine rich domain (blue) and four zinc fingers that serve as the DNA binding domain (green) of the WT1 protein (36). (Adapted from (7))

### Wt1 in Kidney Development

At E9.5 in mice, the first expression of *Wt1* is in the intermediate mesenchyme lateral to the coelomic cavity (37). *Wt1* is later expressed in the metanephric mesenchyme (uncondensed and condensed) (**Figure 1.4**) and its maximum expression is in the S- and comma-shaped bodies (9, 38-42). In the adult kidney, it is restricted to the podocytes of the glomerulus (**Figure 1.4**) (10, 42). The essential role of *Wt1* during kidney development was not recognized until a mouse model was generated for the purpose of investigating the role of *Wt1* in development, homeostasis, and WT development.

Kreidberg *et. al.* (1993) first generated a *Wt1* mouse model (43). Heterozygous mutant *Wt1* mice appeared to be normal; however homozygous mutant *Wt1* mice were typically embryonic lethal between E13 and E15 (43). Homozygous mutant *Wt1* mice failed to complete gestation due to systemic edema and pericardial bleeding (43). Besides embryonic lethality, the major consequence of *Wt1* mutation was renal agenesis (43). Hu *et. al.* (2011) also demonstrated that the *in vivo* ablation of *Wt1* in a conditional inducible knock out mouse model resulted in a complete block in nephron development, in which no glomeruli were present, and there was no differentiation of condensed mesenchyme past the comma-shaped body stage (44). All together, *Wt1* has an essential role, direct or indirect, in the survival and differentiation of the metanephric mesenchymal cells during kidney development.




**Figure 1.4. Wt1 Protein Expression during Murine Kidney Development**

Wt1 is expressed in the uncondensed and condensed mesenchyme (**left**) and expression remains high throughout kidney development, especially in the S- and comma-shaped bodies (**middle**). Its expression is restricted to the podocytes of the glomerulus (**right**) (45). (Provided by Gao, F. and Hu, Q.).

### WT1 in Wilms Tumor

Germline and sporadic mutations of *WT1* in WTs consist of alterations such as deletions or insertions leading to frameshift and/or truncation, missense and nonsense mutation, or splice site mutations that result in the inactivation of the WT1 protein (**Figure 1.5**) (1, 32, 46). These mutations lead to the inactivation of the protein (1, 32). *WT1* germline mutations occur in children with congenital anomaly syndromes, such as WAGR and DDS, and have a high risk of developing WTs (1, 32). In regards to *WT1* mutations in sporadic WTs, numerous of studies have gathered data providing different frequencies of *WT1* mutations ranging from 10-20% (47-50). Furthermore, patients with homozygous mutations within *WT1* provided evidence that mutations in both alleles at the *WT1* locus are critical in WT development and define the functional role of *WT1* in WT as a tumor suppressor gene (1, 47). Even though genetic studies identified germline and somatic mutations that result in the inactivation of *WT1* in WTs, the effect of the loss of *WT1* during Wilms tumorigenesis remains unknown. Since WT1 is a transcriptional factor that regulates the activation or repression of genes, the driving hypothesis is that the inactivation of *WT1* results in the dysregulation of downstream genes that are involved in key cellular pathways during embryogenesis that contribute to Wilms tumorigenesis.



**Figure 1.5. *WT1* mutations detected in WTs**

Germline mutations, dispersed throughout the entire *WT1* gene, detected in patients with Wilms' tumor-associated phenotypes. Similar mutations were detected in somatically mutated WTs. Location of gene mutations are designated by red (insertion or deletions and frameshift mutations), blue (nonsense mutations), and green (splice site mutations). (Adapted from (51)).

## Genes Regulated by WT1

Genes regulated by WT1 have been identified and are involved in pathways that are essential in cellular proliferation, apoptosis, cell survival, and cellular differentiation. The identification of the majority of genes regulated by WT1 was done by promoter assays, electrophoretic mobility shift assay (EMSA), co-immunoprecipitation (co-IP), microarray analyses, and co-transfection assays (52-124). These *in vitro* assays were conducted with human cell lines derived from normal and cancerous tissues and/or murine cell lines (52-124). A collection of published genes thought to be regulated by WT1 is displayed in **Appendix Table 1**. However, it is unclear whether the inactivation of *WT1* has an influence on their gene expression during Wilms tumorigenesis. To determine the effect of the loss of *WT1* on the downstream genes, a previous Affymetrix GeneChip Human Genome Array was used as an approach to identify genes that are differentially expressed in human *WT1*-mutant WTs when compared to *WT1*-wildtype WTs. Approximately 700 genes were significantly differently expressed from this array (Ruteshouser, E. C.). These genes were prioritized by their False Discovery Rate (FDR, 0.05), statistical significance (p-Value <0.05), fold change (>2X), chromosomal region, and gene expression pattern. Moreover, extensive literature searches on their known or putative cellular functions narrowed the number of genes for further investigation to 223 genes. The effect of the loss of *WT1* resulted in the differential expression of a variety of genes; however, it remains unknown whether the effect of the inactivation of *WT1* dysregulates their gene expression during early kidney development, which could lead to the development of WTs.

## Rationale

WT1 is known to be both an essential gene during kidney development and a tumor suppressor of WT. *WT1* is reported mutated in WTs, however the effect of the loss of *WT1* during Wilms tumorigenesis remains uncertain. Potentially, WT1 transcriptionally regulates genes involved in signaling pathways that control cellular processes, such as cellular proliferation, differentiation, apoptosis, and cell adhesion. The hypothesis of the inactivation of *WT1* in WT is that it results in the transcriptional dysregulation of downstream genes that lead to Wilms tumorigenesis. To test the hypothesis, a microarray was conducted and identified numerous genes that were either upregulated or downregulated in *WT1*-mutant WTs. The microarray data support the hypothesis that the inactivation of *WT1* transcriptionally dysregulates downstream genes. I have propose to investigate the loss of *Wt1* during murine kidney development to determine whether the gene is expressed during early kidney development, whether *Wt1* regulates, directly or indirectly, the gene during kidney development, and whether their dysregulation in gene expression following a *Wt1* mutation lead to the formation of WT.

I have taken the first step to determine the effect of inactivation of *WT1* on the genes dysregulated in WTs by validating the microarray data by quantitative PCR (qPCR) using an independent set of WTs to rule out any false discoveries from the microarray data (**Chapter 2**). The qPCR validation revealed a stronger gene regulation relationship between *WT1* and *MEOX2*. Therefore, the goal of my project is to further assess the effect *MEOX2* gene expression following a *WT1* mutation during kidney development.

Because it is thought that a disruption in the cellular differentiation process during early kidney development leads to Wilms tumorigenesis, I generated a number of conditional *Wt1* immortalized undifferentiated metanephric mesenchymal cell lines (MMCs) (**Chapter 3**). I used this cell line to determine the expression level of *Meox2* before and after the ablation of *Wt1*. To assess the dysregulation of *Meox2* following a *Wt1* mutation during early kidney development *in vivo*, I used the *Wt1* conditional knockout mouse model to ablate *Wt1* at E11.5 and determined the dysregulation of *Meox2* at E13.5 (Chapter 4). Results from both the *in vivo* and *in vitro* studies indicated that the inactivation of *Wt1* results in the upregulation of *Meox2* during early kidney development. Furthermore, the upregulation of *Meox2* from the *in vitro* analyses showed an effect on *Meox2* established downstream genes, *p21* and *Itgb5*. My studies of the effect of gene expression of *Meox2* following a *Wt1* mutation suggest that *Wt1* regulates *Meox2* during both early kidney development and Wilms tumorigenesis.



## **Chapter 2: Upregulation of *MEOX2* in *WT1*-mutant tumors: Validation of *WT1* Putative Transcriptional Targets by qPCR**

*WT1* is expressed during early kidney development and encodes a transcriptional factor and tumor suppressor that recognizes several DNA motifs such as the early growth response-1 (EGR-1) binding site, TCC repeats, and a high affinity WTE site (55, 71, 101, 125, 126). As displayed in **Appendix Table 1**, *WT1* putatively activates or represses its downstream genes. Thus, the effect of the loss of *WT1* during Wilms tumorigenesis could be the increased or decreased expression of downstream genes during early kidney development. Our recent microarray data identified genes that were dysregulated due to the inactivation of *WT1* in WT<sub>s</sub>. Some of these dysregulated genes are known to be normally expressed in the kidney, while others were not.

Our previous Affymetrix GeneChip array data revealed numerous genes that were significantly dysregulated in *WT1*-mutant WT<sub>s</sub>. As previously mentioned in **Chapter 1**, stringent statistical analysis of the microarray data and literature searches further narrowed down the number of genes to 21 as the most significant dysregulated genes in *WT1*-mutant WT<sub>s</sub>. Seven out of the 21 genes were selected, based on their biological relevance(s) in kidney development and cancer, for further investigation (**Table 2.1**). Two of these genes, Discoidin domain receptor family, member 1 (*DDR1*) and Homeodomain-interacting protein kinase 2 (*HIPK2*), are involved in cellular proliferation and/or apoptosis (**Table 2.2**) (127). Three of these genes, CXXC-type zinc finger protein 4 (*CXXC4*), Follistatin (*FST*), and *HIPK2*, play roles in signaling pathways which promote cellular proliferation and differentiation (**Table 2.2**) (127). Another gene, Myeloid/lymphoid or mixed-lineage leukemia, translocated to 3

(*MLLT3*), is involved in transcriptional regulation (**Table 2.2**) (127). Additionally, two genes, Mesenchyme Homeobox gene 2 (*MEOX2*), and Paired related homeobox 1 (*PRRX1*), are located in chromosomes 1 and 7, which have been identified in WTs with genetic aberrations. The dysregulation of these genes following mutation *WT1* suggests that *WT1* potentially regulates them during kidney development. Like *WT1*, these genes are expressed during embryogenesis, which also supports the hypothesis that *WT1* transcriptionally regulates these genes. Before we can further study the effect of inactivation of *WT1* on the gene expression of *CXXC4*, *DDR1*, *FST*, *HIPK2*, *MEOX2*, *MLLT3*, and *PRRX1*, the problem of false discovery from the array data has to be resolved.

A method to assess any false discoveries from the microarray is by quantitative PCR (qPCR) that quantifies the relative or absolute gene expression level within a given sample. In order to further investigate the effect of inactivation of *WT1* on the gene expression of *CXXC4*, *DDR1*, *FST*, *HIPK2*, *MEOX2*, *MLLT3*, and *PRRX1*, I validated all these genes with RNA isolated from an independent set of WTs by qPCR. Here, it is reported that the gene expression of *MEOX2*, one of the seven selected genes, stood out from the other validated genes and was significantly upregulated in *WT1*-mutant tumors. This suggested that *WT1* transcriptionally, directly or indirectly, regulates *MEOX2* during early kidney development and Wilms tumorigenesis.

**Table 2.1. Selected Differentially Expressed Genes from the Affymetrix GeneChip Array**

<b>Gene ID</b>	<b>Protein</b>	<b>Location</b>	<b>Fold Change</b>
<i>CXXC4</i>	CXXC-type zinc finger protein 4	4q22-q24	2.5-5X ↓
<i>DDR1</i>	Discoidin domain receptor family, member 1	6p21.3	2.5-3X ↓
<i>FST</i>	Follistatin	5q11.2	6-17X ↑
<i>HIPK2</i>	Homeodomain-interacting protein kinase 2	7q32-q34	2-5X ↓
<i>MEOX2</i>	Mesenchyme Homeobox gene 2	7p22-p21	19X ↑
<i>MLLT3</i>	Myeloid/lymphoid or mixed-lineage leukemia, translocated to 3	9p22	5-7X ↑
<i>PRRX1</i>	Paired related homeobox 1	1q24	4-7X ↑

**Table 2.2. Biological and Cellular Description of the Selected Genes**

Gene ID	Function*	Process*	Component*
Cxxc4	DNA binding	Wnt Signaling Pathway	Cytoplasm
	PDZ domain binding		Cytoplasmic membrane
	Metal ion binding		Cytoplasmic vesicle
	Zinc ion binding		
Ddr1	ATP binding	Cell adhesion	Basolateral plasma membrane
	Kinase activity	Ear development	Integral to membrane
	Nucleotide binding	Embryo implantation	Integral to plasma membrane
	Protein binding	Cell proliferation	Membrane
	Protein kinase activity	Mammary gland alveolus development	Plasma membrane
	Protein tyrosine kinase activity	Peptidyl-tyrosone phosphorylation	
	Receptor activity	Protein phosphorylation	
	Transferase activity	Transmembrane receptor protein tyrosine kinase signaling pathway	
Fst	Transmembrane receptor protein tyrosine kinase activity		
	Activin binding	BMP Signaling Pathway	Extracellular region
	Heparan sulfate proteoglycan binding	Female gonadal development	
		Gamete development	
		Hair follicle development	
		Hemopoietic progenitor cell differentiation	
		Keratinocyte proliferation	
		Cell differentiation	
Hipk2		Pattern specification process	
	ATP binding	DNA damage response	PML body
	SMAD binding	SMAD protein signal transduction	Centrosome
	Nucleotide binding	Anterior/posterior pattern formation	Cytoplasm
	Protein binding	Apoptosis	Nuclear body
	Protein kinase activity	BMP Signaling Pathway	Nuclear membrane
	Transcription corepressor activity	Phosphorylation	Nucleus
	Transferase activity	DNA binding	
	Virion binding	JNK cascade	
		Cell Proliferation	
		Smoothen signaling pathway	
		TGF-beta receptor signaling pathway	
Meox2	DNA binding	Angiogenesis	Cytoplasm
	Sequence-specific DNA binding	Limb development	Nucleus
	Sequence-specific DNA binding transcription factor activity	Multicellular organismal development	
	Transcription regulator activity	Palate development	
		Regulation of transcription, DNA-dependent	
		Skeletal muscle tissue development	
		Somite specification	

Gene ID	Function*	Process*	Component*
Mlt3	Protein binding	Anterior/posterior pattern formation	Nucleus
		Regulation of transcription, DNA-dependent	Cytoplasm
		Segment specification	
Prrx1	DNA binding	Artery morphogenesis	Nucleus
	Sequence-specific DNA binding	Cartilage development	
	Sequence-specific DNA binding transcription factor activity	Embryonic cranial skeleton morphogenesis	
	Transcription regulator activity	Embryonic skeletal system morphogenesis	
	Transcription repressor activity	Inner ear morphogenesis	
		Middle ear morphogenesis	
		Multicellular organismal development	
		Palate development	
		Regulation of transcription, DNA-dependent	

**Information provided by:**

Mouse Genome Informatics (MGI)\*

## Materials and Methods

### Wilms Tumor Samples

Twelve WT samples were used to validate the microarray data. Previously, WT samples, after informed consent, were collected from sporadic WT patients (50, 128). The DNA extracted from the tumors were analyzed for mutations by PCR amplification by the use of primers flanking each of the 10 exons of *WT1*, and followed by single-strand conformation polymorphism (SSCP) to confirm the abnormality (50, 129). The identification of *WT1* mutations was by the direct sequencing of aberrant PCR products (50)..

### Isolation of RNA from Human Wilms Tumor Samples

The isolation of RNA was previously achieved by either acid guanidinium thiocyanate-phenol-chloroform extraction, described by Chomczynski, P. and N. Sacchi (130), RNAqueous®-4PCR Kit from Applied Biosystems (Carlsbad, California), or RNeasy Midi Kit from Qiagen (Valencia, CA). Depending on the method used to isolate the RNA, it was kept in solution in either diethyl pyrocarbonate (DEPC) treated water or in ethanol. Samples were stored at -80°C.

## Quantitative PCR

qPCR is an efficient quantitative approach to validate the dysregulated genes from the microarray data. cDNAs were prepared by reverse transcription of 500 ng of total RNA from human *WT1*-wildtype and *WT1*-mutant Wilms tumors using TaqMan Reverse Transcription reagents with random hexamers from Applied Biosystems (Carlsbad, California). cDNAs were then amplified in duplicate in an ABI 7900HT Sequence Detection System thermal cycler using SYBR Green PCR Master Mix (Applied Biosystems). Primer Express v.3 software (Applied Biosystems) was used to design primer sets for the individual genes (**Table 2.3**). To determine the specificity of these primer sets, a dissociation curve of each primer set was assessed using human fetal kidney by qPCR analysis. Expression of glyceraldehyde phosphate dehydrogenase (*GAPDH*) was determined as an endogenous gene expression control. The standard deviation (SD) was determined for duplicate measurements of gene expression.

**Table 2.3. Primer Sets designed by Primer Express v.3 Software for the qPCR Validation**

<b>Name</b>	<b>Sequence (5'-3')</b>
<i>CXXC4 F</i>	AAAACCTGGCACTTCACTAGAGAGA
<i>CXXC4 R</i>	CTTTAAAAGAACCATCGGAATGCT
<i>DDR1 F</i>	CCCACCATCAGCTACCCAAT
<i>DDR1 R</i>	AAGTCTGCGATTTTGATGGTGAA
<i>FST F</i>	TGCTGGGCAGATCTATTGGAT
<i>FST R</i>	GATATCTTCACAGGACTTTGCTTTGA
<i>HIPK2 F</i>	CACCATGACACACTTACTCGATTTT
<i>HIPK2 R</i>	CACCCGACGCTTGCAGAT
<i>MEOX2 F</i>	CTGCGGAGGCGGAGAA
<i>MEOX2 R</i>	CTTGTAATTTCTTCCTGGGAGTCT
<i>MLLT3 F</i>	GCAGCAGATCGTGAACCTTATAGA
<i>MLLT3 R</i>	CGAGCAAAGATCAAATCAAATGT
<i>PRRX1 F</i>	CAGCGTCTCCGTACAGATCCT
<i>PRRX1 R</i>	CGTTATGAAGCCCCTCGTGTA




## Results

### Validation of the Affymetrix GeneChip Human Genome U133 Plus 2.0 Array

*CXXC4*, *DDR1*, *FST*, *HIPK2*, *MEOX2*, *MLLT3*, and *PRRX1* were validated by qPCR with a total of 12 WT samples, six *WT1*-wildtype WTs and six *WT1*-mutant WTs. *CXXC4*, *DDR1*, and *HIPK2* were downregulated and *FST*, *MEOX2*, *MLLT3*, and *PRRX1* were upregulated in *WT1*-mutant WTs by qPCR analysis (**Figure 2.1 and Table 2.4**). The upregulation of *MEOX2* stood out far from the rest of the dysregulated genes in the *WT1*-mutant WTs, therefore suggesting that *WT1* would normally transcriptionally repress the gene expression of *MEOX2*, but the loss of *WT1* results in the overexpression of *MEOX2*.

## qPCR Validation of the Affymetrix GeneChip Human Genome Array Data



	CXXC4	DDR1	FST	HIPK2	MEOX2	MLLT3	PRRX1
Relative Gene Expression	0.11	0.63	12.94	0.45	41.07	10.09	14.1
p-Value	0.06675	0.2017	0.00385	0.06469	0.00411	0.04114	0.06611

**Figure 2.1. Validation of the Dysregulated Genes in the *WT1*-mutant WTs**

In this graph, the y-axis represents the gene expression detected by the specific designed primer sets relative to the gene expression in the *WT1*-wildtype WTs. The individual color of the bars represents the gene found upregulated or downregulated in the *WT1*-mutant WTs.

## Discussion

It is important to identify genes that play a role in regulating cellular proliferation and differentiation. Upstream genetic alterations, like mutant *WT1*, result in the dysregulation of genes which may contribute to tumorigenesis. Our recent array analysis revealed that there are potentially 223 genes that were dysregulated following a *WT1* mutation. Literature searches, assessment of gene expression pattern(s), and functional role(s)) in kidney development and/or cancer focused studies to 21 genes. Seven out of the 21 genes, *CXXC4*, *DDR1*, *FST*, *HIPK2*, *MEOX2*, *MLLT3*, and *PRRX1*, were validated by qPCR. The validation data confirmed that these genes were dysregulated in *WT1*-mutant WTs.

When I compared the qPCR data to the array data, the dysregulation of their gene expression remained consistent and validated the array data. However, some the genes, *CCCX4*, *DDR1*, *HIPK2*, and *PRRX1*, were no longer significantly dysregulated in the *WT1*-mutant tumors.

One reason why there was a change in their significance may have resulted from the overall technique. Both microarray and qPCR use a reverse transcriptase enzyme (RT) that generates a complementary cDNA to the mRNA. However, in microarrays, the fluorescent labeled cDNAs are hybridized to their synthetic complementary DNAs attached on the microarray slide. The probes used in microarrays vary in their length, ranging from 10 to 99 bps. Instead of using a probe to hybridize to the complementary cDNA, primers or probes are designed to amplify a specific region of the gene (100-200 bps) in qPCR. Depending on the length and location of the probes or primers, the assessment of the significance in their gene

expression by qPCR may be slightly and/or drastically different from the microarray data.

Another potential reason for the differences in the degree of gene expression change may be due to the cell variation within the WT sample. As mentioned in Chapter 1, Wilms tumor could be made up of a mixture of mesenchymal, stromal, and epithelial cells, in which the relative proportions of these cells vary broadly in each tumor. The RNA isolated from these tumor samples may have one cell type predominant over the other cell types residing in the tumor. There is a possibility the genes from **Table 2.1** are expressed in a specific cell type.

Regardless, *FST*, *MEOX2*, and *MLLT3* were significantly dysregulated in *WT1*-mutant WTs. However, the dysregulation of *MEOX2* gene expression stood out more compared to the other genes. *MEOX2* is a member of a novel subclass of the homeobox gene super family (131). In humans, *MEOX2* is located at chromosomal region 7p21, one of the chromosomal regions found altered in WTs (131, 132). Homeobox genes, such as *MEOX2*, regulate cellular growth and differentiation during embryonic development and disease progression (133, 134).

Additionally, *MEOX2* is expressed during early embryonic development. Between E9-E9.5, *Meox2* is expressed in all of the formed somites and in the sclerotome, a population of cells lying between the paraxial mesoderm and the limb buds (131, 132). At E14.5, *Meox2* is expressed in the condensed mesenchyme of the kidney. *Meox2* gene expression in the developing kidney decreases as the metanephric mesenchymal cells differentiate into epithelial structures of the developing kidney (42).

Because of *MEOX2* drastic and significant change in mutant-*WT1* WTs, biological functions, and embryonic gene expression, *MEOX2* is potentially a direct candidate downstream gene of *WT1* during Wilms tumorigenesis. These suggest that there is a transcriptional relationship between *WT1* and *MEOX2* expression in WT. However, more investigation is needed to completely understand the relationship between *WT1* and *MEOX2*, such as when and where does the loss of *WT1* affect the gene expression of *MEOX2* during kidney development.

### **Chapter 3: Generation of a Kidney Mesenchymal Cell Line**

Mammalian kidney development results from reciprocal interactions between the ureteric bud and the undifferentiated metanephric mesenchyme. Because WT is thought to arise from undifferentiated metanephric mesenchyme, it is important to study Wilms tumorigenesis during early kidney development. In order to study Wilms tumorigenesis at the cellular and molecular level, we intended to use human embryonic kidney (HEK) 293 cells, which were transformed with sheared adenovirus 5 DNA to be cultured and passaged for a long period of time (135). However, experiments to determine the effect of the loss of *WT1* in the HEK293 cells were not feasible, because these cells did not expression high amounts of endogenous WT1 by IHC. Therefore, I generated an immortalized undifferentiated kidney mesenchymal cell line in which endogenous WT1 is robustly expressed and can be genetically ablated to efficiently study the effect of the inactivation of *Wt1* on putative downstream target genes.


## Materials and Methods

### Mouse Strains and Breeding

The  $Wt1^{flox/flox}$  mouse generated and described by Gao et, al, (112) and the Immortomouse<sup>®</sup> (CBA;B10-Tg(H2Kb-tsA58)6Kio/Crl) purchased from Charles River (Wilmington, MA) were crossed together to obtain  $Wt1^{flox/flox}$ ; *Immorto* mice. The purpose of this mating cross was to generate a mouse  $Wt1^{flox/flox}$  strain that carried the Immorto allele for conditional cellular immortalization as described by Jat et. al (136). The mating cross consisted of  $Wt1^{flox/flox}$ ; *Immorto* female mice and  $Wt1^{+/-}; Cre-ER^{TM}$  male mice was to generate the following genotypes:  $Wt1^{-/flox}$ ; *Immorto*,  $Wt1^{+/flox}$ ; *Immorto*,  $Wt1^{-/flox}$ ; *Cre-ER^{TM}*,  $Wt1^{+/flox}$ ; *Cre-ER^{TM}*,  $Wt1^{-/flox}$ ; *Immorto*; *Cre-ER^{TM}* and  $Wt1^{+/flox}$ ; *Immorto*; *Cre-ER^{TM}*. The *Cre-ER^{TM}* inducible model will allow the Cre recombinase to recognize the loxP sites around exons 8 and 9 of *Wt1*, which, in turn, will result in a mutant, non-functional WT1 protein (**Figure 3.1**) in mice with the genotype of  $Wt1^{-/flox}$ ; *Immorto*; *Cre-ER^{TM}*. All of the embryos from these crosses will be genotype as described below.

### Dissociation of the Kidney Mesenchymal Cells

The morning of the vaginal plug was defined as E.5. All of the embryos were harvested at E13.5 and the kidney rudiments were dissected out. Both of the kidney rudiments from each embryo were temporarily stored on ice in 50  $\mu$ L of 1X trypsin from MediaTech (Manassas, VA). Once all the kidneys were dissected out from each embryo, the kidney rudiments in the 1X trypsin were placed in the water bath at 37°C for 5 minutes. The cells from kidney rudiments were then dissociated into single cell suspensions by the use of the 10  $\mu$ L pipette.



**Figure 3.1. Schematic Diagram of the Cre-Lox System**

The 1.55 and 1.75 primers (black arrows) are located around the loxP sites around exons 8 and 9 on the floxed allele. Once the Cre-recombinase recognizes these loxP sites, an in-frame deletion is created, resulting in the deletion of exons 8 and 9 to generate a mutant, non-functional WT1 protein.



### Culturing of the Metanephric Mesenchymal Cells (MMCs)

Cells were seeded in a 24-well plate and cultured in permissive conditions: Dulbecco's modified Eagle's medium (DMEM) supplemented with 60 U/mL of interferon  $\gamma$  (IFN- $\gamma$ ) from Sigma-Aldrich (St. Louis, MO) and 100 U/mL of antibiotics (penicillin, streptomycin, and amphotericin B) (MediaTech), 10% Fetal Bovine Serum (FBS), and cultured at 33°C. At this temperature, the temperature sensitive large T-antigen (TA-g) was produced to ultimately result in the immortalization of the MMCs (136). The MMCs were subsequently passaged after enzymatic dissociation in 1X trypsin (MediaTech) in DMEM for 5 minutes at 37°C. Once the MMCs reached 40 passages, they were considered an immortalized cell line (136).

### Molecular Characterization of the MMCs

A total of 20 clones (the *Wt1*<sup>+/*fl*</sup>; *Immorto* MMCs) were subcloned by end point dilution. Colonies were selected based on their cell morphology that were consistent with the established mesenchymal cell morphological traits, such as being elongated, spindle-shaped, and fibroblastic looking (136). The isolated cells did not resemble epithelial cells, which appear to be cuboidal-shaped and well organized. The F11249 *Wt1*<sup>-/*flox*</sup>; *Immorto*; *Cre-ER*<sup>TM</sup>, F11254 *Wt1*<sup>-/*flox*</sup>; *Immorto*; *Cre-ER*<sup>TM</sup>, and F12055 *Wt1*<sup>-/*flox*</sup>; *Immorto* cell lines were not yet subcloned. Immunocytochemistry analyses of all the cell lines indicated whether the cell lines were mesenchymal using two mesenchymal markers, Vimentin and Cited1, and an epithelial marker, Cytokeratin. The MMC cell lines were seeded onto round, glass cover slips (Fisher) and fixed with methanol for 10 minutes and washed off with 1X PBS three times for 10 minutes each. Cells were incubated with 10% goat serum in PBS for 1-2 hour at room temperature or

overnight at 4°C, depending on the antibody. The fixed cells were then incubated overnight at 4°C with either anti-WT1 monoclonal antibody (1:100) (DAKO), anti-MOX2 monoclonal (1:50) (Santa Cruz), anti-Vimentin monoclonal (1:100), anti-Cited1 (1:100) (Fisher) or anti-Cytokeratin monoclonal (1:100) from Sigma-Alrich. After being washed three times with PBS, the fixed cells were incubated with conjugated Alexa Fluor 594 anti-mouse secondary antibody (Invitrogen). The cells on the cover slips were mounted in VECTASHIELD mounting media (Vector Laboratories) and images were captured by a Leica Epifluorescence/Brightfield Microscope.

### Adenovirus Vectors

The Cre recombinase and enhanced green fluorescent protein (eGFP) within an internal ribosome entry site (IRES) expression cassette (Ad-Cre-IRES-eGFP) packaged inside an Adenovirus vector (type 5, DE1/E3) was purchased from Vector Biolabs (Philadelphia, PA). The viral infection control was a recombinant human adenovirus type 5 expressing eGFP under the control of a cytomegalovirus (CMV) promoter (Ad-CMV-GFP) purchased from Vector Biolabs. The purpose of eGFP was to monitor the expression of the viral construct after the viral infection. The Adeno-Cre-eGFP and Adeno-eGFP stocks (titer of  $1 \times 10^{10}$  PFU/ml) were stored in DMEM with 2% Bovine Serum Albumin (BSA), and 2.5% Glycerol.

### Conditional Ablation of *Wt1* in the MMCs

After 24hrs of seeding the F11249 *Wt1*<sup>-flox</sup>; *Immorto*; *Cre-ER*<sup>TM</sup> cells or the F11254 *Wt1*<sup>-flox</sup>; *Immorto*; *Cre-ER*<sup>TM</sup> cells into a 24-well plate, the cells were treated with 1 $\mu$ M of 4-hydroxy tamoxifen (4-OH tamoxifen) for 48hrs. The mutant form of the ligand binding domain of the estrogen receptor (ER<sup>TM</sup>) prevents binding of its natural ligand, 17 $\beta$ -estradiol, at normal physiological concentrations, but allows the ER<sup>TM</sup> domain responsive to 4-OH tamoxifen (137-139). The fusion of *Cre* with *ER*<sup>TM</sup> results in the ER<sup>TM</sup>-dependent cytoplasmic sequestration of *Cre* by Hsp90, which prevents *Cre*-mediated recombination (140, 141). However, binding of 4-OH tamoxifen disrupts the interaction Hsp90 that permits access of *Cre-ER*<sup>TM</sup> to the nucleus and initiation of recombination. The *Cre* recombinase recognizes the loxP sites flanking exons 8 and 9 of *Wt1*. The enzymatic reaction by *Cre* recombinase results in a mutant, non-functional WT1 protein (**Figure 3.1**).

The ablation of *Wt1* in the F12055 *Wt1*<sup>-flox</sup>; *Immorto* MMC cell line was done by adenoviral infection that carried the *Cre* recombinase. The *Wt1*<sup>-fl</sup>; *Immorto* MMCs were infected with 10 multiplicity of infection (MOI) of Adeno *Cre* for 48 hours. The medium was aspirated and replaced with fresh DMEM supplemented with 60 U/mL of IFN- $\gamma$  and 100 U/mL of antibiotics.

## Genotyping

DNA isolated from the MMCs was used for genotyping by PCR amplification conditions (**Table 3.1**) by the primer sets in (**Table 3.2**). The detection of *Wt1*-null allele was accomplished as described by Kreidberg et, al (43) The presence of the *Wt1* flox allele was determined using the 1.75/1.55 primer set, in which primers are located around the loxP sites flanked around exons 8 and 9. The *Cre-ER<sup>TM</sup>* transgene was detected by the *ZP3-se* and *ZP3-as* primer set (44). The *Immorto* allele was detected by the primer set *Immo I* and *Immo II*. The recombined *Wt1* flox allele (*Wt1<sup>Δ</sup>*) was detected by PCR amplification using the *ckodeIF* and 1.55 (112).

**Table 3.1. PCR Conditions for Animal Genotyping**

	<b>Initial Denature</b>	<b>Cycle Denature</b>	<b>Cycle Anneal</b>	<b>Cycle Extension</b>	<b>Cycles</b>	<b>Final Extension</b>	<b>Product Size</b>
<b>Primer Set</b>							
<i>Wt1-F/PGK</i>	94°C - 3 min	94°C - 45 sec	58°C - 45 sec	72°C - 50 sec	35	72°C - 7 min	250 bps
<i>1.55/1.75</i>	94°C - 3 min	94°C - 45 sec	58°C - 45 sec	72°C - 50 sec	35	72°C - 7 min	150 and 200 bps
<i>ckodeIF /1.55</i>	94°C - 3 min	94°C - 45 sec	58°C - 45 sec	72°C - 50 sec	35	72°C - 7 min	450 bps
<i>ZP3-se/ZP3-as</i>	94°C - 3 min	94°C - 45 sec	58°C - 45 sec	72°C - 50 sec	35	72°C - 7 min	600 bps
<i>Immo I/Immo II</i>	94°C - 4 min	94°C - 30 sec	58°C - 1 min	72°C - 1 min	35	72°C - 7 min	1 kb

**Table 3.2. Primer Sequences for Animal Genotyping**

<b>Name</b>	<b>Sequence (5'-3')</b>
<i>Wt1-F</i>	GTG ACC CCG CAG CTA GCC
<i>PGK</i>	CCA TTT GTC ACG TCC TGC
<i>1.55</i>	TGC CTA CCC AAT GCT CAT TG
<i>1.75</i>	GAA ACT GTT TGT AAC GAG AG
<i>ckodellF</i>	GCT AAC ATA TGG GAG ACA TT
<i>ZP3-se</i>	TCC AAT TTA CTG ACC GTA CAC CAA
<i>ZP3-as</i>	CCT GAT CCT GGC AAT TTC GGC TA
<i>Immo I</i>	AGC GCT TGT GTC GCC ATT GTA TTC
<i>Immo II</i>	GTC ACA CCA CAG AAG TAA GGT TCC


## Results

### Genotyping of the MMCs

Genotyping confirmed that clones # 1, 2, 3, 4, 7, 10, 11, 12, 15, 16, and 18 were *Wt1*<sup>+/fl</sup>; *Immorto* MMCs and F11249 E3 and F11254 E2 cell lines were *Wt1*<sup>-/floX</sup>; *Immorto*; *Cre-ER*<sup>TM</sup> MMCs (**Figure 3.2**). Cell line F11254 E3 was *Wt1*<sup>fl/fl</sup>; *Immorto*; *Cre-ER*<sup>TM</sup>, and F12055 was *Wt1*<sup>-/fl</sup>; *Immorto* (**Figure 3.5**).

Two of the cell lines, F11249 *Wt1*<sup>-/floX</sup>; *Immorto*; *Cre-ER*<sup>TM</sup> and F11254 *Wt1*<sup>-/floX</sup>; *Immorto*; *Cre-ER*<sup>TM</sup> were tamoxifen-inducible cell lines, which *Wt1* can be genetically ablated by the Cre-recombinase. By PCR analysis, the ablation of *Wt1* occurred after 48 and 72 hrs as shown in **Figure 3.3**. However, the ablation of *Wt1* in these cells was incomplete, as indicated by the residual presence of the floxed allele.

The ablation of *Wt1* in the F12055 *Wt1*<sup>-/floX</sup>; *Immorto* MMCs cell line was achieved by adenoviral infection. As shown in **Figure 3.5**, GFP positive cells indicated that the cells were successfully infected by the adenovirus. PCR analysis confirmed the presence of the *Cre-ER*<sup>TM</sup> transgene and demonstrated the recombination of the *Wt1*<sup>fl</sup> allele 48 hrs after of the adenoviral infection (**Figure 3.5**). Most important, the ablation of *Wt1* after the adeno-viral infection of Cre-recombinase was complete; the floxed allele was no longer present (**Figure 3.5**).




Lane#	MMCs DNA Samples	Genotype
1	F736 E6 clone#1	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
2	F736 E6 clone#2	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
3	F736 E6 clone#3	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
4	F736 E6 clone#4	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
5	F736 E6 clone#7	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
6	F736 E6 clone#10	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
7	F736 E6 clone#11	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
8	F736 E6 clone#12	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
9	F736 E6 clone#15	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
10	F736 E6 clone#16	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
11	F736 E6 clone#18	<i>Wt1</i> <sup>+/-fl</sup> ; <i>Immorto</i>
12	F11249 E3 #1 (15)	<i>Wt1</i> <sup>-/-lox</sup> ; <i>Immorto</i> ; <i>Cre-ER</i> <sup>TM</sup>
13	F11249 E3 #2 (14)	<i>Wt1</i> <sup>-/-lox</sup> ; <i>Immorto</i> ; <i>Cre-ER</i> <sup>TM</sup>
14	F11254 E2 #5 (15)	<i>Wt1</i> <sup>-/-lox</sup> ; <i>Immorto</i> ; <i>Cre-ER</i> <sup>TM</sup>
15	F11254 E3 #2 (16)	<i>Wt1</i> <sup>-/-lox</sup> ; <i>Immorto</i> ; <i>Cre-ER</i> <sup>TM</sup>
16	F11254 E3 #5 (17)	<i>Wt1</i> <sup>-/-lox</sup> ; <i>Immorto</i> ; <i>Cre-ER</i> <sup>TM</sup>
17	#52 (Fl allele Control)	<i>Wt1</i> <sup>fl/fl</sup>
18	#70 ( $\Delta$ allele Control)	<i>Wt1</i> <sup>-/<math>\Delta</math></sup>
19	#20 (Cre(-) Control)	<i>Wt1</i> <sup>+/-+</sup>
20	dH <sub>2</sub> O	-

**Figure 3.2. PCR Analyses of the MMCs using *Wt1*-F/PGK, 1.55/1.75, ZP3 se/ZP3 as, and *Immo I*/*Immo II* primer sets**

Cells isolated from E13.5 embryos were genotyped using the established primer sets to detect the *Wt1* null allele, floxed allele, *CreER*<sup>TM</sup> allele, and the *Immorto* allele (**Table 3.1**). Genotypes are shown to the right.






Lane #	Description
1	48 hrs Untreated Set 1
2	48 hrs 1 $\mu$ M 4-OH-Tamoxifen Set 1
3	48 hrs Untreated Set 2
4	48 hrs 1 $\mu$ M 4-OH-Tamoxifen Set 2
5	72hrs Untreated Set 1
6	72 hrs 1 $\mu$ M 4-OH-Tamoxifen Set 1
7	72 hrs Untreated Set 2
8	72 hrs 1 $\mu$ M 4-OH-Tamoxifen Set 2
9	dH <sub>2</sub> O


**Figure 3.3. Ablation of *Wt1* by Tamoxifen Treatment**

As shown in duplicates, F11249 E3 *Wt1*<sup>-fl</sup>; *Immorto*; *Cre*<sup>TM</sup> cell line was treated with 1  $\mu$ M of Tamoxifen for 48 hrs and at 72 hrs the recombined *Wt1*<sup>Δ</sup> allele is detected. However, the floxed allele was still present, which indicated only partial ablation of *Wt1*.



**Figure 3.4. Viral Infection of Adeno-Cre-GFP in the F12055 *Wt1*<sup>fllox</sup>; *Immorto* MMC cell line**

The top panel of images was taken by Differential Interference Contrast microscopy (DIC) and the middle panel of images was taken with a GFP-FITC (GFP-fluorescein isothiocyanate) filter to detect the excitation at 519 nanometers in the green region to indicate the cells that have been infected and expressed GFP. All of the images were taken at 20X magnification. The lower panel of images merged DIC and FITC images together to determine the number of cells that were infected by the adenovirus. NTC – Non Treated Cells



**Figure 3.5. PCR Analyses of the Ablation of *Wt1* in the F12055 *Wt1*<sup>-flox</sup>; *Immorto* MMC cell line infected by Adeno-Cre-eGFP or Adeno-eGFP**

The complete ablation of *Wt1* was detected after 48hrs. All the cells were confirmed to have the *Wt1* null allele (yellow arrow) and the recombined allele (red arrow) by using the primers in **Table 3.1**. (C=Cells infected with Cre and eGFP adenovirus, G= Cells infected with eGFP only, and N= Non-Treated cells)

## Molecular Characterization of the MMCs

All of the cells displayed in **Table 3.3** reached passage 40 and were considered immortalized cell lines. The wild type  $Wt1^{+/fl}$ ; *Immorto* MMCs were sub-cloned based on their cell morphology. Ten clones of  $Wt1^{+/fl}$ ; *Immorto* MMCs were generated, and were vimentin positive and cytokeratin negative (**Appendix Figure 1**). Of the 10 clones, Clone# 10 expressed more of both Wt1 and Meox2 (**Figure 3.6 A**).  $Wt1^{-/flox}$ ; *Immorto* cell line, F12055, expressed Wt1 and Meox2 similar to Clone #10 (**Figure 3.6 B**). An established mesenchymal cell marker, Cbp/p300-interacting transactivator 1 (*Cited1*), was used to ensure that the MMCs were derived from the cap mesenchyme. *Cited1* is known to be specifically expressed in the condensed cap mesenchyme with no expression in the clefts between the ureteric bud or the surrounding stromal mesenchyme (142, 143). As shown in **Figure 3.7**,  $Wt1^{+/flox}$ ; *Immorto* and F12055  $Wt1^{-/flox}$ ; *Immorto* MMCs were positive for *Cited1* further indicating the mesenchymal status of the cell lines.

The  $Wt1^{-/flox}$ ; *Immorto*; *CreER<sup>TM</sup>* (F11249 E3 and F11254 E2) and the  $Wt1^{-/flox}$ ; *Immorto* cell lines have not been sub-cloned based on their cell morphology. These cells are still a heterogeneous population of vimentin and cytokeratin positive cells. In addition,  $Wt1^{-/flox}$ ; *Immorto*; *CreER<sup>TM</sup>* (F11249 E3 and F11254 E2) did not express Wt1 (**Appendix Figure 2**).

**Table 3.3 MMCs Cell Lines Derived from E13.5 Kidney Rudiments**

Wt1, Meox2, Vimentin, Cited1, and Cytokeratin were the markers used to determine whether the cells were mesenchymal.

<b>Cell Line</b>	<b>Genotype</b>	<b>Wt1</b>	<b>Meox2</b>	<b>Vimentin</b>	<b>Cited1</b>	<b>Cytokeratin</b>
Clone # 1	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+	NT	+	NT	-
Clone # 2	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+ weak	NT	+	NT	-
Clone # 3	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+ weak	NT	+	NT	-
Clone # 4	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+	NT	+	NT	-
Clone # 5	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	-	NT	+	NT	-
Clone #7	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+	NT	+	NT	-
Clone # 8	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+ weak	NT	+	NT	-
Clone # 10	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+	+	+	+	-
Clone # 11	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+ weak	NT	+	NT	-
Clone # 15	<i>Wt1</i> <sup>+/-fl.</sup> ; <i>Immorto</i>	+ weak	NT	+	NT	-
F11249 E3	<i>Wt1</i> <sup>-/-fl.</sup> ; <i>Immorto</i> ; <i>CreER</i> <sup>TM</sup>	-	NT	+	NT	+*
F11254 E2	<i>Wt1</i> <sup>-/-fl.</sup> ; <i>Immorto</i> ; <i>CreER</i> <sup>TM</sup>	-	NT	+	NT	+*
F12055	<i>Wt1</i> <sup>-/-fl.</sup> ; <i>Immorto</i>	+	+	+	+	+*


**(+)** - Expressed the marker

**(-)** - Did not express the marker

**Weak** – The intensity of the IF staining was weak compared to the control cell line (MCF-7 cells).

**NT**- Not Tested


\* - The cell line was not subcloned.



**Figure 3.6. IF Images of the Generated MMCs from E13.5 Kidney Rudiments**

**A.** The top panel of images is Clone #10  $Wt1^{+/flox}$ ; *Immorto* MMC cell line and the bottom panel of images is the control, MCF-7 cell line. As shown, Clone #10  $Wt1^{+/flox}$ ; *Immorto* MMCs were WT1, MEIOX2, vimentin positive and cytokeratin negative. DAPI (blue) stains the nucleus.

**B.** The top panel of images is F12055  $Wt1^{-/flox}$ ; *Immorto* MMC cell line and the bottom panel of images is the control, MCF-7 cell line. F12055  $Wt1^{-/flox}$ ; *Immorto* MMC cell line expressed both WT1 and MEIOX2. Because the F12055  $Wt1^{-/flox}$ ; *Immorto* MMC cell line were not sub-cloned, it is presently a heterogeneous population of vimentin and cytokeratin positive cells.



**Figure 3.7. MMCs are Positive for Cited1: Biological Marker for Condensed Cap Mesenchyme**

Clone #10  $Wt1^{+/fl}; Immorto$  MMCs (top panel) and the F12055 E4  $Wt1^{-/fl}; Immorto$  (bottom panel) were positive for CITED1, which confirmed that these cells were isolated from the condensed cap mesenchyme. DAPI (blue) stains the nucleus.

## Discussion

The HEK 293 cell line was the first transformed human cell line. Since these cells were isolated from the kidney, I thought it would serve a good cellular model to study the effect of the loss of *Wt1* on the gene expression of *Meox2*. Hence, I proposed to manipulate the gene expression of *Wt1* either by siRNA and/or shRNA to assess the effect on *Meox2* gene expression. However, these cells were more epithelialized cells and they did not express a robust amount of *Wt1*. Therefore, I generated several *Wt1* kidney mesenchymal cell lines: wildtype *Wt1* MMCs (*Wt1*<sup>+/fl</sup>; *Immorto*) and the mutant *Wt1* MMCs F11249 E3 and F11254 E2 (*Wt1*<sup>fl</sup>; *Immorto*; *Cre*<sup>TM</sup>) and F12055 (*Wt1*<sup>-fl</sup>; *Immorto*).

After the molecular characterization of all the cell lines (**Table 3.2**), I chose two cell lines, Clone #10 *Wt1*<sup>+/flox</sup>; *Immorto* cell line and F12055 *Wt1*<sup>-flox</sup>; *Immorto* cell line for my experiments to study the effect of the inactivation of *Wt1* on the gene expression of *Meox2* *in vitro*. Both of these cell lines displayed mesenchymal morphological traits of being elongated, spindle-shaped, and fibroblastic looking. Additionally, the total ablation of *Wt1* was achieved in the F12055 *Wt1*<sup>-fl</sup>; *Immorto* cell line after infection with Adeno-Cre-GFP. By using these molecular tools, we can understand how a *Wt1* mutation effects *Meox2* and other potential downstream genes in kidney mesenchymal cells that may contribute to Wilms tumorigenesis. We can assess *WT1*-ablated cells for tumorigenic phenotypes, such as increased cellular proliferation, inhibition of apoptosis, disruptions in cellular adhesion and migration. Furthermore, the cell lines generated were not just limited for the purpose of Wilms tumorigenesis, but can be used to experimentally study the loss of *Wt1* in other kidney cancers and during kidney development.




## Chapter 4: Inactivation of *Wt1* effects the Gene Expression of *Meox2* during Kidney Development

Mesenchyme homeobox 2 (*MEOX2*), formally known as *GAX* and *MOX2*, was one of the seven genes from the prioritized microarray data and was significantly up regulated by 19-fold in the *WT1* mutant Wilms tumors. *MEOX2* is a member of a novel subclass of the homeobox gene super family which is located at chromosomal region 7p21 in humans and on chromosome 12 in mice (131, 132). *MEOX2* consists of three exons and two introns. It also contains a nuclear localization signaling (NLS) as shown in **Figure 4.1 A** (131). Homeobox genes like *MEOX2* encode homeodomains that correspond to a 60 amino acid helix-turn-helix DNA binding region, located near the C-terminus of the protein (**Figure 4.1 B**) (131, 133, 144). Homeobox proteins localize to the nucleus and are transcription factors which regulate the expression of lineage specific genes which, in turn, control cellular and organ differentiation (145).

*MEOX2* is expressed during early embryonic development. Between E9-E9.5, *Meox2* is expressed in all of the formed somites and in the sclerotome, a population of cells lying between the paraxial mesoderm and the limb buds (131, 132). At E14.5, *Meox2* is expressed in the condensed mesenchyme of the kidney (**Figure 4.2**). *Meox2* gene expression in the developing kidney decreases as the metanephric mesenchymal cells differentiated into epithelial structures of the developing kidney (42). The embryonic gene expression of *Meox2* suggests that it has a specific role during development.

**A.**




**B.**

MEHPLFGCLRSPHATAQGLHPFSQSSLALHGRSDHMSYPELSTSSSSCIIAGYPNEEGMF  
 ASQHHRGHHHHHHHHHHHHHHHQQQQHQALQSNWHLPMSSPPSAARHSLCLQPDGGPPELG  
 SSPPVLCSSSSSLGSSTPTGAACAPGDYGRQALSPADVEKRSGSKRKSDSSDSQEGNYKS  
 EVNSKPRKERTAFTEQIRELEAEFAHHNYLTRLRRIEIAVNLDLTERQVKVWFQNRRMK  
WKRVKGGQQGAAAREKELVNVKKGTLLPSELGIGAATLQQTGDSLANEDSRDSDHSSEH  
 AHL

**Figure 4.1. Representation of Meox2 Gene and Protein Sequence**

**A.** *Meox2* is a homeobox gene that contains 3 exons (light blue) and 2 large introns. The homeobox domain (dark blue) is encoded by exons 2 and 3.

**B.** Amino acid sequence of *Meox2* gene product. The homeodomain (highlighted in yellow) starts in exon2 at position 187 and ends in exon 3 at position 245. The NLS sequence (underlined in red) is located within exon 3.



**Figure 4.2. *Meox2* Gene Expression in E14.5 Mouse**

At E14.5, *Meox2* is expressed in the condensed metanephric mesenchyme (Adapted from (146)).

In normal mouse mammary epithelial NMuMG cells, *Meox2* expression has been shown to be regulated by Tgf- $\beta$ 1 (147). After 16-24 hrs of Tgf- $\beta$ 1 stimulation, the gene expression of *Meox2* increased 2-8X in NMuMG cells. Dominant-negative mutants of *Smad3* and *Smad4* blocked the induction of *Meox2*, further indicating that the regulation of *Meox2* was dependent on the stimulation of Tgf- $\beta$ 1(147). Furthermore, the endogenous expression of *Meox2* by Tgf- $\beta$ 1 inhibits proliferation by inducing *p21* in the NMuMG cells (147). However, these data do not demonstrate that *Meox2* expression is dependent only by Tgf- $\beta$ 1 stimulation and whether the regulation of *Meox2* is cell specific. Other genes throughout development could regulate *Meox2*. *MEOX2* is upregulated in the *WT1* mutant WTs, and *Meox2* and *Wt1* are both expressed during kidney development in the condensed mesenchyme. These data suggest that *Wt1* may regulate the gene expression of *Meox2* during developing kidney.

In order to assess whether there are any predicted WT1 binding sites in *MEOX2*, the Match™ web-based tool from the BIOBASE Biological Database was used to identify transcription factor binding sites in the *MEOX2* gene by a weight matrix search. The Match™ web-based tool was able to identify one predicted *Wt1* binding site (cCCTCCccc) within the first intron of *MEOX2* suggesting that WT1 may regulate *MEOX2*. Having identified a putative WT1 binding site within the *MEOX2* gene, I sought to determine whether modulation of *Wt1* expression resulted in a corresponding change in *Meox2* expression *in vivo* in developing mouse kidney and also in the kidney mesenchymal cell lines. I further sought to determine the role, if any, of Tgf- $\beta$ 1 in regulating *Meox2* expression in these cells. Moreover, I investigated the phenotypic effect of *Wt1* ablation in this experimental system. Additionally, I

investigated whether the inactivation of *Wt1* affects the gene expression of the established regulator, Tgf- $\beta$ 1, of *Meox2*. The dysregulation of *p21* and *Itgb5* resulting from the upregulation of *Meox2* was also assessed within this study.

Within this chapter, I have demonstrated that following the ablation of *Wt1* in E13.5 kidney rudiments and in the kidney mesenchymal cell lines, *Meox2* expression was upregulated. Data revealed that Tgf- $\beta$ 1 did not contribute to the dysregulation of *Meox2*. Moreover, the established downstream genes of *Meox2* (*p21* and *Itgb5*) were dysregulated after the upregulation of *Meox2* in the kidney mesenchymal cells. A novel phenotype was observed in the presence of a *Wt1* mutation and the upregulation of *Meox2*; the cells decreased in cellular growth and loss the ability to adhere to the cell culture plates. Overall, these significant data further supported my hypothesis that the gene expression of *Meox2* is upregulated both *in vivo* and *in vitro*, following a *Wt1* mutation during early kidney development.

## Materials and Methods

### Mouse Strains

Refer to **Chapter 3** – Materials and Methods – Mouse Strains.

### In Vivo Ablation of *Wt1*

The *Wt1*<sup>-flox</sup> mouse model generated and described by Gao et, al, was used to genetically ablate *Wt1* exons eight and nine *in vivo* (112). Crosses were set up with the females' genotype *Wt1*<sup>flox/flox</sup> and the males' genotype *Wt1*<sup>+/-</sup>; *Cre-ER*<sup>TM</sup>. At E11.5, the females were injected with 3mg of Tamoxifen to ablate *Wt1*.

### RNA Isolation from Embryonic Kidney Tissue

Kidneys were dissected from embryos at E13.5 and were placed into 1mL TRIzol purchased from Invitrogen (Carlsbad, CA). The tissues were completely homogenized in the TRIzol and precipitated by isopropanol. The RNA was rinsed off with 70% ethanol. The RNA pellet was resuspended in RNase-free distilled water. The RNA was treated with 1 µl of DNase 1 (Applied Biosystems) for 30 minutes to remove any traces of DNA contamination.

### Genotyping of the Kidney Rudiments

Limbs of the dissected E13.5 embryos were used to extract DNA. PCR genotyping as previously described in Chapter 3. The PCR conditions are displayed in **Table 3.1**. The primer sets used for the genotyping are displayed in **Table 3.2**.

### In Vitro Ablation of *Wt1* in the MMCs

Immediately after seeding into 60mm plates, F12055 MMCs were infected with 10 multiplicity of infection (MOI) of Adeno Cre for 48 hours. The medium was aspirated and replaced with fresh DMEM supplemented with 60 U/mL of IFN- $\gamma$  and 100 U/mL of antibiotics. As a control, a second plate of cells was infected with Adeno GFP of 10 MOI. The recombined allele was detected by PCR as described in Chapter 3 in Table 3.1.

### RNA Isolation from the Infected MMCs

After each time point (48, 72, and 96 hours), cells were harvested, centrifuged at 7500 revolutions per minute (rpm) for 5 minutes, and resuspended in TRIzol to isolate RNA.. A phenol-chloroform extraction was done to remove the residual salt in the RNA sample. The RNA was re-precipitated by 5M Ammonium Acetate, Linear Acrylamide, and 100% ethanol at -20C. The RNA pellet was resuspend in RNase-free distilled water. The NanoDrop spectrophotometer from Thermo Scientific (Wilmington, DE) was used to quantify the RNA. RNA samples were stored at -80°C.

### qPCR

cDNAs were prepared from E13.5 kidney and MMCs RNA by reverse transcribing 1  $\mu$ g of total RNA as described in **Chapter 2** Materials and Methods. The primer sets for *Wt1*, *Meox2*, *Tgf- $\beta$ 1*, *Smad3*, *p21*, and *intergrin  $\beta$ 5* are displayed in **Table 4.1**.

**Table 4.1. Primer Sets designed by Primer Express v.3 Software for the qPCR analysis using cDNA from E13.5 kidney and MMCs**


<b>Name</b>	<b>Sequence (5'-3')</b>
<i>Wt1 F</i>	CAAGGACTGCGAGAGAAGGTTT
<i>Wt1 R</i>	TGGTGTGGGTCTTCAGATGGT
<i>Meox2 F</i>	GCTGTCACCCGCAGACGTA
<i>Meox2 R</i>	AATCTGAGCTGTCGCTTTTCCT
<i>Tgf-<math>\beta</math>1 F</i>	AAACGGAAGCGCATCGAA
<i>Tgf-<math>\beta</math>1 R</i>	TGGCGAGCCTTAGTTTGG
<i>Smad3 F</i>	GGAATGCAGCCGTGGAAC
<i>Smad 3 R</i>	GACCTCCCCTCCGATGTAGTAGA
<i>p21 F</i>	ACGTGGCCTTGTCGCTGTC
<i>p21 R</i>	AATCTGCGCTTGGAGTGATAG
<i>Itgb5 F</i>	AGGCTGGGACGTCATTCAGA
<i>Itgb5 R</i>	GCGAACCTGTAGCTGAAGGT



## Results

### In Vivo Ablation of *Wt1* during Early Kidney Development


The Cre-mediated ablation of *Wt1* did occur, which resulted in the recombined allele produced by the CKO primer set as shown in **Figure 4.3**. However, the ablation of *Wt1* did not occur in all the cells, because the floxed allele was still present after the kidney rudiments were genotyped by PCR as described in Chapter 3 (**Figure 4.3**). However, *Wt1* was genetically knocked out in ~70% (.0004) of cells at E13.5 and resulted in the increase of *Meox2* relative gene expression by 2X (.018) as shown in **Figure 4.4**.



Lane #	Genotype	Lane #	Genotype
1	<i>Wt1</i> <sup>-fl</sup>	10	<i>Wt1</i> <sup>-fl</sup> ; <i>Cre</i> <sup>TM</sup>
2	<i>Wt1</i> <sup>+fl</sup> ; <i>Cre</i> <sup>TM</sup>	11	<i>Wt1</i> <sup>-fl</sup>
3	<i>Wt1</i> <sup>+fl</sup> ; <i>Cre</i> <sup>TM</sup>	12	<i>Wt1</i> <sup>-fl</sup> ; <i>Cre</i> <sup>TM</sup>
4	<i>Wt1</i> <sup>-fl</sup>	13	<i>Wt1</i> <sup>-fl</sup> ; <i>Cre</i> <sup>TM</sup>
5	<i>Wt1</i> <sup>+fl</sup>	14	<i>Wt1</i> <sup>-fl</sup> ; <i>Cre</i> <sup>TM</sup>
6	<i>Wt1</i> <sup>-fl</sup>	15	<i>Wt1</i> <sup>-fl</sup>
7	<i>Wt1</i> <sup>-fl</sup>	16	<i>Wt1</i> <sup>+fl</sup>
8	<i>Wt1</i> <sup>+fl</sup> ; <i>Cre</i> <sup>TM</sup>	17	(+) <i>Wt1</i> <sup>-fl</sup> ; <i>Cre</i> <sup>TM</sup>
9	<i>Wt1</i> <sup>+fl</sup> ; <i>Cre</i> <sup>TM</sup>	18	(-) ddH <sub>2</sub> O

**Figure 4.3. PCR Analyses of the E13.5 Kidney Rudiments after the *in vivo* Ablation of *Wt1***

The E13.5 embryos were genotyped after 48 hrs of tamoxifen treatment to ablate *Wt1*. Genotypes of these embryos are to the right.




**Figure 4.4. *Meox2* Gene Expression after the *in vivo* Ablation of *Wt1***

In this graph, the y-axis indicates the gene expression *Wt1* and *Meox2* in the *Wt1*<sup>-/ $\Delta$</sup>  kidney rudiments relative to the *Wt1*<sup>+/ $\Delta$</sup>  kidney rudiments. The blue bars represent the control samples *Wt1*<sup>+/flox</sup>; *Cre*<sup>TM</sup> (n=4) and the red bars represents the mutant samples *Wt1*<sup>-flox</sup>; *Cre*<sup>TM</sup> (n=4). After the ablation, *Wt1* is knocked down by 70% (.0004) and *Meox2* gene expression increased by 2X (.018).

### Downstream Effects Due to the *In Vitro* Ablation of *Wt1*


In order to assess the loss of *Wt1* on the gene expression of *Meox2*, other potential downstream genes, and tumorigenic phenotypes, such as over cellular proliferation, inhibition of apoptosis, disruptions in cellular adhesion and migration, *Wt1* was genetically knocked down in the generated F12055 *Wt1*<sup>-fl</sup>; *Immorto* kidney mesenchymal cells by adenoviral infection as described in the **Materials and Methods**. As shown in **Figure 4.6**, *Wt1* was knocked down by ~90% after 48 hrs of infection by qPCR analysis. After 48hrs and 72hrs of infection, there was no significant dysregulation of *Meox2*. After 96 hrs of infection, the gene expression of *Meox2* significantly increased by 61X (0.001) as shown in **Figure 4.5**.

It has been reported that the Tgf- $\beta$ /Smad pathway mediates the activation of *Meox2* expression by Tgf- $\beta$ 1 (147). Additionally, a dominant negative mutant *Smad3* had an effect on the on blocking the endogenous Tgf- $\beta$ /Smad signaling, which resulted in the late induction of *Meox2* mRNA levels (147). Thus, *Tgf- $\beta$ 1* and *Smad3* are known to be involved induction of *Meox2* expression. After 48 hrs of the knockdown of *Wt1*, *Tgf- $\beta$ 1* had a relative gene expression of ~2X (0.039) (**Figure 4.6**). However after 72 hrs and 96 hrs of the knockdown of *Wt1*, *Tgf- $\beta$ 1* expression was knocked down by approximately 40% as shown in **Figure 4.6**. For *Smad3*, its expression was at 1.58X (0.243) after 48 hrs of the knockdown of *Wt1*. *Smad3* expression fluctuated between 72 hrs and 96 hrs. After 72 hrs, *Smad3* expression was knocked down approximately 50%. After 96 hrs after the knockdown of *Wt1*, *Smad3* expression increased to 1.23 (0.799). Overall, this data indicated that the loss of *Wt1* has an effect on *Tgf- $\beta$ 1* expression, but not on *Smad3* expression. Moreover, it also suggest that the upregulation of *Meox2* expression was due to the loss of *Wt1*.



**Figure 4.5. *Meox2* is Upregulated following the *in vitro* Ablation of *Wt1***

In the figure above, the y-axis indicates the gene expression *Wt1* (blue) and *Meox2* (red) in the *Wt1*<sup>-/-</sup>; *Immorto* MMCs relative to the *Wt1*<sup>+/+</sup>; *Immorto* MMCs. The x-axis is a table that provides the values of how much *Wt1* and *Meox2* were relatively expressed. The ablation of *Wt1* in MMCs resulted in the upregulation of *Meox2* expression after 96 hrs.




**Figure 4.6. The Effect of Loss of *Wt1* on *Tgf-β1* and *Smad3* Gene Expression**

In the figure above, the y-axis indicates the gene expression *Wt1* (blue), *Tgfβ-1* (red), and *Smad3* (pink) in the *Wt1*<sup>-/-</sup>; *Immorto* MMCs relative to the *Wt1*<sup>+/+</sup>; *Immorto* MMCs. The x-axis is a table that provides the values of how much *Wt1*, *Tgf-β1*, and *Smad3* were relatively expressed after the ablation of *Wt1* in the MMCs.

## Dysregulation of Meox2 Targeted Genes and Cellular Phenotype

I have demonstrated that *MEOX2* is upregulated following the ablation of *WT1* in WTs, during early kidney development, and in kidney mesenchymal cells. It is known that the upregulation of *Meox2* results in the induction of *p21* and the repression of *Itgb5* (145, 148-150) (151). By qPCR analysis, after 96 hrs of infection, the upregulation of *Meox2* in MMC cells resulted in the significant induction of *p21* (0.0002) and decreased the expression of *Itgb5* (0.01) (**Figure 4.7**). In addition to this, an abnormal cellular phenotype was observed. As shown in **Figure 4.8**, the number of the *Wt1*<sup>-Δ</sup>; *Immorto* MMCs cells on the plate appeared to be less after 72 hrs of infection compared to the control cells. After 96 hrs of infection, there was a significant decrease in the number of adherent cells and an increase of floating cells.


To determine whether these cells were apoptotic, the floating cells were stained with Trypan blue reagent which stains dead cells. Surprisingly, the majority of the floating cells were viable. In **Figure 4.9**, the number of *Wt1*<sup>-Δ</sup>; *Immorto* MMCs was compared to the number of the *Wt1*<sup>-fl</sup>; *Immorto* MMCs by counting the number of cells after they were stained with the Trypan blue reagent. The total number of *Wt1*<sup>-Δ</sup>; *Immorto* MMCs started to decrease after 72hrs (0.080) (**Figure 4.9**). After 96 hrs of infection, there was a significant decrease in the total number of *Wt1*<sup>-Δ</sup>; *Immorto* MMCs compared to the control (0.044), which ~40% of the *Wt1*<sup>-Δ</sup>; *Immorto* MMCs total number of cells were viable and in suspension (**Figure 4.9**).



**Figure 4.7. Induction of *p21* and the Repression of *Itgb5***

In the figure above, the y-axis indicates the gene expression *Meox2* (red), *p21* (black), and *Itgb5* (gray) in the *Wt1*<sup>-/-</sup>; Immorto MMCs relative to the *Wt1*<sup>+/+</sup>; Immorto MMCs. The x-axis is a table that provides the values of how much *Meox2*, *p21*, and *Itgb5* were relatively expressed. After 96hrs of *Wt1* ablation, the overexpression of *Meox2* resulted in the significant induction of *p21* (0.0002) and the knockdown of *Itgb5* (0.01).






**Figure 4.8. The Effect of *Wt1* Ablation in *Wt1*<sup>fllox</sup>; *Immorto* MMCs: Visual Observation in Decrease Cell Number and Loss of Cell Adhesion**

As shown above, the top panel of images is infected cells with Adeno Cre (*Wt1*<sup>Δ</sup>; *Immorto* MMCs) and the bottom panel of images is infected cells with Adeno GFP (*Wt1*<sup>fl</sup>; *Immorto* MMCs) at different time points.

## The Effect of *Wt1* Ablation on Cell Growth and Adhesion



**Figure 4.9. The effect of *Wt1* Ablation in *Wt1*<sup>fllox</sup>; *Immorto* MMCs: Quantification of Cell Proliferation and Adhesion**

In this figure, the red lines indicate the mutant, *Wt1*<sup>-/-</sup>; *Immorto* MMCs and the blue lines indicate the *Wt1*<sup>fl</sup>; *Immorto* MMCs (control). Lines with circle indicate the total number of cells, lines with triangles indicate the number of adherent cells, and lines with squares indicate the number of suspended cells at 48 hrs, 72 hrs, and 96 hrs.

## Discussion

The functional role of *Meox2* has been studied in vascular smooth muscle cells (VSMCs), endothelial cells, and cardiomyocytes. Within these cell types, *Meox2* induces the cyclin-dependent kinase inhibitor, *p21* (145, 148-150). Moreover, *Meox2* has shown to induce apoptosis in VSMCs (150). *Meox2* specifically repressed  $\beta_3$  and  $\beta_5$  integrins in VSMC and may lead to the disruption of integrin-regulated signaling pathways (151). In endothelial cells, *Meox2* inhibits cellular proliferation and endothelial cell tube formation (152). All together, *Meox2* regulates genes that are involved in cellular proliferation and cell adhesion. However, the role of *Meox2* has not been extensively studied during kidney development, in particular, kidney mesenchymal cells.

According to the literature, the mediated-regulation of *Meox2* results from the stimulation of Tgf- $\beta$ 1 (147). It is known that TGF- $\beta$  inhibits cellular proliferation of specific cells types and mutated in various human cancers; TGF- $\beta$  has the functional role of being a tumor suppressors (153-155). *TGF- $\beta$ 2* and *TGF- $\beta$ 3* were up-regulated in mutant WTs in the microarray data (156). Could it be that during kidney development, *TGF- $\beta$ 2* and *TGF- $\beta$ 3* were transcriptionally repressed by *WT1*? During Wilms tumorigenesis, could the loss of *WT1* upregulate the gene expression of *TGF- $\beta$ 2* and *TGF- $\beta$ 3* which resulted in the upregulation of *MEOX2*? TGF- $\beta$ 1 was the only TGF- $\beta$  gene assessed on the array. Therefore, the other TGF- $\beta$  genes need to be assessed following *WT1* mutation as discussed in **Chapter 5**.

Because *Meox2* was expressed at high levels in WTs and during kidney development following a *Wt1* mutation, it may play a role as an oncogene, which

would cause cells to survive and overproliferate. The overexpression of *Meox2* following a *Wt1* mutation resulted in the induction of p21 and the downregulation of *Itgb5*. In addition, there was a decrease in cell number and an increase in the number of floating cells which goes against the concept of an oncogenic phenotype. However, the floating cells were viable and potentially still growing. If these cells were considered tumorigenic and remain viable and proliferative, they could metastasize throughout the body, which would define *Meox2* as an oncogene. However, we do not know whether this cellular phenotype was due to the overexpression of *Meox2* or from the loss of *Wt1*.

Overall, my data supports my hypothesis that the upregulation of *Meox2* is *Wt1*-dependent during early kidney development (*in vivo studies*), especially in the condensed mesenchyme (*in vitro studies*). The dysregulation of downstream targeted genes of *Meox2* followed the upregulation of *Meox2* in the MMCs. A novel cellular phenotype was discovered, in which the cells decreased in number and lose their ability to adhere after the ablation of *Wt1* and/or the upregulation of *Meox2*. Of course there needs to be more analyses done to further investigate the effect of the loss of *Wt1* on *Meox2* gene expression and the role of over expressed *Meox2* in the kidney mesenchymal cells as discussed in **Chapter 5**.

## **Chapter 5: Conclusions and Future Directions**

### **Conclusions**

Wilms tumor is a childhood cancer of the kidney that represents approximately 95% of all pediatric kidney tumors and accounts for 6% of all pediatric cancers (1-4). The classic histology of WT in both humans and mice is triphasic, which consists of mesenchymal, stromal, and epithelial cells. The data from WT histology suggest that there was a disruption during cellular differentiation of the mesenchymal cells during early kidney development. Most of our understanding of the association between WTs and kidney development derived from the investigation of *WT1* and the histology of WTs. However, the role of *WT1* during Wilms tumorigenesis is unclear. *WT1* is a transcriptional factor that is known to activate or repress genes involved in cellular proliferation, differentiation, apoptosis, cellular migration, and cellular adhesion. Therefore, it is hypothesized that the inactivation of *WT1* has a reverse effect on downstream genes that are involved in cellular proliferation, differentiation, apoptosis, cellular migration, and cellular adhesion that potentially lead to Wilms tumorigenesis.

Recently, an Affymetrix GeneChip Human Genome Array was conducted with human with *WT1*-wildtype and *WT1*-mutant WTs. Numerous genes were dysregulated in *WT1*-mutant WTs, which *MEOX2* was one of the significantly dysregulated genes upregulated by 19X in *WT1*-mutant tumors. *MEOX2* was validated by qPCR and remained significantly upregulated in *WT1*-mutant tumor by 41X.

In order to test the hypothesis that *Meox2* is upregulated in a *Wt1*-dependent manner, *Wt1* was genetically ablated. A qPCR analysis was conducted comparing the control kidneys to the *Wt1* ablated kidneys, which resulted in the upregulation of *Meox2*. Since *Wt1* and *Meox2* were both expressed in the mesenchyme, a kidney mesenchymal cells line was generated in which *Wt1* could be genetically ablated with Cre recombinase to determine the effect of *Wt1* mutation on *Meox2* gene expression. Once again, the effect of the ablation of *Wt1* resulted in the significant upregulation of *Meox2*. The significant upregulation of *Meox2* following a *Wt1* mutation, specifically in the mesenchyme, further supported the hypothesis that the upregulation of *Meox2* is *Wt1*-dependent. The inactivation of *Wt1* and/or the upregulation of *Meox2* induced *p21*, repressed *Itgb5* and revealed a unique cellular phenotype in which the cells displayed a loss of cellular adhesion and a decrease in cellular growth. It is not certain whether the induction of *p21* and the repression of *Itgb5* are the cause of the cellular phenotype. However, there is more to investigate how the loss of *Wt1* leads to the formation of WT by first further investigating the actual functional role of *Meox2* during kidney development as later discussed. By further studying the dysregulated genes in WTs, such as *MEOX2*, we can develop therapeutic strategies to prevent and improve treating Wilms tumors. Additionally, we can understand how the remaining 80% of WTs were formed independent of *WT1* mutation.

## Future Directions

Even though the upregulation of *MEOX2* expression following a *WT1* mutation was quantitatively assessed in WT1s during developing kidney at E13.5 and in the mesenchymal cell line, there are still gaps remaining in understanding the mechanistic relationship between WT1 protein and *MEOX2* gene. In order to determine how the Wt1 protein and *Meox2* gene interact, directly or indirectly, a Chromatin Immunoprecipitation (ChIP) sequence would provide evidence that Wt1 is directly regulating the gene expression of *Meox2*.

At E13.5, *Meox2* was significantly upregulated in the kidney rudiments following *Wt1* ablation. To determine whether the loss of *Wt1* affected *Meox2* gene expression throughout kidney development, more developmental time points are required. These studies will provide data to specifically show when and where the upregulation of *Meox2* occur during kidney development after the loss of *Wt1*.

It has been reported that the TGF- $\beta$ /Smad pathway positively mediates the gene expression of *Meox2*. I sought to determine whether Wt1 regulates *Meox2* expression by altering the TGF- $\beta$ /Smad pathway. According to my data, the gene expression of *Tgf- $\beta$ 1* was downregulated when *Wt1* was ablated in the mesenchymal cells. According to our microarray data, TGF- $\beta$ 2 and TGF- $\beta$ 3 were both upregulated in the *Wt1*-mutant WT1s. Thus, TGF- $\beta$ 2 and TGF- $\beta$ 3 could also regulate the gene expression of *MEOX2*. For future experiments, a qPCR will be conducted to analyze whether Wt1 dysregulates Tgf- $\beta$ 2 or Tgf- $\beta$ 3 in kidney rudiments and in the MMC cell lines that could lead to upregulation of *Meox2*. If either of these Tgf- $\beta$  proteins are upregulated after the ablation of *Wt1*, it would suggest two possibilities: 1) that Wt1

represses the expression Tgf- $\beta$ 2 and Tgf- $\beta$ 3 and/or 2) that the regulation of *Meox2* is dependent on the isoforms of Tgf- $\beta$  proteins. The combination of the two possibilities would suggest that *Wt1* is upstream and transcriptionally regulates the TGF- $\beta$ /Smad pathway and *Meox2* expression.

The ablation of *Wt1* is known to slightly induce apoptosis during kidney development (157). In the MMCs, the ablation of *Wt1* after 96hrs resulted in the significant increase of *Meox2*, decrease of adhesive cells, and the decrease in cell growth. The floating cells were viable, however, they could be undergoing early induction of apoptosis. In the future, this experiment needs to be repeated and extended to determine whether these cells will eventually become apoptotic by Fluorescence Activated Cell Sorting (FACS) or transferase dUTP nick end labeling (TUNEL) assay. In addition, testing different cellular adhesion coatings, such as gelatin, collagen, fibronectin, poly-D-lysine, or poly-L-lysine, with *Wt1* ablated cells may reduce the loss of cellular adhesion.

In order to assess whether the decrease in cell growth and loss of cellular adhesion was solely achieved by the upregulation of *Meox2*, future experiments would also include overexpressing *Meox2* independently of *Wt1*. A construct with a pCMV promoter that overexpresses full-length *Meox2* has been generated in the lab. The pCMV-*Meox2* construct could be transiently transfected into a wildtype *Wt1*<sup>+/fl</sup>; *Immorto* MMCs cell line to determine whether these phenotypes resulted solely from the overexpression of *Meox2*.

These future directions will set the stage to study the relationship between *Wt1* and *Meox2* during kidney development, to understand the role of *Meox2* during kidney



development, and to develop strategies to reverse the phenotypes, such as the dysregulation of downstream genes of Meox2, decrease in cell growth and loss of cellular adhesion. Once the relationship between Wt1 and Meox2 are fully understood, we will have an insight on the functional role(s) of WT1 and MEOX2 during Wilms tumorigenesis and have the advantage of being able to develop better therapeutics for WTs.

## Appendix

### Appendix Table 1. Putative Targets of WT1

A, Activate and R, Repress

Gene ID	Protein	Location	Regulation	Reference
<i>ABCB1</i>	ATP-bndg cassette sub-fam B mbr 1 (MDR-1)	7q21.12	R	McCoy, 1999
<i>AMHR2</i>	anti-Mullerian hormone receptor 2	12q13.13	A	Klattig, 2007
<i>AR</i>	androgen receptor	Xq12	R	Shimamura, 1997
<i>AREG</i>	amphiregulin	4q13.3	A	Lee, 1999
<i>BAG3</i>	BCL2-associated athanogene 3	10q26.11	A	Cesaro, 2010
<i>BAK1</i>	BCL2 antagonist/killer 1	6p21.31	A	Morrison, 2005
<i>BCL2</i>	B-cell lymphoma prot 2 beta isoform	18q21.33	A	Mayo, 1999
<i>BCL2A1</i>	BCL2-related protein A1	15q25.1	A	Simpson, 2006
<i>CCNE1</i>	cyclin E1	19q12	R	Loeb, 2002
<i>CCNG1</i>	cyclin G1	5q34	A	Wagner, 2001
<i>CDH1</i>	cadherin 1, type 1; E-cadherin	16q22.1	A	Hosono, 2000
<i>CDH5</i>	cadherin 5, vascular endothelial cadherin	16q21	A	Kirschner, 2010
<i>CDKN1A</i>	cyclin-dependent kinase inhibitor 1A (p21)	6p21.31	A	Englert, 1997
<i>COL4A1</i>	alpha 1 type IV collagen	13q34	A	Hosono, 1999
<i>COL4A2</i>	alpha 2 type IV collagen	13q34	A	Hosono, 1999
<i>CSF1</i>	colony stimulating factor 1	1p13.3	R	Harrington, 1993
<i>CTGF</i>	connective tissue growth factor	6q23.2	R	Stanhope-Baker, 2000
<i>CXXC5</i> (WID)	CXXC-type zinc finger protein 5	5q31.3	A	Kim, 2010
<i>CX3CL1</i>	chemokine (C-X3-C) ligand1	16q13	A	Kim, 2007
<i>DMTF1</i>	cyclin-dependent myb-like protein 1, WT1-induced inhibitor of DSH	7q21.12	R	Tschan, 2008
<i>EGFR</i>	epidermal growth factor receptor	7p11.2	R	Englert, 1995
<i>EGR1</i>	early growth response 1	5q31.2	R	Madden, 1991
<i>EPO</i>	erythropoietin	7q22.1	A	Dame, 2006
<i>EPOR</i>	erythropoietin receptor	19p13.2	A	Kirschner, 2008
<i>EREG</i>	epiregulin	4q13.3	A	Kim, 2007
<i>ESR</i>	estrogen receptor	6q25.1	R	Han, 2008
<i>GNAI2</i>	G-protein alpha i-2	3p21.31	R	Kinane, 1996
<i>HBEGF</i>	heparin-binding EGF-like growth factor	5q31.3	A	Kim, 2007
<i>HOXA10</i>	homeobox gene	7p15.2	R	Andikyan, 2009
<i>HSPA4</i>	heat shock 70kDa protein 4 isoform a	5q31.1	A	Maheswaran, 1998

<i>HSPG2</i>	heparan sulfate proteoglycan 2 (perlecan)	1p36.12	A	Hosono, 1999
<i>IFI16</i>	interferon-inducible protein 16	1q23.1	A	Kim, 2008
<i>IGF1R</i>	insulin-like growth factor 1 receptor	15q26.3	R	Werner, 1993
<i>IGF2</i>	insulin-like growth factor 2	11p15.5	R	Drummond, 1992
<i>IGFBP4</i>	insulin-like growth factor binding prot 4	17q21.2	R	Wagner, 2001
<i>IL10</i>	interleukin-10	1q32.1	A	Sciesielski, 2010
<i>INHHA</i>	inhibin alpha	2q35	R	Hsu, 1995
<i>IRF8</i>	interferon regulatory factor 8	16q24.1	R	Vidovic, 2010
<i>ITGA4</i>	integrin alpha 4	2q31.3	A	Kirschner, 2006
<i>ITGA8</i>	integrin alpha 8	10p13	R	Hosono, 1999
<i>JUNB</i>	Proto-oncogene JunB	19p13.13	A	Kim, 2007
<i>MDK</i>	midkine (neurite growth-promoting factor 2)	11p11.2	R	Adachi, 1996
<i>MIS</i>	Mullerian inhibiting substance	19p13.3	A	Nachtigal, 1998
<i>MKP3</i>	mitogen-activ prot kinase phosph 3 (DUSP6)	12q21.33	A	Morrison, 2008
<i>MMP9</i>	matrix metalloproteinase-9	20q13.12	R	Marcet-Palacios, 2007
<i>MYB</i>	c-myb myeloblastosis viral onc hom	6q23.3	R	McCann, 1995
<i>MYC</i>	c-myc oncogene	8q24.21	A	Han, 2004
<i>MYCN</i>	n-myc myelocytomatosis viral related onc	2p24.3	R	Zhang, 1999
<i>NDRG2</i>	N-myc downstream regulated gene 2	14q11.2	A	Svensson, 2007
<i>NES</i>	nestin	1q23.1	A	Wagner, 2006
<i>NOV</i>	novH, nephroblastoma overexpressed gene	8q24.12	R	Martinerie, 1996
<i>NPHS1</i>	nephrin	19q13.12	A	Wagner, 2004
<i>NR0B1</i>	nucl recep subfamily 0, grp B, mbr 1 (Dax-1)	Xp21.2	A	Kim, 1999
<i>NR5A1/SF1</i>	nuclear receptor subfamily 5, group A (SF-1)	9q33.3	A	Wilhelm, 2002
<i>NTRK2</i>	neurotrophin receptor TrkB	9q21.33	A	Wagner, 2005
<i>ODC</i>	ornithine decarboxylase 1	2p25.1	R	Moshier, 1996
<i>PAX2</i>	paired box protein 2	10q24.31	R	Ryan, 1995
<i>PDGFA</i>	platelet-derived growth factor alpha	7p22	R	Wang, 1992
<i>PODXL</i>	podocalyxin-like (podocalyxin)	7q32.3	A	Palmer, 2001
<i>POU4F2</i>	POU domain, class 4, transc factor 2	4q31.22	A	Wagner, 2003
<i>RARA</i>	retinoic acid receptor, alpha	17q21.2	R	Goodyer, 1995
<i>RBBP7</i>	Rb binding protein 7 (RbAp46)	Xp22.2	A	Guan, 1998
<i>SALL2</i>	Sal-like 2 (HSAL2)	14q11.2	R	Ma, 2001
<i>SCRIB</i>	Scribble	8q24.3	A	Wells, 2010
<i>SDC1</i>	syndecan 1	2p24.1	A	Cook, 1996
<i>SLC6A6</i>	solute carrier fam 6 (neurotrans transp, taurine, TauT)	3p25.1	A	Han, 2003
<i>SNAI1</i>	Snail	20q13.13	A	Martinez-Estrada, 2010

<i>SNAI2</i>	Slug	8q11.21	A	Kim, 2007
<i>SOD1</i>	superoxide dismutase 1	21q22.11	A	Minc, 1999
<i>SOX9</i>	transcription factor SOX9	17q24.3	A	Gao, 2006
<i>SPRY1</i>	Sprouty homolog 1	4q28.1	A	Gross, 2003
<i>SREBP2</i>	sterol reg element binding transc factor	17p11.2	A	Rae, 2004
<i>SRY</i>	sex determining region Y	Yp11.31	A	Matsuzawa-Watanabe, 2003
<i>STAT3</i>	signal transducer and activator of transcription 3	17q21.2	A	Inoue, 1998
<i>STIM1</i>	stromal interaction molecule1	11p15.5	R	Ritchie, 2010
<i>TBXA2R</i>	thromboxane A2 receptor	19p13.3	R	Gannon, 2008
<i>TERT</i>	telomerase reverse transcriptase	5p15.33	R	Oh, 1999
<i>TGFB</i>	transforming growth factor, beta 1	19q13.2	R	Dey, 1994
<i>THBS1</i>	thrombospondin 1	15q14	R	Dejong, 1999
<i>VDR</i>	vit D (1,25- dihydroxyvitamin D3) receptor	12q13.11	A	Wagner, 2001
<i>VEGF</i>	vascular endothelial growth factor	6p21.1	A	Hanson, 2007
<i>WNT4</i>	wingless-type MMTV integ site fam, member 4	1p36.12	A	Sim, 2002
<i>WT1</i>	Wilms tumor 1	11p13	R	Rupprecht, 1994

**A**, Activate and **R**, Repress

**Appendix Table 2.** Calculating the Relative Gene Expression of Validated Genes  
(n=12)

**CXXC4:**

<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>WT-MT</b>	<b>Relative Gene Expression</b>
7.009	7.009	10.856	10.856	-3.223	0.107
6.850	6.850	10.914	10.914		<b>-9.337</b>
7.176	7.176	9.937	9.937		
7.241	7.241	11.008	11.008		
10.029	10.029	12.556	12.556		
9.908	9.908	12.653	12.653		
6.287	6.287	8.368	8.368		
6.102	6.102	8.498	8.498		
6.511	6.511	12.169	12.169		
6.616	6.616	15.053	15.053		
8.126	8.126	8.111	8.111		
7.987	7.987	8.396	8.396		
<b>7.487</b>	<b>1.309</b>	<b>10.710</b>	<b>2.162</b>		

**DDR1:**

<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>WT-MT</b>	<b>Relative Gene Expression</b>
5.978	5.978	5.242	5.242	-0.661	0.633
6.041	6.041	5.241	5.241		-1.581
6.047	6.047	7.374	7.374		
5.976	5.976	7.400	7.400		
5.898	5.898	4.349	4.349		
6.040	6.040	4.279	4.279		
6.771	6.771	5.243	5.243		
6.807	6.807	5.217	5.217		
4.870	4.870	4.008	4.008		
5.000	5.000	4.040	4.040		
5.382	5.382	4.781	4.781		
5.337	5.337	5.046	5.046		
<b>5.846</b>	<b>0.609</b>	<b>5.185</b>	<b>1.132</b>		

**FST:**

<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>WT-MT</b>	<b>Relative Gene Expression</b>
9.966	9.966	6.678	6.678	3.694	12.938
10.074	10.074	6.810	6.810		
12.819	12.819	8.787	8.787		
12.951	12.951	7.781	7.781		
12.666	12.666	7.597	7.597		
13.033	13.033	8.333	8.333		
10.870	10.870	8.305	8.305		
15.658	15.658	9.423	9.423		
14.482	14.482	9.360	9.360		
9.735	9.735	9.119	9.119		
9.804	9.804	9.236	9.236		
<b>12.005</b>	<b>2.042</b>	<b>8.312</b>	<b>0.989</b>		

**HIPK2:**

<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>WT-MT</b>	<b>Relative Gene Expression</b>
3.610	3.610	4.184	4.184	-1.176	0.443
3.629	3.629	4.285	4.285		-2.260
1.780	1.780	4.038	4.038		
1.749	1.749	4.253	4.253		
4.315	4.315	3.109	3.109		
4.473	4.473	3.207	3.207		
2.546	2.546	5.863	5.863		
2.514	2.514	5.953	5.953		
1.880	1.880	2.244	2.244		
1.938	1.938	2.341	2.341		
<b>2.843</b>	<b>1.070</b>	4.341	4.341		
		4.413	4.413		
		<b>4.019</b>	<b>1.168</b>		

**MEOX2:**

<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>WT-MT</b>	<b>Relative Gene Expression</b>
13.369	13.369	5.934	5.934	5.360	41.069
13.473	13.473	6.361	6.361		
12.099	12.099	4.404	4.404		
12.305	12.305	5.087	5.087		
11.029	11.029	5.227	5.227		
10.810	10.810	5.996	5.996		
9.914	9.914	4.608	4.608		
9.179	9.179	5.214	5.214		
13.068	13.068	6.477	6.477		
13.152	13.152	6.621	6.621		
7.082	7.082	6.370	6.370		
7.104	7.104	5.962	5.962		
<b>11.049</b>	<b>2.308</b>	<b>5.689</b>	<b>0.754</b>		

**MLLT3:**

<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>WT-MT</b>	<b>Relative Gene Expression</b>
8.550	8.550	7.819	7.819	3.335	10.091
8.570	8.570	8.714	8.714		
14.635	14.635	7.666	7.666		
15.724	15.724	8.366	8.366		
7.384	7.384	7.832	7.832		
7.705	7.705	7.737	7.737		
12.549	12.549	8.303	8.303		
14.656	14.656	8.488	8.488		
15.321	15.321	10.149	10.149		
12.439	12.439	10.531	10.531		
13.323	13.323	8.562	8.562		
<b>11.896</b>	<b>3.232</b>	8.567	8.567		
		<b>8.561</b>	<b>0.909</b>		


**PRRX1:**

<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>WT-MT</b>	<b>Relative Gene Expression</b>
11.536	11.536	12.771	12.771	3.818	14.099
11.396	11.396	13.306	13.306		
11.826	11.826	6.368	6.368		
15.108	15.108	6.314	6.314		
15.303	15.303	7.699	7.699		
20.143	20.143	8.038	8.038		
11.795	11.795	7.693	7.693		
12.610	12.610	8.246	8.246		
12.896	12.896	9.331	9.331		
9.411	9.411	10.200	10.200		
9.648	9.648	9.129	9.129		
<b>12.879</b>	<b>3.040</b>	9.643	9.643		
		<b>9.062</b>	<b>2.210</b>		




**Appendix Figure 1. IF Images of the Subcloned Kidney MMCs**


**Clone #1:**




**Clone #2:**




**Clone #3:**




F7346 Clone 3 P51



F7346 Clone 3 P51:  
Negative Control




F7346 Clone 3 P51:  
WT1




F7346 Clone 3 P51:  
Vimentin


**Clone #4:**




F7346 Clone 4 P51




F7346 Clone 4 P51




F7346 Clone 4 P52



F7346 Clone 4 P51:  
Negative Control




F7346 Clone 4 P51:  
WT1




F7346 Clone 4 P51:  
Vimentin

**Clone #5:**




**Clone #7:**







**Clone #8:**




F7346 Clone 8 P51




F7346 Clone 8 P51




F7346 Clone 8 P52



F7346 Clone 8 P51:  
Negative Control





F7346 Clone 8 P51:  
WT1




F7346 Clone 8 P51:  
Vimentin


**Clone #11:**




F7346 Clone 11 P51



F7346 Clone 11 P51:  
Negative Control




F7346 Clone 11 P51:  
WT1




F7346 Clone 11 P51:  
Vimentin


**Clone #12:**




F7346 Clone 12 P51




F7346 Clone 12 P51




F7346 Clone 12 P52



F7346 Clone 12 P51:  
Negative Control




F7346 Clone 12 P51:  
WT1




F7346 Clone 12 P51:  
Vimentin


**Clone #15:**




F7346 Clone 15 P51




F7346 Clone 15 P51




F7346 Clone 15 P52



F7346 Clone 15 P51:  
Negative Control




F7346 Clone 15 P51:  
WT1




F7346 Clone 15 P51:  
Vimentin

**Appendix Figure 2: IF images of the MMCs with the *CreER<sup>TM</sup>* Transgene**

**F11249 E3 *Wt1<sup>-fl</sup>*; *Immorto*; *CreER<sup>TM</sup>***



**F11254 E2 *Wt1<sup>-fl</sup>*; *Immorto*; *CreER<sup>TM</sup>***



**Appendix Table 3.** Calculating the relative gene expression of *the Wt1*, *Tgf- $\beta$ 1*, *Smad3*, *Meox2*, *p21* and *Itgb5* after adenoviral infection

**Wt1**

<b>48 Cre</b>		<b>48 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
8.474	8.474	4.705	4.705	-3.224	0.107
7.634	7.634	4.291	4.291		
6.790	6.790	4.231	4.231		
7.633	0.842	4.409	0.258		
<b>72 Cre</b>		<b>72 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
7.059	7.059	5.519	5.519	-3.390	0.095
7.250	7.250	4.647	4.647		
10.483	10.483	4.458	4.458		
8.264	1.924	4.874	0.566		
<b>96 Cre</b>		<b>96 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
4.475	4.475	4.280	4.280	-2.474	0.180
6.599	6.599	3.757	3.757		
8.293	8.293	3.908	3.908		
6.456	1.913	3.982	0.269		

## Tgf- $\beta$ 1

<b>48 Cre</b>		<b>48 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
7.185	7.185	8.021	8.021	0.942	1.921
6.078	6.078	6.605	6.605		
5.966	5.966	7.428	7.428		
6.410	0.674	7.351	0.711		
<b>72 Cre</b>		<b>72 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
6.123	6.123	6.646	6.646	-0.666	0.630
5.858	5.858	4.120	4.120		-1.587
6.600	6.600	5.818	5.818		
6.194	0.376	5.528	1.288		
<b>96 Cre</b>		<b>96 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
6.488	6.488	5.875	5.875	-0.616	0.652
6.186	6.186	5.920	5.920		-1.533
6.421	6.421	5.450	5.450		
6.365	0.159	5.749	0.259		



### Smad3

<b>48 Cre</b>		<b>48 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
8.144	8.144	7.037	7.037	-1.584	0.334
9.148	9.148	7.086	7.086		-2.998
8.646	0.710	7.062	0.034		
<b>72 Cre</b>		<b>72 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
9.384	9.384	8.236	8.236	-0.541	0.687
8.685	8.685	8.808	8.808		-1.455
8.844	8.844	8.246	8.246		
8.971	0.366	8.430	0.328		
<b>96 Cre</b>		<b>96 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
9.549	9.549	7.396	7.396	-1.227	0.427
9.143	9.143	7.643	7.643		-2.341
8.425	8.425	8.397	8.397		
9.039	0.570	7.812	0.522		

## Meox2

<b>48 Cre</b>		<b>48 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
11.250	11.250	11.392	11.392	0.494	1.409
11.435	11.435	12.061	12.061		
11.014	11.014	11.730	11.730		
11.233	0.211	11.728	0.335		
<b>72 Cre</b>		<b>72 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
8.186	8.186	6.156	6.156	-1.028	0.490
7.137	7.137	7.120	7.120		-2.039
7.370	7.370	6.333	6.333		
7.564	0.551	6.536	0.513		
<b>96 Cre</b>		<b>96 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
7.070	7.070	12.489	12.489	5.936	61.204
7.338	7.338	13.589	13.589		
6.953	6.953	13.088	13.088		
7.120	0.197	13.056	0.551		

p21

<b>48 Cre</b>		<b>48 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
5.319	5.319	5.705	5.705	0.366	1.289
5.192	5.192	5.478	5.478		
4.984	4.984	5.410	5.410		
5.165	0.169	5.531	0.155		
<b>72 Cre</b>		<b>72 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
4.848	4.848	5.507	5.507	0.367	1.290
4.824	4.824	5.331	5.331		
5.056	5.056	4.992	4.992		
4.909	0.127	5.277	0.262		
<b>96 Cre</b>		<b>96 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
4.945	4.945	6.272	6.272	1.294	2.452
5.278	5.278	6.575	6.575		
5.141	5.141	6.399	6.399		
5.121	0.167	6.415	0.152		

## Itgb5

<b>48 Cre</b>		<b>48 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
7.532	7.532	8.897	8.897	0.436	1.353
7.782	7.782	7.623	7.623		
7.861	7.861	7.961	7.961		
7.725	0.172	8.160	0.660		
<b>72 Cre</b>		<b>72 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
8.833	8.833	7.327	7.327	-0.911	0.532
8.909	8.909	8.036	8.036		-1.880
8.765	8.765	8.411	8.411		
8.836	0.072	7.925	0.551		
<b>96 Cre</b>		<b>96 GFP</b>			
<b>Average</b>	<b>STDEV</b>	<b>Average</b>	<b>STDEV</b>	<b>Control-Infected</b>	<b>Relative Gene Expr</b>
8.485	8.485	7.234	7.234	-1.100	0.467
8.559	8.559	7.674	7.674		-2.143
8.507	8.507	7.345	7.345		
8.517	0.038	7.417	0.229		

## **Bibliography**

1. Huff, V. 1998. Wilms tumor genetics. *Am J Med Genet* 79:260-267.
2. McLean, T. W., and K. S. Buckley. Pediatric genitourinary tumors. *Curr Opin Oncol* 22:268-273.
3. Davidoff, A. M. 2009. Wilms' tumor. *Curr Opin Pediatr* 21:357-364.
4. Ehrlich, P. F. 2009. Bilateral Wilms' tumor: the need to improve outcomes. *Expert Rev Anticancer Ther* 9:963-973.
5. Breslow, N., J. B. Beckwith, M. Ciol, and K. Sharples. 1988. Age distribution of Wilms' tumor: report from the National Wilms' Tumor Study. *Cancer Res* 48:1653-1657.
6. Sebire, N. J., and G. M. Vujanic. 2009. Paediatric renal tumours: recent developments, new entities and pathological features. *Histopathology* 54:516-528.
7. Rivera, M. N., and D. A. Haber. 2005. Wilms' tumour: connecting tumorigenesis and organ development in the kidney. *Nat Rev Cancer* 5:699-712.
8. Dressler, G. R. 1999. Kidney development branches out. *Dev Genet* 24:189-193.
9. Costantini, F., and R. Kopan. Patterning a complex organ: branching morphogenesis and nephron segmentation in kidney development. *Dev Cell* 18:698-712.
10. Rossant, J. a. T., P. P. L. 2002. *Mouse Development: Patterning, Morphogenesis, and Organogenesis*. Academic Press, San Diego.

11. Mizuno, T., and S. Yasugi. 1990. Susceptibility of epithelia to directive influences of mesenchymes during organogenesis: uncoupling of morphogenesis and cytodifferentiation. *Cell Differ Dev* 31:151-159.
12. Kuure, S., R. Vuolteenaho, and S. Vainio. 2000. Kidney morphogenesis: cellular and molecular regulation. *Mech Dev* 92:31-45.
13. Knudson, A. G., Jr., and L. C. Strong. 1972. Mutation and cancer: a model for Wilms' tumor of the kidney. *J Natl Cancer Inst* 48:313-324.
14. Bown, N., S. J. Cotterill, P. Roberts, M. Griffiths, S. Larkins, S. Hibbert, H. Middleton, A. Kelsey, D. Tritton, and C. Mitchell. 2002. Cytogenetic abnormalities and clinical outcome in Wilms tumor: a study by the U.K. cancer cytogenetics group and the U.K. Children's Cancer Study Group. *Med Pediatr Oncol* 38:11-21.
15. Dome, J. S., and M. J. Coppes. 2002. Recent advances in Wilms tumor genetics. *Curr Opin Pediatr* 14:5-11.
16. Gow, K. W., and J. J. Murphy. 2002. Cytogenetic and histologic findings in Wilms' tumor. *J Pediatr Surg* 37:823-827.
17. Riccardi, V. M., E. Sujansky, A. C. Smith, and U. Francke. 1978. Chromosomal imbalance in the Aniridia-Wilms' tumor association: 11p interstitial deletion. *Pediatrics* 61:604-610.
18. Rahman, N., L. Arbour, P. Tonin, J. Renshaw, J. Pelletier, S. Baruchel, K. Pritchard-Jones, M. R. Stratton, and S. A. Narod. 1996. Evidence for a familial Wilms' tumour gene (FWT1) on chromosome 17q12-q21. *Nat Genet* 13:461-463.

19. McDonald, J. M., E. C. Douglass, R. Fisher, C. F. Geiser, C. E. Krill, L. C. Strong, D. Virshup, and V. Huff. 1998. Linkage of familial Wilms' tumor predisposition to chromosome 19 and a two-locus model for the etiology of familial tumors. *Cancer Res* 58:1387-1390.
20. Koufos, A., M. F. Hansen, B. C. Lampkin, M. L. Workman, N. G. Copeland, N. A. Jenkins, and W. K. Cavenee. 1984. Loss of alleles at loci on human chromosome 11 during genesis of Wilms' tumour. *Nature* 309:170-172.
21. Orkin, S. H., D. S. Goldman, and S. E. Sallan. 1984. Development of homozygosity for chromosome 11p markers in Wilms' tumour. *Nature* 309:172-174.
22. Reeve, A. E., P. J. Housiaux, R. J. Gardner, W. E. Chewings, R. M. Grindley, and L. J. Millow. 1984. Loss of a Harvey ras allele in sporadic Wilms' tumour. *Nature* 309:174-176.
23. Kaneko, Y., C. Homma, N. Maseki, M. Sakurai, and J. Hata. 1991. Correlation of chromosome abnormalities with histological and clinical features in Wilms' and other childhood renal tumors. *Cancer Res* 51:5937-5942.
24. Call, K. M., T. Glaser, C. Y. Ito, A. J. Buckler, J. Pelletier, D. A. Haber, E. A. Rose, A. Kral, H. Yeger, W. H. Lewis, and et al. 1990. Isolation and characterization of a zinc finger polypeptide gene at the human chromosome 11 Wilms' tumor locus. *Cell* 60:509-520.
25. Takeuchi, S., C. R. Bartram, R. Ludwig, B. Royer-Pokora, S. Schneider, J. Imamura, and H. P. Koeffler. 1995. Mutations of p53 in Wilms' tumors. *Mod Pathol* 8:483-487.

26. Bardeesy, N., J. B. Beckwith, and J. Pelletier. 1995. Clonal expansion and attenuated apoptosis in Wilms' tumors are associated with p53 gene mutations. *Cancer Res* 55:215-219.
27. Koesters, R., R. Ridder, A. Kopp-Schneider, D. Betts, V. Adams, F. Niggli, J. Briner, and M. von Knebel Doeberitz. 1999. Mutational activation of the beta-catenin proto-oncogene is a common event in the development of Wilms' tumors. *Cancer Res* 59:3880-3882.
28. Maiti, S., R. Alam, C. I. Amos, and V. Huff. 2000. Frequent association of beta-catenin and WT1 mutations in Wilms tumors. *Cancer Res* 60:6288-6292.
29. Li, C. M., C. E. Kim, A. A. Margolin, M. Guo, J. Zhu, J. M. Mason, T. W. Hensle, V. V. Murty, P. E. Grundy, E. R. Fearon, V. D'Agati, J. D. Licht, and B. Tycko. 2004. CTNNB1 mutations and overexpression of Wnt/beta-catenin target genes in WT1-mutant Wilms' tumors. *Am J Pathol* 165:1943-1953.
30. Fukuzawa, R., R. W. Heathcott, H. E. More, and A. E. Reeve. 2007. Sequential WT1 and CTNNB1 mutations and alterations of beta-catenin localisation in intralobar nephrogenic rests and associated Wilms tumours: two case studies. *J Clin Pathol* 60:1013-1016.
31. Rivera, M. N., W. J. Kim, J. Wells, D. R. Driscoll, B. W. Brannigan, M. Han, J. C. Kim, A. P. Feinberg, W. L. Gerald, S. O. Vargas, L. Chin, A. J. Iafrate, D. W. Bell, and D. A. Haber. 2007. An X chromosome gene, WTX, is commonly inactivated in Wilms tumor. *Science* 315:642-645.
32. Ruteshouser, E. C., S. M. Robinson, and V. Huff. 2008. Wilms tumor genetics: mutations in WT1, WTX, and CTNNB1 account for only about one-third of tumors. *Genes Chromosomes Cancer* 47:461-470.



33. Perotti, D., B. Gamba, M. Sardella, F. Spreafico, M. Terenziani, P. Collini, A. Pession, M. Nantron, F. Fossati-Bellani, and P. Radice. 2008. Functional inactivation of the WTX gene is not a frequent event in Wilms' tumors. *Oncogene* 27:4625-4632.
34. Fukuzawa, R., S. K. Holman, C. W. Chow, R. Savarirayan, A. E. Reeve, and S. P. Robertson. 2010. WTX mutations can occur both early and late in the pathogenesis of Wilms tumour. *J Med Genet*.
35. Gessler, M., A. Poustka, W. Cavenee, R. L. Neve, S. H. Orkin, and G. A. Bruns. 1990. Homozygous deletion in Wilms tumours of a zinc-finger gene identified by chromosome jumping. *Nature* 343:774-778.
36. Haber, D. A., R. L. Sohn, A. J. Buckler, J. Pelletier, K. M. Call, and D. E. Housman. 1991. Alternative splicing and genomic structure of the Wilms tumor gene WT1. *Proc Natl Acad Sci U S A* 88:9618-9622.
37. Armstrong JF, P.-J. K., Bickmore WA, Hastie ND, Bard JB 1992. The expression of the Wilms' tumour gene, WT1, in the developing mammalian embryo. *Mech Dev* 40:85-97.
38. Pritchard-Jones, K., S. Fleming, D. Davidson, W. Bickmore, D. Porteous, C. Gosden, J. Bard, A. Buckler, J. Pelletier, D. Housman, and et al. 1990. The candidate Wilms' tumour gene is involved in genitourinary development. *Nature* 346:194-197.
39. Rauscher, F. J., 3rd. 1993. The WT1 Wilms tumor gene product: a developmentally regulated transcription factor in the kidney that functions as a tumor suppressor. *FASEB J* 7:896-903.

40. Hastie, N. D. 1994. The genetics of Wilms' tumor--a case of disrupted development. *Annu Rev Genet* 28:523-558.
41. Vainio, S., and Y. Lin. 2002. Coordinating early kidney development: lessons from gene targeting. *Nat Rev Genet* 3:533-543.
42. Brunskill, E. W., B. J. Aronow, K. Georgas, B. Rumballe, M. T. Valerius, J. Aronow, V. Kaimal, A. G. Jegga, J. Yu, S. Grimmond, A. P. McMahon, L. T. Patterson, M. H. Little, and S. S. Potter. 2008. Atlas of gene expression in the developing kidney at microanatomic resolution. *Dev Cell* 15:781-791.
43. Kreidberg, J. A., H. Sariola, J. M. Loring, M. Maeda, J. Pelletier, D. Housman, and R. Jaenisch. 1993. WT-1 is required for early kidney development. *Cell* 74:679-691.
44. Hu, Q., F. Gao, W. Tian, E. C. Ruteshouser, Y. Wang, A. Lazar, J. Stewart, L. C. Strong, R. R. Behringer, and V. Huff. Wt1 ablation and Igf2 upregulation in mice result in Wilms tumors with elevated ERK1/2 phosphorylation. *J Clin Invest* 121:174-183.
45. Buckler, A. J., J. Pelletier, D. A. Haber, T. Glaser, and D. E. Housman. 1991. Isolation, characterization, and expression of the murine Wilms' tumor gene (WT1) during kidney development. *Mol Cell Biol* 11:1707-1712.
46. Royer-Pokora, B., M. Beier, M. Henzler, R. Alam, V. Schumacher, A. Weirich, and V. Huff. 2004. Twenty-four new cases of WT1 germline mutations and review of the literature: genotype/phenotype correlations for Wilms tumor development. *Am J Med Genet A* 127A:249-257.

47. Huff, V., H. Miwa, D. A. Haber, K. M. Call, D. Housman, L. C. Strong, and G. F. Saunders. 1991. Evidence for WT1 as a Wilms tumor (WT) gene: intragenic germinal deletion in bilateral WT. *Am J Hum Genet* 48:997-1003.
48. Varanasi, R., N. Bardeesy, M. Ghahremani, M. J. Petruzzi, N. Nowak, M. A. Adam, P. Grundy, T. B. Shows, and J. Pelletier. 1994. Fine structure analysis of the WT1 gene in sporadic Wilms tumors. *Proc Natl Acad Sci U S A* 91:3554-3558.
49. Gessler, M., A. Konig, K. Arden, P. Grundy, S. Orkin, S. Sallan, C. Peters, S. Ruyle, J. Mandell, F. Li, and et al. 1994. Infrequent mutation of the WT1 gene in 77 Wilms' Tumors. *Hum Mutat* 3:212-222.
50. Huff, V., N. Jaffe, G. F. Saunders, L. C. Strong, F. Villalba, and E. C. Ruteshouser. 1995. WT1 exon 1 deletion/insertion mutations in Wilms tumor patients, associated with di- and trinucleotide repeats and deletion hotspot consensus sequences. *Am J Hum Genet* 56:84-90.
51. Huff, V. Wilms' tumours: about tumour suppressor genes, an oncogene and a chameleon gene. *Nat Rev Cancer* 11:111-121.
52. McCoy, C., S. B. McGee, and M. M. Cornwell. 1999. The Wilms' tumor suppressor, WT1, inhibits 12-O-tetradecanoylphorbol-13-acetate activation of the multidrug resistance-1 promoter. *Cell Growth Differ* 10:377-386.
53. Klattig, J., R. Sierig, D. Kruspe, B. Besenbeck, and C. Englert. 2007. Wilms' tumor protein Wt1 is an activator of the anti-Mullerian hormone receptor gene *Amhr2*. *Mol Cell Biol* 27:4355-4364.
54. Shimamura, R., G. C. Fraizer, J. Trapman, C. Lau Yf, and G. F. Saunders. 1997. The Wilms' tumor gene WT1 can regulate genes involved in sex

- determination and differentiation: SRY, Mullerian-inhibiting substance, and the androgen receptor. *Clin Cancer Res* 3:2571-2580.
55. Lee, S. B., K. Huang, R. Palmer, V. B. Truong, D. Herzlinger, K. A. Kolquist, J. Wong, C. Paulding, S. K. Yoon, W. Gerald, J. D. Oliner, and D. A. Haber. 1999. The Wilms tumor suppressor WT1 encodes a transcriptional activator of amphiregulin. *Cell* 98:663-673.
  56. Cesaro, E., G. Montano, A. Rosati, R. Crescitelli, P. Izzo, M. C. Turco, and P. Costanzo. WT1 protein is a transcriptional activator of the antiapoptotic bag3 gene. *Leukemia* 24:1204-1206.
  57. Morrison, D. J., M. A. English, and J. D. Licht. 2005. WT1 induces apoptosis through transcriptional regulation of the proapoptotic Bcl-2 family member Bak. *Cancer Res* 65:8174-8182.
  58. Mayo, M. W., C. Y. Wang, S. S. Drouin, L. V. Madrid, A. F. Marshall, J. C. Reed, B. E. Weissman, and A. S. Baldwin. 1999. WT1 modulates apoptosis by transcriptionally upregulating the bcl-2 proto-oncogene. *EMBO J* 18:3990-4003.
  59. Simpson, L. A., E. A. Burwell, K. A. Thompson, S. Shahnaz, A. R. Chen, and D. M. Loeb. 2006. The antiapoptotic gene A1/BFL1 is a WT1 target gene that mediates granulocytic differentiation and resistance to chemotherapy. *Blood* 107:4695-4702.
  60. Loeb, D. M., D. Korz, M. Katsnelson, E. A. Burwell, A. D. Friedman, and S. Sukumar. 2002. Cyclin E is a target of WT1 transcriptional repression. *J Biol Chem* 277:19627-19632.

61. Wagner, K. J., C. E. Patek, C. Miles, S. Christie, A. J. Brookes, and M. L. Hooper. 2001. Truncation of WT1 results in downregulation of cyclin G1 and IGFBP-4 expression. *Biochem Biophys Res Commun* 287:977-982.
62. Hosono, S., I. Gross, M. A. English, K. M. Hajra, E. R. Fearon, and J. D. Licht. 2000. E-cadherin is a WT1 target gene. *J Biol Chem* 275:10943-10953.
63. Kirschner, K. M., L. K. Sciesielski, and H. Scholz. Wilms' tumour protein Wt1 stimulates transcription of the gene encoding vascular endothelial cadherin. *Pflugers Arch* 460:1051-1061.
64. Englert, C., S. Maheswaran, A. J. Garvin, J. Kreidberg, and D. A. Haber. 1997. Induction of p21 by the Wilms' tumor suppressor gene WT1. *Cancer Res* 57:1429-1434.
65. Hosono, S., X. Luo, D. P. Hyink, L. M. Schnapp, P. D. Wilson, C. R. Burrow, J. C. Reddy, G. F. Atweh, and J. D. Licht. 1999. WT1 expression induces features of renal epithelial differentiation in mesenchymal fibroblasts. *Oncogene* 18:417-427.
66. Harrington, M. A., B. Konicek, A. Song, X. L. Xia, W. J. Fredericks, and F. J. Rauscher, 3rd. 1993. Inhibition of colony-stimulating factor-1 promoter activity by the product of the Wilms' tumor locus. *J Biol Chem* 268:21271-21275.
67. Stanhope-Baker, P., and B. R. Williams. 2000. Identification of connective tissue growth factor as a target of WT1 transcriptional regulation. *J Biol Chem* 275:38139-38150.
68. Kim, M. S., S. K. Yoon, F. Bollig, J. Kitagaki, W. Hur, N. J. Whye, Y. P. Wu, M. N. Rivera, J. Y. Park, H. S. Kim, K. Malik, D. W. Bell, C. Englert, A. O.

- Perantoni, and S. B. Lee. A novel Wilms tumor 1 (WT1) target gene negatively regulates the WNT signaling pathway. *J Biol Chem* 285:14585-14593.
69. Kim, H. S., M. S. Kim, A. L. Hancock, J. C. Harper, J. Y. Park, G. Poy, A. O. Perantoni, M. Cam, K. Malik, and S. B. Lee. 2007. Identification of novel Wilms' tumor suppressor gene target genes implicated in kidney development. *J Biol Chem* 282:16278-16287.
  70. Tschan, M. P., U. Gullberg, D. Shan, B. E. Torbett, M. F. Fey, and A. Tobler. 2008. The hDMP1 tumor suppressor is a new WT1 target in myeloid leukemias. *Leukemia* 22:1087-1090.
  71. Englert, C., X. Hou, S. Maheswaran, P. Bennett, C. Ngwu, G. G. Re, A. J. Garvin, M. R. Rosner, and D. A. Haber. 1995. WT1 suppresses synthesis of the epidermal growth factor receptor and induces apoptosis. *Embo J* 14:4662-4675.
  72. Madden, S. L., D. M. Cook, J. F. Morris, A. Gashler, V. P. Sukhatme, and F. J. Rauscher, 3rd. 1991. Transcriptional repression mediated by the WT1 Wilms tumor gene product. *Science* 253:1550-1553.
  73. Dame, C., K. M. Kirschner, K. V. Bartz, T. Wallach, C. S. Hussels, and H. Scholz. 2006. Wilms tumor suppressor, Wt1, is a transcriptional activator of the erythropoietin gene. *Blood* 107:4282-4290.
  74. Han, Y., L. Yang, F. Suarez-Saiz, S. San-Marina, J. Cui, and M. D. Minden. 2008. Wilms' tumor 1 suppressor gene mediates antiestrogen resistance via down-regulation of estrogen receptor-alpha expression in breast cancer cells. *Mol Cancer Res* 6:1347-1355.
  75. Kirschner, K. M., P. Hagen, C. S. Hussels, M. Ballmaier, H. Scholz, and C. Dame. 2008. The Wilms' tumor suppressor Wt1 activates transcription of the

- erythropoietin receptor in hematopoietic progenitor cells. *FASEB J* 22:2690-2701.
76. Kinane, B., Finder, J. B., Kawashima, A., Brown, D., Abbate, M., Fredericks, W.J., Sukhatme, V. P., Rauscher III, F.J., and Ercolani, L. . 1996. Expand+LLC-PK Cell Growth Is Repressed by WT1 Inhibition of G-protein Protooncogene Transcription. *Journal of Biological Chemistry* 270:30760-30764.
77. Andikyan, V., and H. S. Taylor. 2009. WT1 represses HOX gene expression in the regulation of gynaecologic tumour histologic type. *J Cell Mol Med* 13:4522-4531.
78. Maheswaran, S., C. Englert, G. Zheng, S. B. Lee, J. Wong, D. P. Harkin, J. Bean, R. Ezzell, A. J. Garvin, R. T. McCluskey, J. A. DeCaprio, and D. A. Haber. 1998. Inhibition of cellular proliferation by the Wilms tumor suppressor WT1 requires association with the inducible chaperone Hsp70. *Genes Dev* 12:1108-1120.
79. Kim, M. K., J. M. Mason, C. M. Li, W. Berkofsky-Fessler, L. Jiang, D. Choubey, P. E. Grundy, B. Tycko, and J. D. Licht. 2008. A pathologic link between Wilms tumor suppressor gene, WT1, and IFI16. *Neoplasia* 10:69-78.
80. Werner, H., C. T. Roberts, Jr., and D. LeRoith. 1993. The regulation of IGF-I receptor gene expression by positive and negative zinc-finger transcription factors. *Adv Exp Med Biol* 343:91-103.
81. Drummond, I. A., S. L. Madden, P. Rohwer-Nutter, G. I. Bell, V. P. Sukhatme, and F. J. Rauscher, 3rd. 1992. Repression of the insulin-like growth factor II gene by the Wilms tumor suppressor WT1. *Science* 257:674-678.

82. Hsu, S. Y., M. Kubo, S. Y. Chun, F. G. Haluska, D. E. Housman, and A. J. Hsueh. 1995. Wilms' tumor protein WT1 as an ovarian transcription factor: decreases in expression during follicle development and repression of inhibin-alpha gene promoter. *Mol Endocrinol* 9:1356-1366.
83. Vidovic, K., E. Svensson, B. Nilsson, B. Thuresson, T. Olofsson, A. Lennartsson, and U. Gullberg. Wilms' tumor gene 1 protein represses the expression of the tumor suppressor interferon regulatory factor 8 in human hematopoietic progenitors and in leukemic cells. *Leukemia* 24:992-1000.
84. Kirschner, K. M., N. Wagner, K. D. Wagner, S. Wellmann, and H. Scholz. 2006. The Wilms tumor suppressor Wt1 promotes cell adhesion through transcriptional activation of the alpha4integrin gene. *J Biol Chem* 281:31930-31939.
85. Adachi, Y., S. Matsubara, C. Pedraza, M. Ozawa, J. Tsutsui, H. Takamatsu, H. Noguchi, T. Akiyama, and T. Muramatsu. 1996. Midkine as a novel target gene for the Wilms' tumor suppressor gene (WT1). *Oncogene* 13:2197-2203.
86. Nachtigal, M. W., Y. Hirokawa, D. L. Enyeart-VanHouten, J. N. Flanagan, G. D. Hammer, and H. A. Ingraham. 1998. Wilms' tumor 1 and Dax-1 modulate the orphan nuclear receptor SF-1 in sex-specific gene expression. *Cell* 93:445-454.
87. Morrison, D. J., M. K. Kim, W. Berkofsky-Fessler, and J. D. Licht. 2008. WT1 induction of mitogen-activated protein kinase phosphatase 3 represents a novel mechanism of growth suppression. *Mol Cancer Res* 6:1225-1231.
88. Marcet-Palacios, M., M. Ulanova, F. Duta, L. Puttagunta, S. Munoz, D. Gibbins, M. Radomski, L. Cameron, I. Mayers, and A. D. Befus. 2007. The



- transcription factor Wilms tumor 1 regulates matrix metalloproteinase-9 through a nitric oxide-mediated pathway. *J Immunol* 179:256-265.
89. McCann, S., J. Sullivan, J. Guerra, M. Arcinas, and L. M. Boxer. 1995. Repression of the c-myb gene by WT1 protein in T and B cell lines. *J Biol Chem* 270:23785-23789.
  90. Han, Y., S. San-Marina, J. Liu, and M. D. Minden. 2004. Transcriptional activation of c-myc proto-oncogene by WT1 protein. *Oncogene* 23:6933-6941.
  91. Zhang, X., G. Xing, and G. F. Saunders. 1999. Proto-oncogene N-myc promoter is down regulated by the Wilms' tumor suppressor gene WT1. *Anticancer Res* 19:1641-1648.
  92. Svensson, E., K. Vidovic, T. Olofsson, J. Vallon-Christersson, A. Borg, and U. Gullberg. 2007. The Wilms' tumor gene 1 (WT1) induces expression of the N-myc downstream regulated gene 2 (NDRG2). *DNA Cell Biol* 26:589-597.
  93. Wagner, N., K. D. Wagner, H. Scholz, K. M. Kirschner, and A. Schedl. 2006. Intermediate filament protein nestin is expressed in developing kidney and heart and might be regulated by the Wilms' tumor suppressor Wt1. *Am J Physiol Regul Integr Comp Physiol* 291:R779-787.
  94. Martinerie, C., G. Chevalier, F. J. Rauscher, 3rd, and B. Perbal. 1996. Regulation of nov by WT1: a potential role for nov in nephrogenesis. *Oncogene* 12:1479-1492.
  95. Wagner, N., K. D. Wagner, Y. Xing, H. Scholz, and A. Schedl. 2004. The major podocyte protein nephrin is transcriptionally activated by the Wilms' tumor suppressor WT1. *J Am Soc Nephrol* 15:3044-3051.

96. Kim, J., D. Prawitt, N. Bardeesy, E. Torban, C. Vicaner, P. Goodyer, B. Zabel, and J. Pelletier. 1999. The Wilms' tumor suppressor gene (wt1) product regulates Dax-1 gene expression during gonadal differentiation. *Mol Cell Biol* 19:2289-2299.
97. Wilhelm, D., and C. Englert. 2002. The Wilms tumor suppressor WT1 regulates early gonad development by activation of Sf1. *Genes Dev* 16:1839-1851.
98. Wagner, N., K. D. Wagner, H. Theres, C. Englert, A. Schedl, and H. Scholz. 2005. Coronary vessel development requires activation of the TrkB neurotrophin receptor by the Wilms' tumor transcription factor Wt1. *Genes Dev* 19:2631-2642.
99. Moshier, J. A., M. Skunca, W. Wu, S. M. Boppana, F. J. Rauscher, 3rd, and J. Dosesu. 1996. Regulation of ornithine decarboxylase gene expression by the Wilms' tumor suppressor WT1. *Nucleic Acids Res* 24:1149-1157.
100. Ryan, G., V. Steele-Perkins, J. F. Morris, F. J. Rauscher, 3rd, and G. R. Dressler. 1995. Repression of Pax-2 by WT1 during normal kidney development. *Development* 121:867-875.
101. Wang, Z. Y., S. L. Madden, T. F. Deuel, and F. J. Rauscher, 3rd. 1992. The Wilms' tumor gene product, WT1, represses transcription of the platelet-derived growth factor A-chain gene. *J Biol Chem* 267:21999-22002.
102. Palmer, R. E., A. Kotsianti, B. Cadman, T. Boyd, W. Gerald, and D. A. Haber. 2001. WT1 regulates the expression of the major glomerular podocyte membrane protein Podocalyxin. *Curr Biol* 11:1805-1809.

103. Wagner, K. D., N. Wagner, G. Schley, H. Theres, and H. Scholz. 2003. The Wilms' tumor suppressor Wt1 encodes a transcriptional activator of the class IV POU-domain factor Pou4f2 (Brn-3b). *Gene* 305:217-223.
104. Goodyer, P., M. Dehbi, E. Torban, W. Bruening, and J. Pelletier. 1995. Repression of the retinoic acid receptor-alpha gene by the Wilms' tumor suppressor gene product, wt1. *Oncogene* 10:1125-1129.
105. Guan, L. S., M. Rauchman, and Z. Y. Wang. 1998. Induction of Rb-associated protein (RbAp46) by Wilms' tumor suppressor WT1 mediates growth inhibition. *J Biol Chem* 273:27047-27050.
106. Ma, Y., D. Li, L. Chai, A. M. Luciani, D. Ford, J. Morgan, and A. L. Maizel. 2001. Cloning and characterization of two promoters for the human HSAL2 gene and their transcriptional repression by the Wilms tumor suppressor gene product. *J Biol Chem* 276:48223-48230.
107. Wells, J., M. N. Rivera, W. J. Kim, K. Starbuck, and D. A. Haber. The predominant WT1 isoform (+KTS) encodes a DNA-binding protein targeting the planar cell polarity gene *Scribble* in renal podocytes. *Mol Cancer Res* 8:975-985.
108. Cook, D. M., M. T. Hinkes, M. Bernfield, and F. J. Rauscher, 3rd. 1996. Transcriptional activation of the syndecan-1 promoter by the Wilms' tumor protein WT1. *Oncogene* 13:1789-1799.
109. Han, X., and R. W. Chesney. 2003. Regulation of taurine transporter gene (TauT) by WT1. *FEBS Lett* 540:71-76.
110. Martinez-Estrada, O. M., L. A. Lettice, A. Essafi, J. A. Guadix, J. Slight, V. Velecela, E. Hall, J. Reichmann, P. S. Devenney, P. Hohenstein, N. Hosen, R.

- E. Hill, R. Munoz-Chapuli, and N. D. Hastie. Wt1 is required for cardiovascular progenitor cell formation through transcriptional control of Snail and E-cadherin. *Nat Genet* 42:89-93.
111. Minc, E., P. de Coppet, P. Masson, L. Thiery, S. Dutertre, M. Amor-Gueret, and C. Jaulin. 1999. The human copper-zinc superoxide dismutase gene (SOD1) proximal promoter is regulated by Sp1, Egr-1, and WT1 via non-canonical binding sites. *J Biol Chem* 274:503-509.
112. Gao, F., S. Maiti, N. Alam, Z. Zhang, J. M. Deng, R. R. Behringer, C. Lecureuil, F. Guillou, and V. Huff. 2006. The Wilms tumor gene, Wt1, is required for Sox9 expression and maintenance of tubular architecture in the developing testis. *Proc Natl Acad Sci U S A* 103:11987-11992.
113. Gross, I., D. J. Morrison, D. P. Hyink, K. Georgas, M. A. English, M. Mericskay, S. Hosono, D. Sassoon, P. D. Wilson, M. Little, and J. D. Licht. 2003. The receptor tyrosine kinase regulator Sprouty1 is a target of the tumor suppressor WT1 and important for kidney development. *J Biol Chem* 278:41420-41430.
114. Rae, F. K., G. Martinez, K. R. Gillinder, A. Smith, G. Shooter, A. R. Forrest, S. M. Grimmond, and M. H. Little. 2004. Analysis of complementary expression profiles following WT1 induction versus repression reveals the cholesterol/fatty acid synthetic pathways as a possible major target of WT1. *Oncogene* 23:3067-3079.
115. Matsuzawa-Watanabe, Y., J. Inoue, and K. Semba. 2003. Transcriptional activity of testis-determining factor SRY is modulated by the Wilms' tumor 1 gene product, WT1. *Oncogene* 22:7900-7904.

116. Inoue, K., H. Tamaki, H. Ogawa, Y. Oka, T. Soma, T. Tatekawa, Y. Oji, A. Tsuboi, E. H. Kim, M. Kawakami, T. Akiyama, T. Kishimoto, and H. Sugiyama. 1998. Wilms' tumor gene (WT1) competes with differentiation-inducing signal in hematopoietic progenitor cells. *Blood* 91:2969-2976.
117. Ritchie, M. F., C. Yue, Y. Zhou, P. J. Houghton, and J. Soboloff. Wilms tumor suppressor 1 (WT1) and early growth response 1 (EGR1) are regulators of STIM1 expression. *J Biol Chem* 285:10591-10596.
118. Gannon, A. M., E. C. Turner, H. M. Reid, and B. T. Kinsella. 2009. Regulated expression of the alpha isoform of the human thromboxane A2 receptor during megakaryocyte differentiation: a coordinated role for WT1, Egr1, and Sp1. *J Mol Biol* 394:29-45.
119. Oh, S., Y. Song, J. Yim, and T. K. Kim. 1999. The Wilms' tumor 1 tumor suppressor gene represses transcription of the human telomerase reverse transcriptase gene. *J Biol Chem* 274:37473-37478.
120. Dey, B. R., V. P. Sukhatme, A. B. Roberts, M. B. Sporn, F. J. Rauscher, 3rd, and S. J. Kim. 1994. Repression of the transforming growth factor-beta 1 gene by the Wilms' tumor suppressor WT1 gene product. *Mol Endocrinol* 8:595-602.
121. Dejong, V., A. Degeorges, S. Filleur, S. Ait-Si-Ali, A. Mettouchi, P. Bornstein, B. Binetruy, and F. Cabon. 1999. The Wilms' tumor gene product represses the transcription of thrombospondin 1 in response to overexpression of c-Jun. *Oncogene* 18:3143-3151.
122. Hanson, J., J. Gorman, J. Reese, and G. Fraizer. 2007. Regulation of vascular endothelial growth factor, VEGF, gene promoter by the tumor suppressor, WT1. *Front Biosci* 12:2279-2290.

123. Sim, E. U., A. Smith, E. Szilagi, F. Rae, P. Ioannou, M. H. Lindsay, and M. H. Little. 2002. Wnt-4 regulation by the Wilms' tumour suppressor gene, WT1. *Oncogene* 21:2948-2960.
124. Rupprecht, H. D., I. A. Drummond, S. L. Madden, F. J. Rauscher, 3rd, and V. P. Sukhatme. 1994. The Wilms' tumor suppressor gene WT1 is negatively autoregulated. *J Biol Chem* 269:6198-6206.
125. Rauscher, F. J., 3rd, J. F. Morris, O. E. Tournay, D. M. Cook, and T. Curran. 1990. Binding of the Wilms' tumor locus zinc finger protein to the EGR-1 consensus sequence. *Science* 250:1259-1262.
126. Wang, Z. Y., Q. Q. Qiu, J. Huang, M. Gurrieri, and T. F. Deuel. 1995. Products of alternatively spliced transcripts of the Wilms' tumor suppressor gene, wt1, have altered DNA binding specificity and regulate transcription in different ways. *Oncogene* 10:415-422.
127. Ashburner, M., C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A. Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin, and G. Sherlock. 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25:25-29.
128. Ruteshouser, E. C., B. W. Hendrickson, S. Colella, R. Krahe, L. Pinto, and V. Huff. 2005. Genome-wide loss of heterozygosity analysis of WT1-wild-type and WT1-mutant Wilms tumors. *Genes Chromosomes Cancer* 43:172-180.
129. Jogi, A., F. Abel, R. M. Sjoberg, R. Toftgard, P. G. Zaphiropoulos, S. Pahlman, T. Martinsson, and H. Axelson. 2000. Patched 2, located in 1p32-34, is not mutated in high stage neuroblastoma tumors. *Int J Oncol* 16:943-949.

130. Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156-159.
131. Candia, A. F., J. Hu, J. Crosby, P. A. Lalley, D. Noden, J. H. Nadeau, and C. V. Wright. 1992. Mox-1 and Mox-2 define a novel homeobox gene subfamily and are differentially expressed during early mesodermal patterning in mouse embryos. *Development* 116:1123-1136.
132. Candia, A. F., and C. V. Wright. 1996. Differential localization of Mox-1 and Mox-2 proteins indicates distinct roles during development. *Int J Dev Biol* 40:1179-1184.
133. Scott, M. P., J. W. Tamkun, and G. W. Hartzell, 3rd. 1989. The structure and function of the homeodomain. *Biochim Biophys Acta* 989:25-48.
134. Gehring, W. J., M. Affolter, and T. Burglin. 1994. Homeodomain proteins. *Annu Rev Biochem* 63:487-526.
135. Graham, F. L., J. Smiley, W. C. Russell, and R. Nairn. 1977. Characteristics of a human cell line transformed by DNA from human adenovirus type 5. *J Gen Virol* 36:59-74.
136. Jat, P. S., M. D. Noble, P. Ataliotis, Y. Tanaka, N. Yannoutsos, L. Larsen, and D. Kioussis. 1991. Direct derivation of conditionally immortal cell lines from an H-2Kb-tsA58 transgenic mouse. *Proc Natl Acad Sci U S A* 88:5096-5100.
137. Danielian, P. S., R. White, S. A. Hoare, S. E. Fawell, and M. G. Parker. 1993. Identification of residues in the estrogen receptor that confer differential sensitivity to estrogen and hydroxytamoxifen. *Mol Endocrinol* 7:232-240.

138. Fawell, S. E., J. A. Lees, R. White, and M. G. Parker. 1990. Characterization and colocalization of steroid binding and dimerization activities in the mouse estrogen receptor. *Cell* 60:953-962.
139. Littlewood, T. D., D. C. Hancock, P. S. Danielian, M. G. Parker, and G. I. Evan. 1995. A modified oestrogen receptor ligand-binding domain as an improved switch for the regulation of heterologous proteins. *Nucleic Acids Res* 23:1686-1690.
140. Mattioni, T., J. F. Louvion, and D. Picard. 1994. Regulation of protein activities by fusion to steroid binding domains. *Methods Cell Biol* 43 Pt A:335-352.
141. Picard, D. 1994. Regulation of protein function through expression of chimaeric proteins. *Curr Opin Biotechnol* 5:511-515.
142. Boyle, S., A. Misfeldt, K. Chandler, K. Deal, E. Southard-Smith, D. Mortlock, H. Baldwin, and M. de Caestecker. 2008. Fate mapping using Cited1-CreER mice demonstrates that the cap mesenchyme contains self-renewing progenitor cells and gives rise exclusively to nephronic epithelia. *Development Biology* 313:234-245.
143. Boyle, S., T. Shioda, A. Perantoni, and M. de Caestecker. 2007. Cited1 and Cited2 are differentially expressed in the developing kidney but are not required for nephrogenesis. *Development Dynamics* 236:2321-2330.
144. LePage, D. F., D. A. Altomare, J. R. Testa, and K. Walsh. 1994. Molecular cloning and localization of the human GAX gene to 7p21. *Genomics* 24:535-540.
145. Gorski, D. H., and K. Walsh. 2000. The role of homeobox genes in vascular remodeling and angiogenesis. *Circ Res* 87:865-872.



146. Chemistry, M.-P.-I. o. B. 2005. Genepaint.org.
147. Valcourt, U., S. Thuault, K. Pardali, C. H. Heldin, and A. Moustakas. 2007. Functional role of Meox2 during the epithelial cytostatic response to TGF-beta. *Mol Oncol* 1:55-71.
148. Gorski, D. H., D. F. LePage, C. V. Patel, N. G. Copeland, N. A. Jenkins, and K. Walsh. 1993. Molecular cloning of a diverged homeobox gene that is rapidly down-regulated during the G0/G1 transition in vascular smooth muscle cells. *Mol Cell Biol* 13:3722-3733.
149. Smith, R. C., D. Branellec, D. H. Gorski, K. Guo, H. Perlman, J. F. Dedieu, C. Pastore, A. Mahfoudi, P. Deneffe, J. M. Isner, and K. Walsh. 1997. p21CIP1-mediated inhibition of cell proliferation by overexpression of the gax homeodomain gene. *Genes Dev* 11:1674-1689.
150. Perlman, H., Z. Luo, K. Krasinski, A. Le Roux, A. Mahfoudi, R. C. Smith, D. Branellec, and K. Walsh. 1999. Adenovirus-mediated delivery of the Gax transcription factor to rat carotid arteries inhibits smooth muscle proliferation and induces apoptosis. *Gene Ther* 6:758-763.
151. Witzenbichler, B., Y. Kureishi, Z. Luo, A. Le Roux, D. Branellec, and K. Walsh. 1999. Regulation of smooth muscle cell migration and integrin expression by the Gax transcription factor. *J Clin Invest* 104:1469-1480.
152. Gorski, D. H., and A. J. Leal. 2003. Inhibition of endothelial cell activation by the homeobox gene Gax. *J Surg Res* 111:91-99.
153. Ten Dijke, P., M. J. Goumans, F. Itoh, and S. Itoh. 2002. Regulation of cell proliferation by Smad proteins. *J Cell Physiol* 191:1-16.

154. Siegel, P. M., and J. Massague. 2003. Cytostatic and apoptotic actions of TGF-beta in homeostasis and cancer. *Nat Rev Cancer* 3:807-821.
155. Pardali, K., and A. Moustakas. 2007. Actions of TGF-beta as tumor suppressor and pro-metastatic factor in human cancer. *Biochim Biophys Acta* 1775:21-62.
156. Udtha, M., S. J. Lee, R. Alam, K. Coombes, and V. Huff. 2003. Upregulation of c-MYC in WT1-mutant tumors: assessment of WT1 putative transcriptional targets using cDNA microarray expression profiling of genetically defined Wilms' tumors. *Oncogene* 22:3821-3826.
157. Hu, Q., Gao, F., Tian, W., Ruteshouser, E. C., Wang, Y., Lazar, A., Stewart, J., Strong, L., Behringer, R., and Huff, V. . 2010 (In Press). Wt1 Ablation and Igf2-Up-regulation in Mice Result in Wilms Tumors with Activated pERK1/2. *The Journal of Clinical Investigation*.

## **Vitae**

LaGina Merie Nosavanh was born on May 15, 1985 in Salisbury, Maryland and raised in Frankford, Delaware. She attended Delaware State University located in Dover, Delaware beginning in 2003. LaGina received a NIH MARC (Minority Access to Research Careers) undergraduate training grant, and participated in the summer research programs at the following institutions: University of Texas – Health Science Center at Houston (2004), Stanford University (2005), and University of Michigan – Ann Arbor (2006). LaGina gave a poster presentation at the ABRMS (Annual Biomedical Conference for Minority Students). She conducted research in Dr. Fatma Helmy's laboratory at Delaware State University from Fall 2005 - Spring 2007. She presented her data at the Experimental Biology Conference. LaGina graduated Magna Cum Laude with a B.S. in Biology in 2007 and continued her education in Genetics at the University of Texas – Graduate School for Biomedical Sciences. In 2008, she joined Dr. Vicki Huff's laboratory, and gained research experience in development biology, pediatric cancer biology/genetics and molecular genetics to address the important questions in kidney development and Wilms tumorigenesis. LaGina attended the AACR (American Association for Cancer Research) National meeting to gain more knowledge in cancer research in 2009 at Denver Colorado. She presented a poster at the Genes and Development Spring Retreat in 2009 at Port Aransas, Texas. She also presented a poster at the 2009 Molecular Carcinogenesis and Genes and Development Joint Retreat at M.D. Anderson South Campus. In 2009, she received the Gigli Family Endowed Scholarship for 2009-2010. LaGina was granted a Masters in Genes and Development in December of 2011.

Her abstracts include:

**Nosavanh, L. M.**, Ruteshouser, C. and Huff, V. Genes Dysregulated Following *WT1* Mutation and their Role in Wilms Tumorigenesis and Nephrogenesis. Abstract for poster presentation, Genes and Development Annual Retreat, 2009

**Nosavanh, L. M.**, and Huff, V. Genes Dysregulated Following *WT1* Mutation and their Role in Wilms Tumorigenesis and Nephrogenesis. Abstract for poster presentation, Genes and Development and Molecular Carcinogenesis Joint Retreat, 2009

Helmy, F., **Nosavanh, L.**, Haynes, H. and Juracka, A. Phospholipid Profile of rat Testis, its Unique High level of Monolysocardiolipin and its Lipolytic Capabilities *in Vitro*. A Chromatographic and Densitometric Analysis. Abstract for poster presentation, The Experimental Biology Conference at Washington D.C. on May 3, 2007

**Nosavanh, L. M.**, Tysor, M. Burmeister, M. and Hortsch, M. Developing Molecular Tools and a Simple Genetic System for Investigating Mutations in Caytaxin-type Genes. Abstract for poster presentation, Annual Biomedical Research Conference for Minority Students at Anaheim, CA on November 10, 2006

Helmy, F., Rothenbacher, F., **Nosavanh, L.**, Lowery, J. and Juracka A. A comparative study on the phospholipids profile of Bullfrog cardiac muscle and thigh skeletal muscle and their *in vitro* differential deacylation by endogenous phospholipases. Abstract for poster presentation, The Experimental Biology Conference at San Francisco, California on April 5, 2006

**Nosavanh, L.**, Absher, D., and Myers, R. Targeted Knock-out of the Polyglutamine Tract in Huntingtin. Abstract for poster presentation, Annual Biomedical Research Conference for Minority Students at Atlanta, GA November 3, 2005

**Nosavanh, L.**, Frost, J. A., Tran, N., Wu, X. and Kilpatrick, S. Interaction of Ksr with Extracellular Signal Relating Kinase Pathway Components. Abstract for poster presentation, Annual Biomedical Research Conference for Minority Students at Dallas, TX November 11, 2004

Her publications include:

F. Helmy, **L. Nosavanh**, H. Haynes and A. Juracka. 2008. Phospholipid profile of rat testis, its unique high level of monolysocardiolipin and its lipolytic capabilities in vitro. A chromatographic analysis. Cell Biochem. Funct. Vol. 26 (4). 434-42.

F. Helmy, F. Rothenbacher, **L. Nosavanh**, J. Lowery and A. Juracka. 2007. A Comparative Study of the Phospholipid Profiles of Guinea Pig Cardiac Muscle and Bullfrog Cardiac and Thigh Skeletal Muscle, and their In-Vitro Differential Deacylation by Endogenous Phospholipases. Thin Layer Chromatographic and Densitometric Analysis. J. Planar Chromatogr. Vol. 20. 209 -215.

**ELSEVIER LICENSE  
TERMS AND CONDITIONS**

Dec 14, 2011

---

This is a License Agreement between LaGina M Nosavanh ("You") and Elsevier ("Elsevier") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by Elsevier, and the payment terms and conditions.

**All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.**

Supplier	Elsevier Limited The Boulevard, Langford Lane Kidlington, Oxford, OX5 1GB, UK
Registered Company Number	1982084
Customer name	LaGina M Nosavanh
Customer address	2301 Fountain View Dr. #73 Houston, TX 77057
License number	2807781454181
License date	Dec 14, 2011
Licensed content publisher	Elsevier
Licensed content publication	Mechanisms of Development
Licensed content title	Kidney morphogenesis: cellular and molecular regulation
Licensed content author	Satu Kuure, Reetta Vuolteenaho, Seppo Vainio
Licensed content date	15 March 2000
Licensed content volume number	92
Licensed content issue number	1
Number of pages	15
Start Page	31
End Page	45
Type of Use	reuse in a thesis/dissertation
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
Format	both print and electronic
Are you the author of this Elsevier article?	No
Will you be translating?	No

## Order reference number

Title of your thesis/dissertation      DYSREGULATION OF MEOX2 FOLLOWING WT1 MUTATION IN KIDNEY DEVELOPMENT AND WILMS TUMORIGENESIS

Expected completion date      Dec 2011

Estimated size (number of pages)      117

Elsevier VAT number      GB 494 6272 12

Permissions price      0.00 USD

VAT/Local Sales Tax      0.0 USD / 0.0 GBP

Total      0.00 USD

## Terms and Conditions

## INTRODUCTION

1. The publisher for this copyrighted material is Elsevier. By clicking "accept" in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the Billing and Payment terms and conditions established by Copyright Clearance Center, Inc. ("CCC"), at the time that you opened your Rightslink account and that are available at any time at <http://myaccount.copyright.com>).

## GENERAL TERMS

2. Elsevier hereby grants you permission to reproduce the aforementioned material subject to the terms and conditions indicated.

3. Acknowledgement: If any part of the material to be used (for example, figures) has appeared in our publication with credit or acknowledgement to another source, permission must also be sought from that source. If such permission is not obtained then that material may not be included in your publication/copies. Suitable acknowledgement to the source must be made, either as a footnote or in a reference list at the end of your publication, as follows:

“Reprinted from Publication title, Vol /edition number, Author(s), Title of article / title of chapter, Pages No., Copyright (Year), with permission from Elsevier [OR APPLICABLE SOCIETY COPYRIGHT OWNER].” Also Lancet special credit - “Reprinted from The Lancet, Vol. number, Author(s), Title of article, Pages No., Copyright (Year), with permission from Elsevier.”

4. Reproduction of this material is confined to the purpose and/or media for which permission is hereby given.

5. Altering/Modifying Material: Not Permitted. However figures and illustrations may be altered/adapted minimally to serve your work. Any other abbreviations, additions, deletions and/or any other alterations shall be made only with prior written authorization of Elsevier Ltd. (Please contact Elsevier at [permissions@elsevier.com](mailto:permissions@elsevier.com))

6. If the permission fee for the requested use of our material is waived in this instance,

please be advised that your future requests for Elsevier materials may attract a fee.

7. **Reservation of Rights:** Publisher reserves all rights not specifically granted in the combination of (i) the license details provided by you and accepted in the course of this licensing transaction, (ii) these terms and conditions and (iii) CCC's Billing and Payment terms and conditions.

8. **License Contingent Upon Payment:** While you may exercise the rights licensed immediately upon issuance of the license at the end of the licensing process for the transaction, provided that you have disclosed complete and accurate details of your proposed use, no license is finally effective unless and until full payment is received from you (either by publisher or by CCC) as provided in CCC's Billing and Payment terms and conditions. If full payment is not received on a timely basis, then any license preliminarily granted shall be deemed automatically revoked and shall be void as if never granted. Further, in the event that you breach any of these terms and conditions or any of CCC's Billing and Payment terms and conditions, the license is automatically revoked and shall be void as if never granted. Use of materials as described in a revoked license, as well as any use of the materials beyond the scope of an unrevoked license, may constitute copyright infringement and publisher reserves the right to take any and all action to protect its copyright in the materials.

9. **Warranties:** Publisher makes no representations or warranties with respect to the licensed material.

10. **Indemnity:** You hereby indemnify and agree to hold harmless publisher and CCC, and their respective officers, directors, employees and agents, from and against any and all claims arising out of your use of the licensed material other than as specifically authorized pursuant to this license.

11. **No Transfer of License:** This license is personal to you and may not be sublicensed, assigned, or transferred by you to any other person without publisher's written permission.

12. **No Amendment Except in Writing:** This license may not be amended except in a writing signed by both parties (or, in the case of publisher, by CCC on publisher's behalf).

13. **Objection to Contrary Terms:** Publisher hereby objects to any terms contained in any purchase order, acknowledgment, check endorsement or other writing prepared by you, which terms are inconsistent with these terms and conditions or CCC's Billing and Payment terms and conditions. These terms and conditions, together with CCC's Billing and Payment terms and conditions (which are incorporated herein), comprise the entire agreement between you and publisher (and CCC) concerning this licensing transaction. In the event of any conflict between your obligations established by these terms and conditions and those established by CCC's Billing and Payment terms and conditions, these terms and conditions shall control.

14. **Revocation:** Elsevier or Copyright Clearance Center may deny the permissions described in this License at their sole discretion, for any reason or no reason, with a full refund payable to you. Notice of such denial will be made using the contact information provided by you. Failure to receive such notice will not alter or invalidate the denial. In no event will Elsevier or Copyright Clearance Center be responsible or liable for any costs, expenses or damage



incurred by you as a result of a denial of your permission request, other than a refund of the amount(s) paid by you to Elsevier and/or Copyright Clearance Center for denied permissions.

### LIMITED LICENSE

The following terms and conditions apply only to specific license types:

15. **Translation:** This permission is granted for non-exclusive world **English** rights only unless your license was granted for translation rights. If you licensed translation rights you may only translate this content into the languages you requested. A professional translator must perform all translations and reproduce the content word for word preserving the integrity of the article. If this license is to re-use 1 or 2 figures then permission is granted for non-exclusive world rights in all languages.

16. **Website:** The following terms and conditions apply to electronic reserve and author websites:

**Electronic reserve:** If licensed material is to be posted to website, the web site is to be password-protected and made available only to bona fide students registered on a relevant course if:

This license was made in connection with a course,

This permission is granted for 1 year only. You may obtain a license for future website posting,

All content posted to the web site must maintain the copyright information line on the bottom of each image,

A hyper-text must be included to the Homepage of the journal from which you are licensing at <http://www.sciencedirect.com/science/journal/xxxxx> or the Elsevier homepage for books at <http://www.elsevier.com> , and

Central Storage: This license does not include permission for a scanned version of the material to be stored in a central repository such as that provided by Heron/XanEdu.

17. **Author website** for journals with the following additional clauses:

All content posted to the web site must maintain the copyright information line on the bottom of each image, and

he permission granted is limited to the personal version of your paper. You are not allowed to download and post the published electronic version of your article (whether PDF or HTML, proof or final version), nor may you scan the printed edition to create an electronic version,

A hyper-text must be included to the Homepage of the journal from which you are licensing at <http://www.sciencedirect.com/science/journal/xxxxx> , As part of our normal production process, you will receive an e-mail notice when your article appears on Elsevier's online service ScienceDirect ([www.sciencedirect.com](http://www.sciencedirect.com)). That e-mail will include the article's Digital Object Identifier (DOI). This number provides the electronic link to the published article and should be included in the posting of your personal version. We ask that you wait until you receive this e-mail and have the DOI to do any posting.

Central Storage: This license does not include permission for a scanned version of the material to be stored in a central repository such as that provided by Heron/XanEdu.

18. **Author website** for books with the following additional clauses:

Authors are permitted to place a brief summary of their work online only.  
A hyper-text must be included to the Elsevier homepage at <http://www.elsevier.com>

All content posted to the web site must maintain the copyright information line on the bottom of each image

You are not allowed to download and post the published electronic version of your chapter, nor may you scan the printed edition to create an electronic version.

Central Storage: This license does not include permission for a scanned version of the material to be stored in a central repository such as that provided by Heron/XanEdu.

19. **Website** (regular and for author): A hyper-text must be included to the Homepage of the journal from which you are licensing at <http://www.sciencedirect.com/science/journal/xxxxx>. or for books to the Elsevier homepage at <http://www.elsevier.com>

20. **Thesis/Dissertation**: If your license is for use in a thesis/dissertation your thesis may be submitted to your institution in either print or electronic form. Should your thesis be published commercially, please reapply for permission. These requirements include permission for the Library and Archives of Canada to supply single copies, on demand, of the complete thesis and include permission for UMI to supply single copies, on demand, of the complete thesis. Should your thesis be published commercially, please reapply for permission.

21. **Other Conditions**:

v1.6

If you would like to pay for this license now, please remit this license along with your payment made payable to "COPYRIGHT CLEARANCE CENTER" otherwise you will be invoiced within 48 hours of the license date. Payment should be in the form of a check or money order referencing your account number and this invoice number RLNK500683992.

Once you receive your invoice for this order, you may pay your invoice by credit card. Please follow instructions provided at that time.

Make Payment To:  
Copyright Clearance Center  
Dept 001  
P.O. Box 843006  
Boston, MA 02284-3006

For suggestions or comments regarding this order, contact RightsLink Customer Support: [customer care@copyright.com](mailto:customer care@copyright.com) or +1-877-622-5543 (toll free in the US) or +1-978-646-2777.

Gratis licenses (referencing \$0 in the Total field) are free. Please retain this printable license for your reference. No payment is required.

---

---

## NATURE PUBLISHING GROUP LICENSE TERMS AND CONDITIONS

Dec 14, 2011

This is a License Agreement between LaGina M Nosavanh ("You") and Nature Publishing Group ("Nature Publishing Group") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the payment terms and conditions.

**All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.**

License Number	2807790327843
License date	Dec 14, 2011
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Cancer
Licensed content title	Wilms' tumour: connecting tumorigenesis and organ development in the kidney
Licensed content author	Miguel N. Rivera and Daniel A. Haber
Licensed content date	Sep 1, 2005
Volume number	5
Issue number	9
Type of Use	reuse in a thesis/dissertation
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
High-res required	no
Figures	Figure 3   Alternative splicing of WT1.
Author of this NPG article	no
Your reference number	
Title of your thesis / dissertation	DYSREGULATION OF MEOX2 FOLLOWING WT1 MUTATION IN KIDNEY DEVELOPMENT AND WILMS TUMORIGENESIS
Expected completion date	Dec 2011
Estimated size (number of pages)	117
Total	0.00 USD
Terms and Conditions	

### Terms and Conditions for Permissions

Nature Publishing Group hereby grants you a non-exclusive license to reproduce this

material for this purpose, and for no other use, subject to the conditions below:

1. NPG warrants that it has, to the best of its knowledge, the rights to license reuse of this material. However, you should ensure that the material you are requesting is original to Nature Publishing Group and does not carry the copyright of another entity (as credited in the published version). If the credit line on any part of the material you have requested indicates that it was reprinted or adapted by NPG with permission from another source, then you should also seek permission from that source to reuse the material.
2. Permission granted free of charge for material in print is also usually granted for any electronic version of that work, provided that the material is incidental to the work as a whole and that the electronic version is essentially equivalent to, or substitutes for, the print version. Where print permission has been granted for a fee, separate permission must be obtained for any additional, electronic re-use (unless, as in the case of a full paper, this has already been accounted for during your initial request in the calculation of a print run). NB: In all cases, web-based use of full-text articles must be authorized separately through the 'Use on a Web Site' option when requesting permission.
3. Permission granted for a first edition does not apply to second and subsequent editions and for editions in other languages (except for signatories to the STM Permissions Guidelines, or where the first edition permission was granted for free).
4. Nature Publishing Group's permission must be acknowledged next to the figure, table or abstract in print. In electronic form, this acknowledgement must be visible at the same time as the figure/table/abstract, and must be hyperlinked to the journal's homepage.
5. The credit line should read:  
 Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME]  
 (reference citation), copyright (year of publication)  
 For AOP papers, the credit line should read:  
 Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME],  
 advance online publication, day month year (doi: 10.1038/sj.[JOURNAL  
 ACRONYM].XXXXX)

**Note: For republication from the *British Journal of Cancer*, the following credit lines apply.**

Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication) For AOP papers, the credit line should read:  
 Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME], advance online publication, day month year (doi: 10.1038/sj.[JOURNAL ACRONYM].XXXXX)

6. Adaptations of single figures do not require NPG approval. However, the adaptation should be credited as follows:

Adapted by permission from Macmillan Publishers Ltd: [JOURNAL NAME]  
(reference citation), copyright (year of publication)

**Note: For adaptation from the *British Journal of Cancer*, the following credit line applies.**

Adapted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

7. Translations of 401 words up to a whole article require NPG approval. Please visit <http://www.macmillanmedicalcommunications.com> for more information. Translations of up to a 400 words do not require NPG approval. The translation should be credited as follows:

Translated by permission from Macmillan Publishers Ltd: [JOURNAL NAME]  
(reference citation), copyright (year of publication).

**Note: For translation from the *British Journal of Cancer*, the following credit line applies.**

Translated by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

We are certain that all parties will benefit from this agreement and wish you the best in the use of this material. Thank you.

Special Terms:

v1.1

If you would like to pay for this license now, please remit this license along with your payment made payable to "COPYRIGHT CLEARANCE CENTER" otherwise you will be invoiced within 48 hours of the license date. Payment should be in the form of a check or money order referencing your account number and this invoice number RLNK500683998.

Once you receive your invoice for this order, you may pay your invoice by credit card. Please follow instructions provided at that time.

Make Payment To:  
Copyright Clearance Center  
Dept 001  
P.O. Box 843006  
Boston, MA 02284-3006

For suggestions or comments regarding this order, contact RightsLink Customer Support: [customercare@copyright.com](mailto:customercare@copyright.com) or +1-877-622-5543 (toll free in the US) or +1-978-646-2777.

Gratis licenses (referencing \$0 in the Total field) are free. Please retain this printable license for your reference. No payment is required.

---

---

**NATURE PUBLISHING GROUP LICENSE  
TERMS AND CONDITIONS**

Dec 14, 2011

---

This is a License Agreement between LaGina M Nosavanh ("You") and Nature Publishing Group ("Nature Publishing Group") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the payment terms and conditions.

**All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.**

License Number	2807790564431
License date	Dec 14, 2011
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Cancer
Licensed content title	Wilms' tumours: about tumour suppressor genes, an oncogene and a chameleon gene
Licensed content author	Vicki Huff
Licensed content date	Feb 1, 2011
Volume number	11
Issue number	2
Type of Use	reuse in a thesis/dissertation
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
High-res required	no
Figures	Figure 1   WT1 mutations observed in Wilms' tumour and acute myeloid leukaemia (AML).
Author of this NPG article	no
Your reference number	
Title of your thesis / dissertation	DYSREGULATION OF MEOX2 FOLLOWING WT1 MUTATION IN KIDNEY DEVELOPMENT AND WILMS TUMORIGENESIS
Expected completion date	Dec 2011
Estimated size (number of pages)	117
Total	0.00 USD
Terms and Conditions	

Terms and Conditions for Permissions

Nature Publishing Group hereby grants you a non-exclusive license to reproduce this material for this purpose, and for no other use, subject to the conditions below:

1. NPG warrants that it has, to the best of its knowledge, the rights to license reuse of this material. However, you should ensure that the material you are requesting is original to Nature Publishing Group and does not carry the copyright of another entity (as credited in the published version). If the credit line on any part of the material you have requested indicates that it was reprinted or adapted by NPG with permission from another source, then you should also seek permission from that source to reuse the material.
2. Permission granted free of charge for material in print is also usually granted for any electronic version of that work, provided that the material is incidental to the work as a whole and that the electronic version is essentially equivalent to, or substitutes for, the print version. Where print permission has been granted for a fee, separate permission must be obtained for any additional, electronic re-use (unless, as in the case of a full paper, this has already been accounted for during your initial request in the calculation of a print run). NB: In all cases, web-based use of full-text articles must be authorized separately through the 'Use on a Web Site' option when requesting permission.
3. Permission granted for a first edition does not apply to second and subsequent editions and for editions in other languages (except for signatories to the STM Permissions Guidelines, or where the first edition permission was granted for free).
4. Nature Publishing Group's permission must be acknowledged next to the figure, table or abstract in print. In electronic form, this acknowledgement must be visible at the same time as the figure/table/abstract, and must be hyperlinked to the journal's homepage.
5. The credit line should read:  
Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME]  
(reference citation), copyright (year of publication)  
For AOP papers, the credit line should read:  
Reprinted by permission from Macmillan Publishers Ltd: [JOURNAL NAME],  
advance online publication, day month year (doi: 10.1038/sj.[JOURNAL  
ACRONYM].XXXXX)

**Note: For republication from the *British Journal of Cancer*, the following credit lines apply.**

Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication) For AOP papers, the credit line should read:  
Reprinted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME], advance online publication, day month year (doi: 10.1038/sj.[JOURNAL ACRONYM].XXXXX)

6. Adaptations of single figures do not require NPG approval. However, the adaptation should be credited as follows:

Adapted by permission from Macmillan Publishers Ltd: [JOURNAL NAME]  
(reference citation), copyright (year of publication)

**Note: For adaptation from the *British Journal of Cancer*, the following credit line applies.**

Adapted by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

7. Translations of 401 words up to a whole article require NPG approval. Please visit <http://www.macmillanmedicalcommunications.com> for more information. Translations of up to a 400 words do not require NPG approval. The translation should be credited as follows:

Translated by permission from Macmillan Publishers Ltd: [JOURNAL NAME]  
(reference citation), copyright (year of publication).

**Note: For translation from the *British Journal of Cancer*, the following credit line applies.**

Translated by permission from Macmillan Publishers Ltd on behalf of Cancer Research UK: [JOURNAL NAME] (reference citation), copyright (year of publication)

We are certain that all parties will benefit from this agreement and wish you the best in the use of this material. Thank you.

Special Terms:

v1.1

If you would like to pay for this license now, please remit this license along with your payment made payable to "COPYRIGHT CLEARANCE CENTER" otherwise you will be invoiced within 48 hours of the license date. Payment should be in the form of a check or money order referencing your account number and this invoice number RLNK500684004.

Once you receive your invoice for this order, you may pay your invoice by credit card. Please follow instructions provided at that time.

Make Payment To:  
Copyright Clearance Center  
Dept 001  
P.O. Box 843006  
Boston, MA 02284-3006

For suggestions or comments regarding this order, contact RightsLink Customer Support: [customer care@copyright.com](mailto:customer care@copyright.com) or +1-877-622-5543 (toll free in the US) or +1-978-646-2777.

Gratis licenses (referencing \$0 in the Total field) are free. Please retain this printable license for your reference. No payment is required.

---

---