

5-2012

FZD6, MATN2 AND SLC25A32, POSSIBLE CANDIDATE GENES IN NONSYNDROMIC CLEFT LIP AND PALATE

Nevena Cvjetkovic

Follow this and additional works at: https://digitalcommons.library.tmc.edu/utgsbs_dissertations



Part of the [Genetics and Genomics Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Cvjetkovic, Nevena, "FZD6, MATN2 AND SLC25A32, POSSIBLE CANDIDATE GENES IN NONSYNDROMIC CLEFT LIP AND PALATE" (2012). *The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences Dissertations and Theses (Open Access)*. 281.
https://digitalcommons.library.tmc.edu/utgsbs_dissertations/281

This Thesis (MS) is brought to you for free and open access by the The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences at DigitalCommons@TMC. It has been accepted for inclusion in The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences Dissertations and Theses (Open Access) by an authorized administrator of DigitalCommons@TMC. For more information, please contact digitalcommons@library.tmc.edu.

*FZD6, MATN2 AND SLC25A32, POSSIBLE CANDIDATE GENES IN
NONSyndromic Cleft Lip and Palate*

by

Nevena Cyjetkovic

APPROVED:

Supervisory Professor
Jacqueline T. Hecht, PhD

Eric C. Swindell, PhD

John F. Teichgraeber, MD

Ariadne M. Letra, DDS, MS, PhD

Marianna Raia, MS, CGC

APPROVED:

Dean, The University of Texas
Graduate School of Biomedical Sciences at Houston

*FZD6, MATN2 AND SLC25A32, POSSIBLE CANDIDATE GENES IN
NONSyndromic Cleft Lip and Palate*

A
THESIS

Presented to the Faculty of
The University of Texas
Health Science Center at Houston
and
The University of Texas
M. D. Anderson Cancer Center
Graduate School of Biomedical Sciences
in Partial Fulfillment
of the Requirements
for the Degree of
MASTER OF SCIENCE

by

Nevena Cvjetkovic, BS
Houston, Texas

May, 2012

ACKNOWLEDGEMENTS

First and foremost I would like to thank Dr. Hecht, my thesis advisor, for giving me the opportunity to work on a project that I have a personal connection with. Her guidance through the process allowed me to accomplish something I am extremely proud of. I would also like to thank my committee: Dr. Swindell, Dr. Teichgraeber, Dr. Letra, and Marianna Raia for all their support and advice. Additionally, I would like to thank my lab-mates: Katelyn, for helping me from day 1, showing me the ropes, and always answering my questions, no matter how trivial they were, Lori, for her devotion to this project and all the hard work she put into it, Robert for making the time in lab very entertaining, as well as the rest of Dr. Hecht's lab for always being available and willing to help.

I would like to thank the UT Genetic Counseling program, Claire Singletary, Sarah Jane Noblin, my mentor Jennifer Czerwinski, and the numerous supervisors for their dedication to my training and education over the past two years. I would also like to thank Norma Leal and Patricia McDonald for their help behind the scenes. I would especially like to thank my wonderful classmates: Ann, Carolyn, Katie, Roya, Salma, and Sam. I could not imagine a better group of girls to experience graduate school with. When the going got tough having them right there beside me made everything more bearable. Additionally I'd like to thank my second year Emily who proved to be an awesome mentor and a true friend.

I would like to thank my boyfriend Tihomir for sticking by me through this trying time for any relationship. I would also like to thank all my family who always believed in me and encouraged me in whatever I decided to do. And most importantly, I'd like to thank my father Nenad, mother Ljubenka, and sister Jovana for their unconditional love and support. It is very simple fact of life that without them I would not be who or where I am.

*FZD6, MATN2 AND SLC25A32, POSSIBLE CANDIDATE GENES IN
NONSyndROMIC CLEFT LIP AND PALATE*

Publication No. _____

Nevena Cvjetkovic, B.S.

Nonsyndromic cleft lip with or without cleft palate (NSCLP) is a common birth defect with a multifactorial etiology. Despite decades of research, the genetic underpinnings of NSCLP still remain largely unexplained. A genome wide association study (GWAS) of a large NSCLP African American family with seven affected individuals across three generations found evidence for linkage at 8q21.3-24.12 (LOD = 2.98). This region contained three biologically relevant candidate genes: *Frizzled-6 (FZD6)* (LOD = 2.8), *Matrilin-2 (MATN2)* (LOD = 2.3), and *Solute Carrier Family 25, Member 32 (SLC26A32)* (LOD = 1.6). Sequencing of the coding regions and the 5' and 3' UTRs of these genes in two affected family members identified a rare intronic variant, rs138557689 (c.-153+432A>C), in *FZD6*. The rs138557689/C allele segregated with the NSCLP phenotype; *in silico* analysis predicted and EMSA analysis showed that the 138557689/C allele creates new DNA binding sites. *FZD6* is part of the WNT pathway, which is involved in craniofacial development, including midface development and upper lip fusion. Our novel findings suggest that an alteration in *FZD6* gene regulation may perturb this tightly controlled biological pathway and in turn contribute to the development of NSCLP in this family. Studies are underway to further define how the rs138557689/C variant affects expression of *FZD6*.

TABLE OF CONTENTS

Acknowledgements	iii
Abstract	iv
List of Figures	vi
List of Tables	vii
Background	1
Materials and Methods	18
Results	21
Discussion	25
Appendix	30
References	33
Vita	51

LIST OF FIGURES

Figure 1. Development of lip and palate	6
Figure 2. Pedigree of the African American NSCLP family	10
Figure 3. Canonical WNT/ β -catenin signaling pathway	13
Figure 4. Segregation of rs138557689 in the African American NSCLP family	21
Figure 5. Conservation analysis of rs138557689	23
Figure 6. EMSA of rs138557689	24
Figure 7. Schematic representation of WNT Signaling Pathway	27

LIST OF TABLES

Table 1. Syndromes with CLP or CPO	3
Table 2. Frequency of rs138557689 C allele	22
Supplementary Table 1. Primers for <i>FZD6</i>	30
Supplementary Table 2. Primers for <i>MATN2</i>	31
Supplementary Table 3. Primers for <i>SLC25A32</i>	32

INTRODUCTION

Isolated or nonsyndromic cleft lip with or without cleft palate (NSCLP) is the fourth most common birth defect affecting more than 4,000 births each year in the United States [1,2,3]. Despite the improvements in treatment, NSCLP has considerable medical and financial implications for the affected individuals and their families, and the effects on speech, hearing, and appearance can lead to adverse psychosocial and psychiatric outcomes [4,5,6]. Individuals with NSCLP require multidisciplinary team care through adulthood and continue to demonstrate increased morbidity and mortality rates compared to the general population [7,8].

Cleft lip and palate has been recognized and documented throughout history. First evidence of clefting was noted in an Egyptian mummy dating from 2400 to 1300 BC [9]. A 2000-year-old statue of a Columbian king, a 2000-year old African mask, a Ming dynasty painting, and a 20th century Russian painting all depicted orofacial clefts [9]. What appears to be the first documented treatment of cleft lip and palate was performed in 390AD by a Chinese physician [10].

Many different explanations for causation of cleft lip and palate have evolved among the various cultures and populations in the world. Causes are often attributed to maternal impressions or supernatural events [11]. The term “hare-lip” to describe cleft lip originated from beliefs that clefting results from eating or looking at a rabbit [12,13]. A prevailing belief in many cultures is that a child will be born with a birth defect, including cleft lip and palate, if a woman pities or makes fun of an affected individual while pregnant [13,14,15]. Supernatural causes, such as the effect of the moon (lunar eclipse), are alleged causes of cleft lip and palate in Mexican folklore and among Hispanic populations [14,15,16], while populations in India and Nigeria ascribed the cleft to “God's will”, evil and ancestral spirits,

or sins committed in past lives [17,18]. Thesis work of Fogh-Andersen in 1942 was the first to widely recognize the hereditary component of CLP [19].

Despite the fact that NSCLP has been part of human life for many centuries, and the fact that many beliefs for the causation of this birth defect exist, the etiology of NSCLP is still largely unknown. Continual research in this area is needed to help elucidate the underlying etiologies of this common birth defect in order to improve genetic counseling for recurrence, diagnosis, prevention, and treatment, as well as to improve our understanding of the development of clefts.

Classification of Clefts

Orofacial clefts are divided into two groups based on developmental origin: anterior and posterior clefts [20]. Anterior cleft anomalies include cleft lip (CL) only, or cleft lip and primary palate (CLP) [20]. Anterior clefts can extend through the lip and alveolar part of the maxilla (primary or hard palate) to the incisive fossa [20]. Posterior cleft anomalies, which include clefts of the secondary or soft palate, present as cleft palate only (CPO) [9,20]. Posterior clefts extend through the soft and hard sections of the palate to the incisive fossa [20].

Clefts can further be divided into syndromic or nonsyndromic. Syndromic clefts are distinguished by the presence of other congenital malformations and may be associated with specific genes, chromosomal abnormalities, or maternal teratogenic exposures [21].

Approximately 30% of CLP and 50% of CPO are associated with other syndromes or anomalies, and there are currently over 400 different conditions listed on Online Mendelian Inheritance of Man in which clefts occur (<http://www.ncbi.nlm.nih.gov/omim>) [21]. Table 1

is an abbreviated list of syndromes which have CLP or CPO as a phenotypic finding and the genes associated with them. The remaining 70% of cases of CLP and 50% of cases of CPO are isolated, and therefore are referred to as nonsyndromic [21]. Nonsyndromic CLP (NSCLP) is the focus of this study.

Table 1.

Syndromes with CLP		Syndromes with CPO	
Syndrome	Gene	Syndrome	Gene
Autosomal dominant developmental malformations, deafness, and dystonia	<i>ACTB</i>	Oculofaciocardiodental	BCOR
Craniofrontonasal	<i>EFNB1</i>	CHARGE	CHD7
Roberts	<i>ESCO2</i>	Escobar multiple pterygium	CHRNA
Holoprosencephaly	<i>GLI2, SHH, SLX3, TGIF</i>	Stickler	COL2A1, COL11A1, COL11A2
Oro-facial-digital	<i>GLI3</i>	Smith-Lemli-Opitz	DHCR7
Van der Woude	<i>IRF6</i>	Crouzon/Apert	FGFR2
X-linked mental retardation and CL/P	<i>PHF8</i>	Kabuki	MLL2
Gorlin	<i>PTCH1</i>	Cornelia de Lange	NIPBL
CLP – ectodermal dysplasia	<i>PVRL1</i>	Diastrophic dysplasia	SLC26A2
Branchio-oculo-facial	<i>TFAP2A</i>	Campomelic dysplasia	SOX9
Ectrodactyly-ectodermal dysplasia-clefting-	<i>TP63</i>	Treacher Collins	TCOF1
Tetra-amelia with CLP	<i>WNT3</i>	Loeys-Dietz	TGFBR1, TGFBR2

Adapted from M.J. Dixon, M.L. Marazita, T.H. Beaty, J.C. Murray, 2011

A cleft can be unilateral, occurring on one side of the face, or bilateral, occurring on both sides of the face. Unilateral clefts account for 90% of CLP (~60% of which occur on the left side) while bilateral clefts account for the remaining 10% [21].

Recurrence of syndromic CLP depends on the underlying genetic condition, while recurrence risk for NSCLP depends on several factors including number of affected individuals in family, relation of affected family member, and laterality of the cleft [22].

The empirical recurrence risk for siblings is 2-3% in a case of unilateral CLP and 3-5% in the case of bilateral CLP [22].

Birth Prevalence

Cleft lip with or without palate is estimated to occur in approximately 1/700 to 1/1000 births but the prevalence of clefts varies by ethnicity [21,23]. The frequency of NSCLP appears to be highest among Native Americans (~3.6/1000 births) followed by Japanese (~2.1/1000 births) and Chinese (~1.7/1000 births) populations, and appears to be the lowest among African derived populations (~0.3/1000 births) [9]. In Texas, the prevalence of NSCLP is 0.67/1000 for Non-Hispanic Whites, 0.63/1000 for US-born Hispanics, 0.65/1000 for non-US born Hispanics, and 0.40/1000 for African Americans [1].

Birth prevalence of NSCLP also varies by gender with a 2:1 male to female ratio [24]. An association between NSCLP and socioeconomic status is implicated by studies which show groups from rural, lower socioeconomic areas having a higher birth prevalence of NSCLP compared to ethnically similar groups within higher socioeconomic status [21].

The birth prevalence of isolated CPO is approximately 1/2000 births and does not appear to differ much between various ethnicities [21]. In contrast to NSCLP, there is a female predominance of isolated CPO with a 3:2 female to male ratio [25].

Normal Development of the lip and palate

Understanding of normal development of the upper lip, primary, and secondary palates is important to help explain the reason for the altered development seen in CLP. Normal craniofacial development is a complex process directed by intricate pathways


important for cell induction, differentiation, proliferation, migration, patterning, and apoptosis [8,26]. Molecular pathways, including the Bmp, Fgf, Shh, and Wnt pathways, coordinate craniofacial development through both synergistic and antagonistic signaling [26]. Numerous genes, which code for growth factors, transcription factors, cell adhesion molecules, and signaling molecules, strictly regulate the development of the face [19,27,28,29,30]. Theoretically, perturbations within any of these genes and pathways could alter craniofacial morphogenesis and result in clefting.

The human face begins developing in the 4th week of embryogenesis [28]. Neural crest cells (NCCs), which are the major source of connective tissue components, migrate from the neural folds and combine with the core mesoderm and the epithelia to establish five facial primordia [20,26]. As depicted in Fig. 1a, the primordia consist of a single frontonasal prominence, two mandibular and two maxillary prominences, all of which surround the primitive oral cavity [8,28]. Ectodermal thickenings form nasal placodes and results in the formation of nasal pits by the end of week 4. The nasal pits further divide the frontonasal prominence into two horseshoe-shaped medial and lateral nasal processes (Fig. 1b) [8,26].

Rapid growth of the underlying mesenchyme of the facial primordia causes the medial nasal prominences to merge with each other and with the maxillary and lateral nasal prominences between the 7th to 10th week of embryogenesis [20]. The contact epithelia between the prominences are broken up by cell apoptosis to give rise to the intermaxillary segment from which the philtrum of the upper lip, the premaxillary part of the maxilla and the primary palate are formed (Fig. 1c) [20,26].

The secondary palate, which comprises majority of the soft palate, except for the anterior portion that holds the incisor teeth, has a different developmental origin [20].

Palatal shelves, which originate from the maxillary processes, appear in the 6th week of development and grow downward, vertical to the sides of the developing tongue (Fig. 1d) [8,9]. The palatal shelves then elevate to a horizontal position above the tongue (Fig. 1e), and fuse together (Fig. 1f) [8,20]. This is thought to occur earlier in females than in males [9]. Additionally, the two palatal shelves merge with the primary palate and the nasal septum dividing the oral from the nasal cavity [8]. This process is complete by the end of the 10th week [9,20].



Development of Cleft Lip and Palate

Cleft lip and palate can result when any of the multifaceted factors involved in normal facial morphogenesis are disrupted or altered. These may include failure of fusion of

the processes that form the face, distortions in epithelial movement, deficiency in epithelial and mesenchymal transformation (EMT), or failure in apoptosis [26].

Deficiency of mesenchyme in the maxillary prominence(s) and the median palatal process leads to anterior clefts [20]. Failure of one maxillary prominence to unite with the two merged medial nasal prominences results in unilateral cleft of the upper lip, while failure of both maxillary prominences to unite with the merged medial nasal prominences results in bilateral cleft of the upper lip [9,20].

Defective development of the secondary palate leads to posterior palatal clefts [20]. This is mainly due to the failure of mesenchymal masses in the lateral palatal processes to meet and fuse with each other and the nasal septum [9,20].

Treatment of NSCLP

Treatment of a patient with CLP requires a multidisciplinary approach which includes craniofacial and maxillofacial surgery, dentistry and orthodontic, audiology, otolaryngology, speech and language, nursing, pediatrics, genetics, and social services [31]. Many different surgical protocols exist to repair CLP and treat ensuing complications [21]. Currently, however, there is not an internationally accepted protocol for CLP repair [32].

Surgical repair of the CLP usually occurs within the first six months of life, the general rule being to perform surgical repair when the child is approximately 10 weeks of age, weighs at least 10 lb, and has achieved serum hemoglobin of 10 mg/ml [33]. There are various techniques of CLP repair which have been refined throughout time and some of these include LeMesurier (1955), Tennison (1952), Millard (1950), Skoog (1969), Randall

(1990) and Brauer (1985). The choice of technique, however, most often depends on surgeon preference and severity of cleft [21].

Etiology of NSCLP

NSCLP does not follow a traditional Mendelian pattern of inheritance but is believed to result from a combination of several genetic variants which act in an additive fashion and interact with the environment to produce the phenotype [21]. NSCLP is therefore considered to be a multifactorial disorder that results from the interaction of both genetic and environmental factors [21]. Genetic evidence for NSCLP comes from multiple studies. These studies show that NSCLP aggregates in families, that there is a family history for clefting in 24-33% of NSCLP patients with a 10 to 32-fold increase in recurrence risk to first degree relatives, that the heritability of NSCLP in the NHW population is approximately 76%, and that the rate of concordance is higher in monozygotic (25-40%) than dizygotic (3-6%) twins [9,21,34,35,36,37,38,39]. Various environmental factors have also been associated with an increased risk for NSCLP including smoking, alcohol, diet, infections, fever, drugs, and teratogenic agents during early pregnancy [21,40,41,42,43].

Only a handful of genes associated with NSCLP have been identified to date and these comprise approximately 20% of the genetic causes of NSCLP [44]. Interferon regulator factor 6 (*IRF6*) is one of the genes that has shown an association with NSCLP in multiple populations [45,46,47,48,49,50]. Mutations in *IRF6* are also known to cause van der Woude syndrome [51]. Various other genes, which include growth factors ($TGF\alpha$, $TGF\beta$), transcription factors (*MSX1*, *TBX22*), genes involved in xenobiotics (*CYP1A1*, *GSTM1*, *NAT2*), and other genes (*RARA*, *MTHFR*, *CRISPLD2*), have also been associated

with NSCLP [52,53,54,55,56,57,58,59,60,61]. The constant identification of novel genes for the pathogenesis of NSCLP across various populations highlights the genetic heterogeneity of this complex birth defect.

Gene identification in NSCLP


Multiple genetic approaches have been used to identify the genes and pathways contributing to NSCLP. These approaches include assessment of mouse models, linkage analysis using large multiplex families and affected relative pairs, association studies using family-based or case-control samples, identification of chromosomal anomalies or microdeletions in affected cases, and candidate gene studies [62].

Recently, genome-wide association studies (GWAS) have expanded the research in NSCLP and markedly helped in the identification of new candidate genes. GWAS studies use known single nucleotide polymorphisms (SNPs) that cover the genome to identify regions that are linked to and/or associated with NSCLP [62]. Identification of candidate genes by GWAS permits further analysis of individual genes through direct sequencing [62]. Sequencing of the candidate genes allows for identification of sequence variants within the gene which could potentially contribute to the development of NSCLP [62]. The sequence variants can be located within protein coding regions as well as noncoding regions, which include promoter/enhancer regions, or introns of the gene [63].

8q21.3-24.12 Chromosomal Region

A GWAS study performed in 2009 identified a major locus for NSCLP on chromosome 8q24 but the region with highest linkage did not contain any protein coding

genes [64]. Subsequent GWAS studies confirmed the association with the 8q24 locus and identified additional loci including 10q25, 7q22, 20q12, and 1p22 [62]. A smaller GWAS study reported by Chiquet et al. evaluated 10 multiplex NSCLP families including one large African-American family with 11 affecteds across 3 generations (Fig. 2). It was performed to confirm existing and identify new NSCLP chromosomal regions [65]. The study found evidence for linkage on 8q21.3-24.12 (LOD=2.98) [65]. Further evaluation of this region using linkage analysis generated a maximum multipoint LOD score of 2.8 for SNPs in *Frizzled 6* (*FZD6*), 2.3 in *Matrilin-2* (*MATN2*), and 1.6 in *Solute carrier family 25, member 32* (*SLC26A32*), prompting further analysis of these three genes and their connection to NSCLP.



***Frizzled-6* and WNT pathway**

Human *Frizzled-6* (*FZD6*) gene maps to chromosome 8q22.3-q23.1 and encodes a 706 amino-acid seven-pass transmembrane protein with a cystine-rich domain in the N-terminal extracellular region, two-N linked glycosylation sites, and two cystine residues in the second and third extracellular loops [66]. A total of 7 exons, 6 of which are coding, comprise the *FZD6* gene [66]. *FZD6* amino acid sequence is well conserved between mouse and human genes with an amino acid identity of 83.3% [66].

FZD6 is part of the *Frizzled* gene family which code for a group of receptors critical for initiation of wingless-type (WNT) signaling pathway [67]. The Frizzled receptors share conserved features which include a cystine-rich domain in the N-terminal extracellular region implicated in the binding of WNT ligands, N-linked glycosylation sites, two cystine residues in the second and third extracellular loops, and a Ser/Thr-x-Val motif in the C-terminus which acts as a binding site for the cytoplasmic protein containing the PDZ domain [66,68,69].

FZD6 codes for three mRNA isoforms that are detected in both adult and fetal tissues [70]. *FZD6* was shown to be important for hair patterning in both *Drosophila* and mice as well as mice claw morphogenesis and appears to act within the WNT/PCP (planar cell polarity) pathway [71,72,73]. Mutations in *FZD6* have been associated with isolated autosomal recessive nail dysplasia and open neural tube defects in humans [71,74,75].


WNT signaling regulates differentiation and proliferation of a variety of cell types during development and is an important regulator of various functions within the cells [76]. Signaling by the WNT pathway directs cell proliferation, cell polarity, and cell fate determination during embryogenesis and homeostasis, regulates multiple developmental

processes, and plays a critical role in embryogenesis through both canonical (β -catenin dependent) and noncanonical (WNT/PCP and Ca^{2+} /CAMKII) signaling [76,77].

The canonical/ β -catenin pathway is the best understood. As depicted in Fig. 3a, when a WNT ligand is absent, cytoplasmic β -catenin forms a complex with the scaffolding protein Axin, the tumor suppressor *adenomatous polyposis coli* gene product (APC), glycogen synthase kinase 3B (GSK3B) and casein kinase 1 (CK1), and is phosphorylated by CK1 and subsequently by GSK3B [67]. Phosphorylated β -catenin is then recognized by the E3 ubiquitin ligase (UB), and is targeted for degradation within the proteasome [78]. This degradation, in turn, prevents β -catenin from reaching the nucleus, where the WNT target genes are repressed by the DNA-bound T cell factor/lymphoid enhancer factor (TCF-TLE)/Groucho and histone deacetylases (HDAC) [67].

Transcription of WNT target genes is activated in the presence of WNT ligand. As shown in Fig. 3b, the WNT ligand binds to a receptor complex formed between Frizzled and low-density lipoprotein receptor related protein 5 or 6 (LRP5/6). This complex then recruits a scaffolding protein Dishevelled (Dvl) and leads to LRP5/6 phosphorylation and Axin recruitment [67,77]. This interrupts Axin-mediated phosphorylation and subsequent degradation of β -catenin, and allows β -catenin to build up in the nucleus where it can activate TCF/LEF and initiate transcription of WNT responsive genes [67].

Figure 3. Canonical WNT/ β -catenin signaling pathway



In contrast to other Frizzled receptors, FZD6 protein lacks the Ser/Thr-X-Val motif at the C-terminal common to other frizzled receptors and has been shown to repress canonical WNT signaling through the noncanonical Ca^{2+} /CaMKII pathway by inhibiting the TCF/LEF binding activity and down-regulating β -catenin targeted transcription of WNT genes [66,70]. Removal of *FZD6*'s N or C terminal sequences, however, abolishes the repressive activity of the protein [70].

Mutations in the WNT pathway have been linked to a variety of birth defects, cancers, and other diseases [67]. Importantly, WNT signaling has been shown to play a role in craniofacial development, including regional specification in the vertebrate face, neural crest induction and differentiation, mid-face development and upper lip fusion in mice, and

facial morphogenesis in mice [26,79,80,81,82,83,84,85]. Craniofacial abnormalities, including orofacial clefts, are found in WNT knockout mice and zebrafish [86,87]. An association with individual WNT genes and NSCLP has been found in humans (*WNT3*, *WNT3A*, *WNT5A*, *WNT7A*, *WNT8A*, *WNT9B* and *WNT11*) [87,88,89,90]. Based on this information, *FZD6* is a strong candidate gene for NSCLP. Coding mutations or variants in the regulatory regions of *FZD6* may alter the tightly regulated WNT pathway, and in turn contribute to NSCLP.

Matrilin-2 and the extracellular matrix assembly

Matrilin-2 (MATN2) gene maps to chromosome 8q22.1-q22.3 and encodes a 956 amino-acid protein whose structure consists of a putative signal peptide, two von Willerbrand factor A (vWFA)-like domains parted by ten epidermal growth factor (EGF)-like domains, a unique segment not seen in other matrilins, and an α -helical coiled-coil (CC) domain [91,92,93]. *MATN2* has 19 exons and is transcribed from two alternative promoters, one of which is an upstream housekeeping-type promoter functional in fibroblasts and other cell types, and a downstream TATA-like promoter restricted to only embryo fibroblast and certain cell lines [94]. Matrilin-2 protein functions as a part of a tightly regulated pathway in the extracellular matrix (ECM) assembly and it is found in a wide variety of connective tissue cells, smooth muscle cells, and both the dermis and the epidermis in humans [93,95]. Human and mice Matrilin-2 proteins show 86.5% identity [93].

Matrilin-2 belongs to a family of four ECM proteins (matrilin-1, -2, -3, and -4), all of which share a structure made of vWFA and EGF domains, and a α -helical CC segment [92].

The matrilin genes also share a phase I, U-12-type AT-AC intron that is located in a strictly conserved position that separates the two exons for the CC domain [94]. While matrilin-1 and matrilin-3 are expressed mainly in cartilage and skeletal tissue, matrilin-2 and -4 have wider tissue distribution, including some non-skeletal tissues [91]. Matrilins -1, -2, and -3 can form filamentous networks that can connect to collagen fibers [93]. While Matrilin-2 is expressed in the mesoderm and epithelium in mice it is expressed in the dermal side of the basement membrane at the dermal-epidermal junction, as well as in keratinocytes and fibroblasts in humans [95].

Mutations in *MATN3* gene were found to be associated with autosomal dominant forms of multiple epiphyseal dysplasia (MED) and mutations in *MATN1* have been associated with osteoarthritis and relapsing polychondritis. *MATN2* and *MATN4*, to date, have not been associated with any disease [92,96]. Knock-out mice lacking *Matn2* have been shown to develop without any obvious abnormalities [97]. Few disorders, including Klippel-Fiel syndrome with laryngeal malformations and Cohen syndrome (which consists of facial, oral, ocular, and limb deformities, and well as intellectual disability), have been linked to the 8q22 position [93]. *MATN2*, however, has not been identified as a candidate gene in these syndromes [93].

Despite the lack of obvious craniofacial phenotype in *Matn2* knock-out mice, interactions of ECM components, growth factors, and embryonic tissues are known to be involved in the normal development of cranial primordia and orofacial structures [98,99] and alterations in the ECM components could potentially lead to NSCLP.

Solute carrier family 25, member 32 and Folate metabolism

Solute carrier family 25, member 32 (SLC25A32) gene, also called *Mitochondrial Folate Transporter (MTF)*, is located on chromosome 8q21.2. *SLC25A32* consists of 7 exons and encodes a 315 amino-acid protein transporter which shuttles folates from the cytoplasm into the mitochondria [100]. The protein consists of six transmembrane domains: three loops facing the mitochondrial matrix and three repeats of an energy transfer sequence [100]. The protein does not contain an ATP-binding motif and probably functions as an ATP-independent transporter [100].

Solute carriers are comprised of 43 identified families which control the movement of various substances across the cell membrane [101]. Members of the solute carrier family 25 (SLC25) are known to transfer a variety of substances across the mitochondrial membrane, and are therefore referred to as mitochondrial solute carriers [102]. The *SLC25A32* protein is shown to transport folate across the mitochondrial membrane [100].

Folate metabolism is a complex process consisting of multiple genes and pathways. Folate is absorbed from the environment, transported across the cell membrane, and compartmentalized between the mitochondria and the cell cytoplasm before it is transported into the mitochondria [103]. The transport of folate into the mitochondria involves various transporters and carriers, including the *SLC25A32* transporter [100,103].

The importance of folate in prevention of birth defects has been investigated since folate fortification reduced the occurrence of spina bifida, an open neural tube defect (ONTD), by approximately 70% [104]. Because ONTDs and NSCLP both develop from migrating neural crest cells, studies have been performed to assess the role of folic acid in NSCLP in hopes to validate the hypothesis that folic acid deficiency may also contribute to

NSCLP [105,106]. Observational studies of folic acid supplementation and the recurrence of NSCLP have not reliably shown the same decrease as with ONTD. A few meta-analysis studies, however, showed decreases of 18-23% in the risk of NSCLP with supplementation of folic acid and/or multivitamins during pregnancy [107,108]. Studies also suggest that variations in the folate pathway genes may contribute to NSCLP [105]. Disturbances in the folate transport, therefore, may also contribute to NSCLP.

Significance of this study

NSCLP is a common birth defect caused by both genetic and environmental factors. Despite decades of research, genetic contributions to NSCLP have yet to be explained, and there remains the need to identify candidate genes. The goal of this project is to further elucidate the genetic etiology of NSCLP by sequencing the coding and 5' and 3' UTR regions of three biologically relevant candidate genes, *FZD6*, *MATN2*, and *SLC25A32*, previously identified in a genome wide scan in a large multiplex African-American family. This family is unique because NSCLP has a low prevalence in the African American population yet there are numerous affected individuals in multiple generations. Sequence variants in any of these three genes may disrupt the highly regulated craniofacial development and give rise to NSCLP. Identification of new genes for NSCLP would not only benefit in diagnosis, prevention, and counseling for this birth defect, but may help in understanding the developmental pathways involved in craniofacial morphogenesis.

MATERIALS AND METHODS

IRB Approval

This study was approved by the Committee for the Protection of Human Subjects at the University of Texas Health Science Center at Houston (HSC-MS-03-090 and HSC-MS-11-0336).

Family

The focus of this study is an African American NSCLP family with 11 affected individuals, 7 of whom were available for evaluation (Fig. 2). All family members were evaluated by one author (JTH). No other anomalies or lip pits were present in any family members. DNA samples from these 7 individuals and 13 connecting or related unaffected family members were subjected to a previously described 6K Illumina IVb genome scan and linkage analysis [65]. A maximum multipoint LOD score of 3.0 can be obtained for this family.

Sequencing

The genomic structure of *FZD6*, *MATN2*, and *SLC25A32* was determined using the NCBI GenBank (www.ncbi.nlm.nih.gov). Forward and reverse primers were designed to capture the sequence of each exon and approximately 50-100 bps upstream and downstream of the intron/exon junction, as well as the complete 5' and 3' untranslated regions (UTRs) for all three isoforms of *FZD6*, two isoforms of *MATN2*, and *SLC25A32* (Supplemental Tables 1-3). DNA samples from two affected family members (Fig 2: III-15 and IV-2) were sequenced for each gene. These individuals were chosen from the opposite sides of the family since they are less likely to share common familial variants. After initial data analysis, 24 additional family members (5 affected and 19 unaffected) were sequenced for

rs138557689 using *FZD6* primer set E1C (Supplementary Table 1). Standard PCR amplification conditions were used and the annealing temperatures for each primer set are shown in the Supplemental Tables 1-3. Amplified PCR product was purified according to manufacturer's protocol (Qiagen, Valencia, CA). Sequencing results were compared to consensus sequences obtained from NCBI public database and analyzed using Sequencer v4.9 (Gene Codes, Ann Arbor, MI).

Variant Analysis

Sequence variants were identified using dbSNP (www.ncbi.nlm.nih.gov/projects/SNP). Only sequence changes shared by both affected individuals (Fig 2: III-15 and IV-2) were considered. SNPs identified in the potential regulatory regions, 5' UTR and the first two introns of the gene, were assessed for their effect on DNA binding using three online *in silico* analyses: Alibaba2, Patch, and Transcription Element Search Software (TESS) [109,110,111]. SNPs identified in the 3'UTR region were assessed for their effect on microRNA binding sites using microRNAMap and miRBase databases [112,113]. SNPs identified in the coding region were analyzed using PolyPhen and SIFT [114,115].

Species Conservation Analysis

SNPs were assessed for evolutionary conservation using the UCSC Genome Browser Multiz Alignments of 46 Vertebrates track (www.genome.ucsc.edu/cgi-bin/hgGateway) and the ECR Browser tool (<http://ecrbrowser.dcode.org>). SNPs were further assessed using the genomic evolutionary rate profiling (GERP) track within the UCSC browser to estimate the evolutionary constraint rates for individual nucleotide positions [116]. GERP scores are positive in constrained regions and negative in neutral DNA [116].

Electrophoretic mobility shift assay (EMSA)

EMSA was used to evaluate whether an ancestral or alternate allele in the SNP of interest altered DNA binding. 20-mer oligonucleotide sequences were hybridized incorporating either the ancestral or alternate allele (Integrated DNA Technologies, Coralville, IA) and labeled with P³² dCTP (40,000cpm). The binding reaction was carried out in 20 uL binding buffer mix (1M Tris (pH 7.5), 1M KCl, 80% glycerol, 10% NP-40, 0.5M EDTA, 100 mM PMSF, 1M DTT and H₂O), containing 1 uL poly dGdC and 1 uL Cos7 cell nuclear extract, and was incubated on ice for 20 mins. The sample was incubated with the radiolabeled probes for 20 mins at room temperature. The complexes were resolved on a 5% polyacrylamide gel at 150V for 3hrs in TBE buffer. Gels were dried and exposed at -80° C for 48 hrs. Negative controls were run with cold oligonucleotides using the labeled probes and binding buffer without the nuclear extract.

Genotyping of NSCLP Probands and Unaffected Controls

A custom TaqMan Genotyping Assay (Applied Biosystems, Foster City, CA) was designed to genotype rs138557689 using our standard protocol and was detected on ABI Viia7 RUO Machine (Applied Biosystems, Foster City, CA). A total of 579 controls and 836 NSCLP probands were genotyped. Allele calls were determined using the Viia7 Software (ABI) and the allele call rate was greater than 98%.

RESULTS

Sequencing of the coding and 5' and 3' UTR regions for *FZD6*, *MATN2*, and *SLC25A32* identified three sequence variants in III-15 and IV-2 (Fig. 4). One sequence change, a homozygous deletion of nucleotide T (c.*164delT), was identified in the 3'UTR of *SLC25A32*. This sequence variation is listed in dbSNP as rs11345830, but the frequency in the general population was not available. However, this homozygous deletion is highly conserved among primates, is not predicted to affect microRNA binding sites, and was therefore excluded from further analysis. One variant, rs113199627 was identified in the 5'UTR of *MATN2*. This is a common variant in the population (MAF=14.5%) and was also excluded from further analysis.

A single base pair c.-153+432A>C change in *FZD6* was present in both individuals (Fig. 4). Rs138557689, a SNP in intron 1, is located 681bps upstream of the start site.




Figure 4. Segregation of rs138557689 in the African American NSCLP family. A large African American family with 11 affected individuals across 3 generations is shown. Filled symbols denote affected individuals. Asterisks (*) denote individuals included in the genome scan reported in Chiquet et al., 2009. The C allele of rs138557689 segregates with the affected individuals and is transmitted by 4 unaffected individuals originating from individual I-3. The C allele is present in 5 additional unaffected individuals.

Based on the availability of DNA samples, 24 additional relatives were sequenced for the same variant. The C allele was found to segregate with the cleft phenotype where 100% of the affected individuals had the CA genotype (Fig 4: II-9, III-2, III-15, III-17, IV-2, IV-6, IV-8). The C allele was transmitted by 4 unaffected individuals (Fig. 4: I-3, II-13, III-4, III-11) and there were 5 other unaffected family members with the CA genotype (Fig. 4: II-12, III-13, IV-5, IV-10, IV-12). The CA genotype was present in 60% of the unaffected family members. There were no affected individuals, however, with the AA genotype.

Despite being submitted by the 1000 Genomes project and listed in NCBI, frequency data on rs138557689 variant was not available. To assess the frequency of this allele, we genotyped 579 controls of Non-Hispanic White (NHW), Hispanic, and African American ethnicity, and found the C allele in 3 individuals (1 NHW and 2 African Americans) who were heterozygous. No homozygous individuals for the C allele were found. The frequency of the C allele was 0.8% in the African American, 0.3% in NHW, and 0% in Hispanic controls (Table 2A). We then genotyped 836 probands from our NSCLP dataset

Table 2. Frequency of the rs138557689/C allele

A. Controls

Alleles	NHW	African A	Hisp	TOTAL
A	299	248	608	1155
C	1	2	0	3
<i>C allele frequency</i>	<i>0.3%</i>	<i>0.8%</i>	<i>0.0%</i>	<i>0.26%</i>

B. NSCLP

Alleles	NHW	African A	Hisp	TOTAL
A	918	168	584	1670
C	0	2	0	2
<i>C allele frequency</i>	<i>0%</i>	<i>1.2%</i>	<i>0%</i>	<i>0.12%</i>

NHW= Non-Hispanic White, African A= African American, Hisp=Hispanic

and found the C allele in two African American individuals, who were both heterozygous. This includes one African American proband from the family described here. The C allele was not found in the NHW and Hispanic

NSCLP probands. The frequency of the C allele in the NSCLP dataset was 1.2% for the African Americans and 0% for the NHW and Hispanics (Table 2B).

To assess whether this SNP was in a conserved region, we compared the sequence in different species. The ancestral A allele of rs138557680 showed conservation in chimps, rhesus monkeys, gorillas, marmoset, mouse lemur, and opossum while the alternate C allele was only conserved in elephants and armadillos (Fig. 5). These results suggest that C is a rare variant. The position of the rs138557689 variant showed moderate evolutionary constraint (GERP score=-2.98). This suggests that the A allele is moderately conserved at this location and that nucleotide changes at this position are tolerated and not selected against through evolution.





Figure 5. Conservation analysis of rs138557689. The UCSC Genome Browser Multiz Alignments of 46 Vertebrates track depicting the nucleotide alignment of the human *FZD6* gene and homologues from other species. The position of the rs138557689 variant is indicated by the arrow. The rs138557689 A allele is conserved. Mouse, rat, *X. tropicalis*, and zebrafish lack the region containing this variant.

In silico analyses predicted that the alternate C allele alters DNA binding. The ancestral allele was predicted to create a C/EB alpha transcription factor binding site by Patch. The alternate allele was predicted to create a Sp1 site by AliBaba2, and NF-1/L binding site by Patch and NIP and NF-1/L binding sites by TESS. EMSA analysis was used to determine whether a DNA binding site was present with either allele. As shown in Fig. 6, there were two bands present for the probe containing the C allele but not for the A allele. These results indicate that the C allele creates new DNA binding sites.



DISCUSSION

We identified a potentially functional variant, rs138557689, in intron 1 of *FZD6* gene segregating with NSCLP phenotype in a large African American family. *In silico* analysis suggested and EMSA results showed that the alternate C allele creates transcription binding sites. Additionally, the ancestral A allele is highly conserved in primates and moderately conserved in other vertebrates, suggesting that it may have important functional significance in the regulation of *FZD6*. While the CA genotype for the rs138557689 variant appears to contribute to the development of NSCLP in this family, analyses are underway to assess how the variant affects gene expression.


The rs138557689/C allele is rare in the general population and was found in less than 1% of NHW and African American controls. The C allele was also rare in the NSCLP probands being identified in only two of 1672 African American chromosomes. Even though there appears to be an enrichment for the C allele in the African American population in our data set, there was no significant difference in the C allele frequency between the African American controls and NSLCP probands (p-value=1.0, Fisher exact test). Based on this information the variant does not appear to be relevant at the population level. However, the large number of individuals with the C allele in the African American family suggests that the rare rs138557689/C genotype may act as a predisposing risk factor for NSCLP when superimposed on the genetic background present in this family. The presence of the C allele in the unaffected family members as well as controls suggests that a person must have a specific combination of variants in one or multiple genes to exhibit the phenotype. This is consistent with the multifactorial model and is supported by the observed reduced penetrance in this African American family. Our results also suggest that different

ethnic-specific genetic risk profiles may contribute to NSCLP, and therefore the underlying disease mechanisms may also be different between ethnic groups. This is consistent with the observed differences in prevalence of NSCLP among different populations and the observed genetic heterogeneity that characterizes this complex disorder. Ethnic-specific differences warrant further studies.

Variants in noncoding regions that modify transcription and expression have increasingly been implicated in complex diseases because they perturb transcription by creating or removing binding sites for specific transcription factors [21,117,118]. We found that the presence of the rs138557689/C allele creates DNA binding sites that could potentially repress or enhance the expression of *FZD6*. Because the development of facial structures is a highly regulated process that relies on crosstalk and convergence of different developmental pathways, alterations in the expression of *FZD6* could affect the cascade of molecular events that the gene is involved in.

While the exact function of *FZD6* in human craniofacial development is not well defined, *FZD6* is diffusely expressed in the craniofacial mesenchyme of zebrafish and chick suggesting that it plays a role during craniofacial development [119,120]. *FZD6* is part of the *Frizzled* family of genes which encode a group of G-coupled receptors critical for initiation of WNT signaling [66,67]. WNT signaling is a highly controlled cellular pathway that regulates multiple functions during development through both canonical/ β -catenin and noncanonical signal transduction [76,77]. Importantly, WNT signaling has been shown to play a role in craniofacial development, including regional specification in the vertebrate face, neural crest induction and differentiation, and mid-face development and upper lip fusion in mice [26,80,81,82,83,84,85,121]. Craniofacial abnormalities, including orofacial

clefts, are found in WNT knock-out mice and zebrafish, while mutations in *WNT3* cause tetra-amelia with CLP in humans [86,87,122]. Additionally, associations between WNTs (*WNT3*, *WNT3A*, *WNT5A*, *WNT7A*, *WNT8A*, *WNT9B* and *WNT11*) and NSCLP have been found, with strongest associations seen in *WNT3* and *WNT3A* [87,88,89,90,123]. Unlike the other Frizzleds, FZD6 represses canonical WNT signaling through the noncanonical Ca²⁺/CaMKII pathway by down-regulating TCF/LEF binding activity and subsequent transcription of WNT target genes (Fig. 7) [70].



An increase in the expression of *FZD6* may further repress canonical WNT signaling crucial for normal craniofacial development while a decrease in *FZD6* expression may cause a loss of negative regulation of the WNT pathway. Negative regulation of the WNT pathway is equally important in craniofacial development since mice that are deficient in the *Dkk1* gene (another negative regulator of the WNT pathway) lack craniofacial structures [124].

FZD6 has also been shown to mediate the non-canonical planar cell polarity (WNT/PCP) pathway in mice (Fig. 7) [72,73,125]. The WNT/PCP pathway controls the polarity and orientation of the migrating neural crest cells [81,126]. During normal craniofacial development NCCs migrate from the neural folds to fill the facial prominences with mesenchyme and contribute to the formation of the nose and upper lip [20,26,28,127,128]. Defects in NCC formation, induction, differentiation, or migration can result in craniofacial abnormalities [129,130,131,132]. Therefore, a decrease in *FZD6* expression may perturb the WNT/PCP pathway and alter neural cell migration or mesenchymal planar cell polarity in craniofacial structures and may potentially lead to NSCLP.

In summary, we report that a rare genetic variant, rs138557689, which alters DNA binding in *FZD6*, may act as a predisposing risk factor for NSCLP in a large African American family with 11 cases of NSCLP segregating in 3 generations. The alteration in *FZD6* gene regulation may perturb highly controlled biological pathways, in particular the WNT pathway, therefore focusing on the genes within the WNT pathway and their interaction with *FZD6* may provide further insight into the genetic etiology of NSCLP. Assessing the affected and unaffected family members who have the CA genotype for the presence of other variants in NSCLP susceptibility genes may delineate the underlying

genetic risk profile contributing to development of NSCLP in this family. This outcome would not only improve genetic counseling for this family but may improve the understanding of the biological mechanisms linking *FZD6* to NSCLP.

APPENDIX

Supplementary Table 1. *FZD6* Primers

Name	Forward	Reverse	Length (bp)	Tm(°C)
E1A	CACTGCTACCTGAGCATCCA	GGA AGC CAC CTC CAC CTT	318	65.1
E1B	AGA GCC AGC GCC AAG AGC TTC A	AGG GTG TAG TGC TGC CGT CGA AA	874	65.1
E1C	GGG AAC CGG CTC TGA AAG GCG	CCT TAA TAA GCT TTC CAA CAG GGG CCC	466	69.4
E1D	CTA GTT GGC CTT ACG AAA ATC GAG	GAC CCA GGA CTC ATT TTC AGG	514	60.2
E2	CAAATGTTGCTGATACACCCTC	CACAACTTGAAGAAATCGGCTC	473	54.3
E3	GAGTTCATAAGTCTGATAGAGGG	CTGTAAGTTCCTGAGAGCAAG	544	54.3
E4A	CCCCATTAACAGCCACAAGTTTT	CGCCTAGCAAAAATCCAATGAAG	593	54.3
E4B	GCAACTCTGTTACATTCCTTAC	GTACAAAGTAGCGAGAAGCATCC	481	54.3
E4C	GCAGTGTGGTTTCATGCTGTTG	GGCTCTTGTTATTTCTCACCTG	538	54.3
E5	GATAAAAAATGTGTTGCACTTAGAGC	GATAAAAAATGTGTTGCACTTAGAGC	486	54.3
E6A	CTGACAAAAGCAACTTAGAGTG	CCTCTCTCATTGATGTTTCTGG	472	54.3
E6C	GGTCATTTCCAAATCCATGGG	CAAACCTCCTTGGGGTAAGAG	427	54.3
E7A	GATAAAGGTGGACACTGGTTAG	GAAGATTCTCTAACTCTGTCC	530	54.3
E7B	GCATTGCCTACTGTTATACTGG	GGACACTCTTTGAGTAAGCACC	592	54.3
E7C	CCCACTTATTGATACCTTACCATC	CAAAAGCATCAGAAAATCTTGCCC	576	54.3
E7D	GCCAATCAAATGGAAAAAAGGTAG	GCATTCTCCTTCAAGGTTAAAAAG	450	54.3

E=Exon

*Betaine was added to primer E1B during the PCR reaction.

*DMSO and magnesium chloride were added to primer E1C during the PCR reaction.

Supplementary Table 2. *MATN2* Primers

Name	Forward	Reverse	Length (bp)	T _m (°C)
E1A	GCT ACT CTG AGG CAG GAG AAT GG	GGC TAA TTG ATG ACA AGC GCC AAG	572	63.5
E1B	GAC TTT CCC TGC TCC CTC GGG GTA GG	C CGG GCG GAA GGA GGG GTG TCC	538	65.1
E1C	GGGAGCGCTCTGGGATGGGAC	CCT GTC CTC GGA GGG GTC GAA G	374	60.2
E1D	CAT CCCCCG C GTC AGT GGG TGC	GAG GAC GGA GCC CCA GCT TCT	499	63.1
E2	CGGCATTTTGAGTGCAAGTGG	CCAGCACATCCAACAAGGAC	417	63.5
E3A	GTAGAGACAGGGTTTTACAGTG	CCTCTGCTTCTGAGAATGCT	465	63.5
E3B	CGTGCTGTCAAGAGGATGCG	CCTATTGCCAGCACAGTACCTG	459	63.5
E4	CTGCCCAGAGGAGAGAGTAG	GGGATGATAACTGGGGTGGG	342	60.2
E5	GTTACTTTTGGTGAGGGCTCTG	CTGAGCACTTTGTGGAAACCC	352	63.5
E6	GCCCTCATCCTACCATTCCC	GTTGAGATACAGTGGTCTATGGTC	383	63.5
E7	CCCCTTTCATGGTGTGACTC	GCTTAAGATTTGCACGAAAAGG	338	60.2
E8	GAATGATCCGTCCTGGCTTG	GACCATCAGGAAACCCGGTG	375	63.5
E9	GACACCTTCCCTGTGGCTTG	GACACCTTCCCTGTGGCTTG	305	63.5
E10	GCATGCCTTCGAGGGAGGGC	GGCCCAGCAAAGCCTGGAGAC	315	63.5
E11	GGCCCAGCAAAGCCTGGAGAC	CCATTTTCTCTTTCAGGGTCC	400	60.2
E12	CCATGGACCACTGAGCTCAGG	GAGCACCAAGGCAGGAAGT	379	63.5
E13	CCCCATCCTGAGTATGAGAC	GCAGTTAAGCCCTCTCTTCC	440	63.5
E14	GGTCCCCAAAGTGGTTATGCC	CTGTACACTCAAAGAAGCTTACCC	625	63.5
E15	GTTGGCATGGACTCTTCAAATC	GCTGAAGCATGAGAATCACTTG	446	63.5
E16	GCTTTTTGAATCTTTGGTGTACC	GTAAATGTAACTATCTTTTGAGGG	372	63.5
E17	CCAAGTGCTAGGAATACCAAG	CTATTTCTACTTCTTCTTTCC	338	56.2
E18	GGTATTACTGGATCTGGCTGC	CTTGCCCTTGCCTCTCAAAC	425	63.5
E19A	GAG TTC TAC AAA TTT ACA AGT CAG GG	CA AAT GAG ATT GCA CAC TAA GGC	542	56.2
E19B	GTGAGAATGAATAAGCTATGCAAG	CCACATCCTTACCAACACTTG	562	62
E19C	GAG AAA TGG CCA ACA TGC CTA TGA AAA AAA TGC TG	CTA CAG GCA CCC ACT ACC ACA CCC G	595	68.2

E=Exon

DMSO and magnesium chloride were added to primer E2 during the PCR reaction.

Supplementary Table 3. *SLC25A32* Primers

Name	Forward	Reverse	Length (bp)	Tm(°C)
E1A	CAT CTC GGT TGC TCT TCC GGC	CGC CCC TTG TGA GCG CAA C	431	67
E1B	CCC CTC CAT CGC GCT TTC CG	CAG GTT AGC CAA CGC GGA CAG	513	67
E2	CGG GAC TCT GAC ACA AAA ATG	GCC ATT TTG TTC TTC CTG ATC C	519	62
E3	GCA AAC CAC TTC CAG CAA ATT C	CTG AGG CAG GAT TAT CGC TTG	530	58.3
E4	CCT TTT GAC CCT AAG ACT GTG C	GAT ACA TCC TAT GTT AAT GGG GAC	537	58.3
E5	GTG CTG TAA TGA GAG AGA AGA G	GAC ATT TGT GTG GCT TCA	374	58.3
E6	GAT GAG TAC TGG CTC TGC CA	CGT AAT AAC TGG GAA AGC AGG	473	58.3
E7A	GAC TGC TGC TTG CTC CAT GCC	GGC AGC CAT TTC AGG CAG AGG	454	58.3
E7B	GAA GCC AGA GAA CTG CTA AGT C	CGA CAA AGC AAG ACT CCA TCT C	404	62
E7C	GTG AGC TTA CTT GCC TGG ATT GC	CGC TAG GTA GTG CAT CCC AAC TG	543	62
E7D	GGATGGTCTCAATCTCCTGAC	CAA CCT GAA TTT GAG AAA CCA ATG AAG	564	58.3
E7E	GTTGACAAGGTAAATGGAAATGAG	GTT CAC TTT TTC CGT GTG GGG	517	58.3
E7F	GCT GTT GCA CTA CCA TCT ATT TG	CTC AGT GCT TGG TGA CGT AC	556	52

E=Exon

REFERENCES

- [1] S.S. Hashmi, D.K. Waller, P. Langlois, M. Canfield, J.T. Hecht, Prevalence of nonsyndromic oral clefts in Texas: 1995-1999, *Am J Med Genet A* 134 (2005) 368-372.
- [2] S.E. Parker, C.T. Mai, M.A. Canfield, R. Rickard, Y. Wang, R.E. Meyer, P. Anderson, C.A. Mason, J.S. Collins, R.S. Kirby, A. Correa, Updated National Birth Prevalence estimates for selected birth defects in the United States, 2004-2006, *Birth Defects Res A Clin Mol Teratol* 88 (2010) 1008-1016.
- [3] Improved national prevalence estimates for 18 selected major birth defects--United States, 1999-2001, *MMWR Morb Mortal Wkly Rep* 54 (2006) 1301-1305.
- [4] T. Demir, G. Karacetin, S. Baghaki, Y. Aydin, Psychiatric assessment of children with nonsyndromic cleft lip and palate, *Gen Hosp Psychiatry* 33 (2011) 594-603.
- [5] R.P. Strauss, "Only skin deep": health, resilience, and craniofacial care, *Cleft Palate Craniofac J* 38 (2001) 226-230.
- [6] G.L. Wehby, C.H. Cassell, The impact of orofacial clefts on quality of life and healthcare use and costs, *Oral Dis* 16 (2010) 3-10.
- [7] K. Christensen, K. Juel, A.M. Herskind, J.C. Murray, Long term follow up study of survival associated with cleft lip and palate at birth, *BMJ* 328 (2004) 1405.
- [8] P.A. Mossey, J. Little, R.G. Munger, M.J. Dixon, W.C. Shaw, Cleft lip and palate, *Lancet* 374 (2009) 1773-1785.
- [9] R.J. Gorlin, Cohen, M.M., Hennekam, R.C.M, *Syndromes of the Head and Neck*, Fourth Edition ed., Oxford University Press, New York 2011.
- [10] K. Boo-Chai, An ancient Chinese text on a cleft lip, *Plast Reconstr Surg* 38 (1966) 89-91.

- [11] L.H. Cohen, Fine, B.A., Pergament, E., An Assessment of Ethnocultural Beliefs Regarding the Causes of Birth Defects and Genetic Disorders *Journal of Genetic Counseling* 7 (1998).
- [12] L.R. Cheng, Asian-American cultural perspectives on birth defects: focus on cleft palate, *Cleft Palate J* 27 (1990) 294-300.
- [13] G. Toliver-Weddington, Cultural considerations in the treatment of craniofacial malformations in African Americans, *Cleft Palate J* 27 (1990) 289-293.
- [14] M.D. Meyerson, Cultural considerations in the treatment of Latinos with craniofacial malformations, *Cleft Palate J* 27 (1990) 279-288.
- [15] L.F. Snow, S.M. Johnson, H.E. Mayhew, The behavioral implications of some old wives' tales, *Obstet Gynecol* 51 (1978) 727-732.
- [16] M.E. Burk, P.C. Wieser, L. Keegan, Cultural beliefs and health behaviors of pregnant Mexican-American women: implications for primary care, *ANS Adv Nurs Sci* 17 (1995) 37-52.
- [17] R.C. Weatherley-White, W. Eiserman, M. Beddoe, R. Vanderberg, Perceptions, expectations, and reactions to cleft lip and palate surgery in native populations: a pilot study in rural India, *Cleft Palate Craniofac J* 42 (2005) 560-564.
- [18] H.O. Olasoji, V.I. Ugboko, G.T. Arotiba, Cultural and religious components in Nigerian parents' perceptions of the aetiology of cleft lip and palate: implications for treatment and rehabilitation, *Br J Oral Maxillofac Surg* 45 (2007) 302-305.
- [19] B.C. Schutte, J.C. Murray, The many faces and factors of orofacial clefts, *Hum Mol Genet* 8 (1999) 1853-1859.

- [20] K.L. Moore, Persaud, T.V.N., The Developing Human: Clinically Oriented Embryology, 8th ed., Saunders Elsevier, Philadelphia, 2008.
- [21] D.F. Wyszynski, Cleft Lip & Palate: From Origin to Treatment, Oxford University Press, New York, 2002.
- [22] P.S. Harper, Practical Genetic Counselling, Sixth Ed. ed., Edward Arnold Ltd., London, 2006.
- [23] R.A. Spritz, The genetics and epigenetics of orofacial clefts, *Curr Opin Pediatr* 13 (2001) 556-560.
- [24] M.M. Tolarova, J. Cervenka, Classification and birth prevalence of orofacial clefts, *Am J Med Genet* 75 (1998) 126-137.
- [25] M.M. Cohen, Etiology and pathogenesis of orofacial clefting, *Oral Maxillofac Clin North Am* (2000) 379-397.
- [26] R. Jiang, J.O. Bush, A.C. Lidral, Development of the upper lip: morphogenetic and molecular mechanisms, *Dev Dyn* 235 (2006) 1152-1166.
- [27] A.C. Lidral, L.M. Moreno, Progress toward discerning the genetics of cleft lip, *Curr Opin Pediatr* 17 (2005) 731-739.
- [28] C.W. Senders, E.C. Peterson, A.G. Hendrickx, M.A. Cukierski, Development of the upper lip, *Arch Facial Plast Surg* 5 (2003) 16-25.
- [29] J.C. Murray, Face facts: genes, environment, and clefts, *Am J Hum Genet* 57 (1995) 227-232.
- [30] M.C. Johnston, P.T. Bronsky, Prenatal craniofacial development: new insights on normal and abnormal mechanisms, *Crit Rev Oral Biol Med* 6 (1995) 368-422.

- [31] Parameters for evaluation and treatment of patients with cleft lip/palate or other craniofacial anomalies. American Cleft Palate-Craniofacial Association. March, 1993, Cleft Palate Craniofac J 30 Suppl (1993) S1-16.
- [32] F. Manna, S. Pensiero, G. Clarich, G.F. Guarneri, P.C. Parodi, Cleft lip and palate: current status from the literature and our experience, J Craniofac Surg 20 (2009) 1383-1387.
- [33] J.L. Marsh, Craniofacial surgery: the experiment on the experiment of nature, Cleft Palate Craniofac J 33 (1996) 1 p.
- [34] D.F. Wyszynski, T.H. Beaty, N.E. Maestri, Genetics of nonsyndromic oral clefts revisited, Cleft Palate Craniofac J 33 (1996) 406-417.
- [35] K. Christensen, P. Fogh-Andersen, Cleft lip (+/- cleft palate) in Danish twins, 1970-1990, Am J Med Genet 47 (1993) 910-916.
- [36] A.C. Lidral, J.C. Murray, Genetic approaches to identify disease genes for birth defects with cleft lip/palate as a model, Birth Defects Res A Clin Mol Teratol 70 (2004) 893-901.
- [37] L.E. Mitchell, N. Risch, Mode of inheritance of nonsyndromic cleft lip with or without cleft palate: a reanalysis, Am J Hum Genet 51 (1992) 323-332.
- [38] A. Sivertsen, A.J. Wilcox, R. Skjaerven, H.A. Vindenes, F. Abyholm, E. Harville, R.T. Lie, Familial risk of oral clefts by morphological type and severity: population based cohort study of first degree relatives, BMJ 336 (2008) 432-434.
- [39] D. Groesen, C. Bille, J.K. Pedersen, A. Skytthe, J.C. Murray, K. Christensen, Recurrence risk for offspring of twins discordant for oral cleft: a population-based cohort study

- of the Danish 1936-2004 cleft twin cohort, *Am J Med Genet A* 152A (2010) 2468-2474.
- [40] M.A. Honein, S.A. Rasmussen, J. Reefhuis, P.A. Romitti, E.J. Lammer, L. Sun, A. Correa, Maternal smoking and environmental tobacco smoke exposure and the risk of orofacial clefts, *Epidemiology* 18 (2007) 226-233.
- [41] L. Stuppia, M. Capogreco, G. Marzo, D. La Rovere, I. Antonucci, V. Gatta, G. Palka, C. Mortellaro, S. Tete, Genetics of syndromic and nonsyndromic cleft lip and palate, *J Craniofac Surg* 22 (2011) 1722-1726.
- [42] J. Murthy, L. Bhaskar, Current concepts in genetics of nonsyndromic clefts, *Indian J Plast Surg* 42 (2009) 68-81.
- [43] S. Shahrukh Hashmi, M.S. Gallaway, D.K. Waller, P.H. Langlois, J.T. Hecht, Maternal fever during early pregnancy and the risk of oral clefts, *Birth Defects Res A Clin Mol Teratol* 88 (2010) 186-194.
- [44] Q. Yuan, S.H. Blanton, J.T. Hecht, Genetic causes of nonsyndromic cleft lip with or without cleft palate, *Adv Otorhinolaryngol* 70 (2011) 107-113.
- [45] T.M. Zuccherro, M.E. Cooper, B.S. Maher, S. Daack-Hirsch, B. Nepomuceno, L. Ribeiro, D. Caprau, K. Christensen, Y. Suzuki, J. Machida, N. Natsume, K. Yoshiura, A.R. Vieira, I.M. Orioli, E.E. Castilla, L. Moreno, M. Arcos-Burgos, A.C. Lidral, L.L. Field, Y.E. Liu, A. Ray, T.H. Goldstein, R.E. Schultz, M. Shi, M.K. Johnson, S. Kondo, B.C. Schutte, M.L. Marazita, J.C. Murray, Interferon regulatory factor 6 (IRF6) gene variants and the risk of isolated cleft lip or palate, *N Engl J Med* 351 (2004) 769-780.

- [46] L. Scapoli, A. Palmieri, M. Martinelli, F. Pezzetti, P. Carinci, M. Tognon, F. Carinci, Strong evidence of linkage disequilibrium between polymorphisms at the IRF6 locus and nonsyndromic cleft lip with or without cleft palate, in an Italian population, *Am J Hum Genet* 76 (2005) 180-183.
- [47] A. Jugessur, F. Rahimov, R.T. Lie, A.J. Wilcox, H.K. Gjessing, R.M. Nilsen, T.T. Nguyen, J.C. Murray, Genetic variants in IRF6 and the risk of facial clefts: single-marker and haplotype-based analyses in a population-based case-control study of facial clefts in Norway, *Genet Epidemiol* 32 (2008) 413-424.
- [48] S. Birnbaum, K.U. Ludwig, H. Reutter, S. Herms, N.A. de Assis, A. Diaz-Lacava, S. Barth, C. Lauster, G. Schmidt, M. Scheer, M. Saffar, M. Martini, R.H. Reich, F. Schiefke, A. Hemprich, S. Potzsch, B. Potzsch, T.F. Wienker, P. Hoffmann, M. Knapp, F.J. Kramer, M.M. Nothen, E. Mangold, IRF6 gene variants in Central European patients with non-syndromic cleft lip with or without cleft palate, *Eur J Oral Sci* 117 (2009) 766-769.
- [49] Y. Pan, J. Ma, W. Zhang, Y. Du, Y. Niu, M. Wang, Z. Zhang, L. Wang, IRF6 polymorphisms are associated with nonsyndromic orofacial clefts in a Chinese Han population, *Am J Med Genet A* 152A (2010) 2505-2511.
- [50] T. Wu, K.Y. Liang, J.B. Hetmanski, I. Ruczinski, M.D. Fallin, R.G. Ingersoll, H. Wang, S. Huang, X. Ye, Y.H. Wu-Chou, P.K. Chen, E.W. Jabs, B. Shi, R. Redett, A.F. Scott, T.H. Beaty, Evidence of gene-environment interaction for the IRF6 gene and maternal multivitamin supplementation in controlling the risk of cleft lip with/without cleft palate, *Hum Genet* 128 (2010) 401-410.

- [51] S. Kondo, B.C. Schutte, R.J. Richardson, B.C. Bjork, A.S. Knight, Y. Watanabe, E. Howard, R.L. de Lima, S. Daack-Hirsch, A. Sander, D.M. McDonald-McGinn, E.H. Zackai, E.J. Lammer, A.S. Aylsworth, H.H. Ardinger, A.C. Lidral, B.R. Pober, L. Moreno, M. Arcos-Burgos, C. Valencia, C. Houdayer, M. Bahuau, D. Moretti-Ferreira, A. Richieri-Costa, M.J. Dixon, J.C. Murray, Mutations in IRF6 cause Van der Woude and popliteal pterygium syndromes, *Nat Genet* 32 (2002) 285-289.
- [52] M.L. Marazita, J.C. Murray, A.C. Lidral, M. Arcos-Burgos, M.E. Cooper, T. Goldstein, B.S. Maher, S. Daack-Hirsch, R. Schultz, M.A. Mansilla, L.L. Field, Y.E. Liu, N. Prescott, S. Malcolm, R. Winter, A. Ray, L. Moreno, C. Valencia, K. Neiswanger, D.F. Wyszynski, J.E. Bailey-Wilson, H. Albacha-Hejazi, T.H. Beaty, I. McIntosh, J.B. Hetmanski, G. Tuncbilek, M. Edwards, L. Harkin, R. Scott, L.G. Roddick, Meta-analysis of 13 genome scans reveals multiple cleft lip/palate genes with novel loci on 9q21 and 2q32-35, *Am J Hum Genet* 75 (2004) 161-173.
- [53] J. Suazo, J.L. Santos, H. Carreno, L. Jara, R. Blanco, Linkage disequilibrium between MSX1 and non-syndromic cleft lip/palate in the Chilean population, *J Dent Res* 83 (2004) 782-785.
- [54] A.C. Marcano, K. Doudney, C. Braybrook, R. Squires, M.A. Patton, M.M. Lees, A. Richieri-Costa, A.C. Lidral, J.C. Murray, G.E. Moore, P. Stanier, TBX22 mutations are a frequent cause of cleft palate, *J Med Genet* 41 (2004) 68-74.
- [55] E.J. Lammer, G.M. Shaw, D.M. Iovannisci, J. Van Waes, R.H. Finnell, Maternal smoking and the risk of orofacial clefts: Susceptibility with NAT1 and NAT2 polymorphisms, *Epidemiology* 15 (2004) 150-156.

- [56] E.J. Lammer, G.M. Shaw, D.M. Iovannisci, R.H. Finnell, Maternal smoking, genetic variation of glutathione s-transferases, and risk for orofacial clefts, *Epidemiology* 16 (2005) 698-701.
- [57] K.K. Hozyasz, A. Mostowska, Z. Surowiec, P.P. Jagodzinski, [Genetic polymorphisms of GSTM1 and GSTT1 in mothers of children with isolated cleft lip with or without cleft palate], *Przegl Lek* 62 (2005) 1019-1022.
- [58] M. Shi, K. Christensen, C.R. Weinberg, P. Romitti, L. Bathum, A. Lozada, R.W. Morris, M. Lovett, J.C. Murray, Orofacial cleft risk is increased with maternal smoking and specific detoxification-gene variants, *Am J Hum Genet* 80 (2007) 76-90.
- [59] V. Shotelersuk, C. Ittiwut, P. Siriwan, A. Angspatt, Maternal 677CT/1298AC genotype of the MTHFR gene as a risk factor for cleft lip, *J Med Genet* 40 (2003) e64.
- [60] J.R. Avila, P.A. Jezewski, A.R. Vieira, I.M. Orioli, E.E. Castilla, K. Christensen, S. Daack-Hirsch, P.A. Romitti, J.C. Murray, PVRL1 variants contribute to non-syndromic cleft lip and palate in multiple populations, *Am J Med Genet A* 140 (2006) 2562-2570.
- [61] B.T. Chiquet, R. Henry, A. Burt, J.B. Mulliken, S. Stal, S.H. Blanton, J.T. Hecht, Nonsyndromic cleft lip and palate: CRISPLD genes and the folate gene pathway connection, *Birth Defects Res A Clin Mol Teratol* 91 (2011) 44-49.
- [62] E. Mangold, K.U. Ludwig, M.M. Nothen, Breakthroughs in the genetics of orofacial clefting, *Trends Mol Med* (2011).
- [63] F. Pagani, F.E. Baralle, Genomic variants in exons and introns: identifying the splicing spoilers, *Nat Rev Genet* 5 (2004) 389-396.

- [64] S. Birnbaum, K.U. Ludwig, H. Reutter, S. Herms, M. Steffens, M. Rubini, C. Baluardo, M. Ferrian, N. Almeida de Assis, M.A. Alblas, S. Barth, J. Freudenberg, C. Lauster, G. Schmidt, M. Scheer, B. Braumann, S.J. Berge, R.H. Reich, F. Schiefke, A. Hemprich, S. Potzsch, R.P. Steegers-Theunissen, B. Potzsch, S. Moebus, B. Horsthemke, F.J. Kramer, T.F. Wienker, P.A. Mossey, P. Propping, S. Cichon, P. Hoffmann, M. Knapp, M.M. Nothen, E. Mangold, Key susceptibility locus for nonsyndromic cleft lip with or without cleft palate on chromosome 8q24, *Nat Genet* 41 (2009) 473-477.
- [65] B.T. Chiquet, S.S. Hashmi, R. Henry, A. Burt, J.B. Mulliken, S. Stal, M. Bray, S.H. Blanton, J.T. Hecht, Genomic screening identifies novel linkages and provides further evidence for a role of MYH9 in nonsyndromic cleft lip and palate, *Eur J Hum Genet* 17 (2009) 195-204.
- [66] M. Tokuwara, M. Hirai, Y. Atomi, M. Terada, M. Katoh, Molecular cloning of human Frizzled-6, *Biochem Biophys Res Commun* 243 (1998) 622-627.
- [67] B.T. MacDonald, K. Tamai, X. He, Wnt/beta-catenin signaling: components, mechanisms, and diseases, *Dev Cell* 17 (2009) 9-26.
- [68] P. Bhanot, M. Brink, C.H. Samos, J.C. Hsieh, Y. Wang, J.P. Macke, D. Andrew, J. Nathans, R. Nusse, A new member of the frizzled family from *Drosophila* functions as a Wingless receptor, *Nature* 382 (1996) 225-230.
- [69] Z. Songyang, A.S. Fanning, C. Fu, J. Xu, S.M. Marfatia, A.H. Chishti, A. Crompton, A.C. Chan, J.M. Anderson, L.C. Cantley, Recognition of unique carboxyl-terminal motifs by distinct PDZ domains, *Science* 275 (1997) 73-77.

- [70] T. Golan, A. Yaniv, A. Bafico, G. Liu, A. Gazit, The human Frizzled 6 (HFz6) acts as a negative regulator of the canonical Wnt. beta-catenin signaling cascade, *J Biol Chem* 279 (2004) 14879-14888.
- [71] A.S. Frojmark, J. Schuster, M. Sobol, M. Entesarian, M.B. Kilander, D. Gabrikova, S. Nawaz, S.M. Baig, G. Schulte, J. Klar, N. Dahl, Mutations in Frizzled 6 cause isolated autosomal-recessive nail dysplasia, *Am J Hum Genet* 88 (2011) 852-860.
- [72] N. Guo, C. Hawkins, J. Nathans, Frizzled6 controls hair patterning in mice, *Proc Natl Acad Sci U S A* 101 (2004) 9277-9281.
- [73] Y. Wang, N. Guo, J. Nathans, The role of Frizzled3 and Frizzled6 in neural tube closure and in the planar polarity of inner-ear sensory hair cells, *J Neurosci* 26 (2006) 2147-2156.
- [74] P. De Marco, E. Merello, A. Rossi, G. Piatelli, A. Cama, Z. Kibar, V. Capra, FZD6 is a novel gene for human neural tube defects, *Hum Mutat* 33 (2012) 384-390.
- [75] G. Naz, S.M. Pasternack, C. Perrin, M. Mattheisen, M. Refke, S. Khan, A. Gul, M. Simons, W. Ahmad, R.C. Betz, FZD6 encoding the Wnt receptor frizzled-6 is mutated in autosomal-recessive nail dysplasia, *Br J Dermatol* (2011).
- [76] C.Y. Logan, R. Nusse, The Wnt signaling pathway in development and disease, *Annu Rev Cell Dev Biol* 20 (2004) 781-810.
- [77] R.T. Moon, A.D. Kohn, G.V. De Ferrari, A. Kaykas, WNT and beta-catenin signalling: diseases and therapies, *Nat Rev Genet* 5 (2004) 691-701.
- [78] X. He, M. Semenov, K. Tamai, X. Zeng, LDL receptor-related proteins 5 and 6 in Wnt/beta-catenin signaling: arrows point the way, *Development* 131 (2004) 1663-1677.

- [79] Y. Wang, L. Song, C.J. Zhou, The canonical Wnt/beta-catenin signaling pathway regulates Fgf signaling for early facial development, *Dev Biol* 349 (2011) 250-260.
- [80] S.A. Brugmann, L.H. Goodnough, A. Gregorieff, P. Leucht, D. ten Berge, C. Fuerer, H. Clevers, R. Nusse, J.A. Helms, Wnt signaling mediates regional specification in the vertebrate face, *Development* 134 (2007) 3283-3295.
- [81] J. De Calisto, C. Araya, L. Marchant, C.F. Riaz, R. Mayor, Essential role of non-canonical Wnt signalling in neural crest migration, *Development* 132 (2005) 2587-2597.
- [82] O. Mazemondet, R. Hubner, J. Frahm, D. Koczan, B.M. Bader, D.G. Weiss, A.M. Uhrmacher, M.J. Frech, A. Rolfs, J. Luo, Quantitative and kinetic profile of Wnt/beta-catenin signaling components during human neural progenitor cell differentiation, *Cell Mol Biol Lett* 16 (2011) 515-538.
- [83] B.S. Reid, H. Yang, V.S. Melvin, M.M. Taketo, T. Williams, Ectodermal Wnt/beta-catenin signaling shapes the mouse face, *Dev Biol* 349 (2011) 261-269.
- [84] L. Song, Y. Li, K. Wang, Y.Z. Wang, A. Molotkov, L. Gao, T. Zhao, T. Yamagami, Y. Wang, Q. Gan, D.E. Pleasure, C.J. Zhou, Lrp6-mediated canonical Wnt signaling is required for lip formation and fusion, *Development* 136 (2009) 3161-3171.
- [85] P. Mani, A. Jarrell, J. Myers, R. Atit, Visualizing canonical Wnt signaling during mouse craniofacial development, *Dev Dyn* 239 (2010) 354-363.
- [86] C.P. Heisenberg, M. Tada, G.J. Rauch, L. Saude, M.L. Concha, R. Geisler, D.L. Stemple, J.C. Smith, S.W. Wilson, Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation, *Nature* 405 (2000) 76-81.

- [87] D.M. Juriloff, M.J. Harris, A.P. McMahon, T.J. Carroll, A.C. Lidral, Wnt9b is the mutated gene involved in multifactorial nonsyndromic cleft lip with or without cleft palate in A/WySn mice, as confirmed by a genetic complementation test, *Birth Defects Res A Clin Mol Teratol* 76 (2006) 574-579.
- [88] B.T. Chiquet, S.H. Blanton, A. Burt, D. Ma, S. Stal, J.B. Mulliken, J.T. Hecht, Variation in WNT genes is associated with non-syndromic cleft lip with or without cleft palate, *Hum Mol Genet* 17 (2008) 2212-2218.
- [89] R. Menezes, A. Letra, A.H. Kim, E.C. Kuchler, A. Day, P.N. Tannure, L. Gomes da Motta, K.B. Paiva, J.M. Granjeiro, A.R. Vieira, Studies with Wnt genes and nonsyndromic cleft lip and palate, *Birth Defects Res A Clin Mol Teratol* 88 (2010) 995-1000.
- [90] T. Yao, L. Yang, P.Q. Li, H. Wu, H.B. Xie, X. Shen, X.D. Xie, Association of Wnt3A gene variants with non-syndromic cleft lip with or without cleft palate in Chinese population, *Arch Oral Biol* 56 (2011) 73-78.
- [91] F. Deak, R. Wagener, I. Kiss, M. Paulsson, The matrilins: a novel family of oligomeric extracellular matrix proteins, *Matrix Biol* 18 (1999) 55-64.
- [92] A.R. Klatt, A.K. Becker, C.D. Neacsu, M. Paulsson, R. Wagener, The matrilins: modulators of extracellular matrix assembly, *Int J Biochem Cell Biol* 43 (2011) 320-330.
- [93] S. Muratoglu, K. Krysan, M. Balazs, H. Sheng, R. Zakany, L. Modis, I. Kiss, F. Deak, Primary structure of human matrilin-2, chromosome location of the MATN2 gene and conservation of an AT-AC intron in matrilin genes, *Cytogenet Cell Genet* 90 (2000) 323-327.

- [94] L. Mates, E. Korpos, F. Deak, Z. Liu, D.R. Beier, A. Aszodi, I. Kiss, Comparative analysis of the mouse and human genes (Matn2 and MATN2) for matrilin-2, a filament-forming protein widely distributed in extracellular matrices, *Matrix Biol* 21 (2002) 163-174.
- [95] D. Piecha, K. Hartmann, B. Kobbe, I. Haase, C. Mauch, T. Krieg, M. Paulsson, Expression of matrilin-2 in human skin, *J Invest Dermatol* 119 (2002) 38-43.
- [96] R. Wagener, H.W. Ehlen, Y.P. Ko, B. Kobbe, H.H. Mann, G. Sengle, M. Paulsson, The matrilins--adaptor proteins in the extracellular matrix, *FEBS Lett* 579 (2005) 3323-3329.
- [97] L. Mates, C. Nicolae, M. Morgelin, F. Deak, I. Kiss, A. Aszodi, Mice lacking the extracellular matrix adaptor protein matrilin-2 develop without obvious abnormalities, *Matrix Biol* 23 (2004) 195-204.
- [98] P. Carinci, E. Becchetti, T. Baroni, F. Carinci, F. Pezzetti, G. Stabellini, P. Locci, L. Scapoli, M. Tognon, S. Volinia, M. Bodo, Extracellular matrix and growth factors in the pathogenesis of some craniofacial malformations, *Eur J Histochem* 51 Suppl 1 (2007) 105-115.
- [99] R. Perris, D. Perissinotto, Role of the extracellular matrix during neural crest cell migration, *Mech Dev* 95 (2000) 3-21.
- [100] S.A. Titus, R.G. Moran, Retrovirally mediated complementation of the glyB phenotype. Cloning of a human gene encoding the carrier for entry of folates into mitochondria, *J Biol Chem* 275 (2000) 36811-36817.

- [101] M.A. Hediger, M.F. Romero, J.B. Peng, A. Rolfs, H. Takanaga, E.A. Bruford, The ABCs of solute carriers: physiological, pathological and therapeutic implications of human membrane transport proteins Introduction, *Pflugers Arch* 447 (2004) 465-468.
- [102] T. Haitina, J. Lindblom, T. Renstrom, R. Fredriksson, Fourteen novel human members of mitochondrial solute carrier family 25 (SLC25) widely expressed in the central nervous system, *Genomics* 88 (2006) 779-790.
- [103] A. Biswas, S.R. Senthilkumar, H.M. Said, Effect of chronic alcohol exposure on folate uptake by liver mitochondria, *Am J Physiol Cell Physiol* (2011).
- [104] L.D. Botto, A. Lisi, C. Bower, M.A. Canfield, N. Dattani, C. De Vigan, H. De Walle, D.J. Erickson, J. Halliday, L.M. Irgens, R.B. Lowry, R. McDonnell, J. Metneki, S. Poetzsch, A. Ritvanen, E. Robert-Gnansia, C. Siffel, C. Stoll, P. Mastroiacovo, Trends of selected malformations in relation to folic acid recommendations and fortification: an international assessment, *Birth Defects Res A Clin Mol Teratol* 76 (2006) 693-705.
- [105] S.H. Blanton, R.R. Henry, Q. Yuan, J.B. Mulliken, S. Stal, R.H. Finnell, J.T. Hecht, Folate pathway and nonsyndromic cleft lip and palate, *Birth Defects Res A Clin Mol Teratol* 91 (2011) 50-60.
- [106] G.L. Wehby, J.C. Murray, Folic acid and orofacial clefts: a review of the evidence, *Oral Dis* 16 (2010) 11-19.
- [107] R.L. Badovinac, M.M. Werler, P.L. Williams, K.T. Kelsey, C. Hayes, Folic acid-containing supplement consumption during pregnancy and risk for oral clefts: a meta-analysis, *Birth Defects Res A Clin Mol Teratol* 79 (2007) 8-15.

- [108] C.Y. Johnson, J. Little, Folate intake, markers of folate status and oral clefts: is the evidence converging?, *Int J Epidemiol* 37 (2008) 1041-1058.
- [109] N. Grabe, AliBaba2: context specific identification of transcription factor binding sites, *In Silico Biol* 2 (2002) S1-15.
- [110] V. Matys, O.V. Kel-Margoulis, E. Fricke, I. Liebich, S. Land, A. Barre-Dirrie, I. Reuter, D. Chekmenev, M. Krull, K. Hornischer, N. Voss, P. Stegmaier, B. Lewicki-Potapov, H. Saxel, A.E. Kel, E. Wingender, TRANSFAC and its module TRANSCompel: transcriptional gene regulation in eukaryotes, *Nucleic Acids Res* 34 (2006) D108-110.
- [111] J. Schug, Using TESS to Predict Transcription Factor Binding Sites in DNA Sequence, *Curr. Protoc. Bioinform.* 21 (2008) 2.6.1-2.6.15.
- [112] S. Griffiths-Jones, H.K. Saini, S. van Dongen, A.J. Enright, miRBase: tools for microRNA genomics, *Nucleic Acids Res* 36 (2008) D154-158.
- [113] S.D. Hsu, C.H. Chu, A.P. Tsou, S.J. Chen, H.C. Chen, P.W. Hsu, Y.H. Wong, Y.H. Chen, G.H. Chen, H.D. Huang, miRNAmap 2.0: genomic maps of microRNAs in metazoan genomes, *Nucleic Acids Res* 36 (2008) D165-169.
- [114] I.A. Adzhubei, S. Schmidt, L. Peshkin, V.E. Ramensky, A. Gerasimova, P. Bork, A.S. Kondrashov, S.R. Sunyaev, A method and server for predicting damaging missense mutations, *Nat Methods* 7 (2010) 248-249.
- [115] P.C. Ng, S. Henikoff, Predicting deleterious amino acid substitutions, *Genome Res* 11 (2001) 863-874.

- [116] G.M. Cooper, E.A. Stone, G. Asimenos, E.D. Green, S. Batzoglou, A. Sidow,
Distribution and intensity of constraint in mammalian genomic sequence, *Genome Res* 15 (2005) 901-913.
- [117] I. Thesleff, The genetic basis of normal and abnormal craniofacial development, *Acta Odontol Scand* 56 (1998) 321-325.
- [118] X. Wang, D.J. Tomso, X. Liu, D.A. Bell, Single nucleotide polymorphism in
transcriptional regulatory regions and expression of environmentally responsive
genes, *Toxicol Appl Pharmacol* 207 (2005) 84-90.
- [119] P. Geetha-Loganathan, S. Nimmagadda, L. Antoni, K. Fu, C.J. Whiting, P. Francis-
West, J.M. Richman, Expression of WNT signalling pathway genes during chicken
craniofacial development, *Dev Dyn* 238 (2009) 1150-1165.
- [120] B.E. Sisson, J. Topczewski, Expression of five frizzleds during zebrafish craniofacial
development, *Gene Expr Patterns* 9 (2009) 520-527.
- [121] D. Wang, F. Liu, L. Wang, S. Huang, J. Yu, Nonsynonymous substitution rate (K_a) is
a relatively consistent parameter for defining fast-evolving and slow-evolving
protein-coding genes, *Biol Direct* 6 (2011) 13.
- [122] S. Niemann, C. Zhao, F. Pascu, U. Stahl, U. Aulepp, L. Niswander, J.L. Weber, U.
Muller, Homozygous WNT3 mutation causes tetra-amelia in a large consanguineous
family, *Am J Hum Genet* 74 (2004) 558-563.
- [123] A. Mostowska, K.K. Hozyasz, B. Biedziak, P. Wojcicki, M. Lianeri, P.P. Jagodzinski,
Genotype and haplotype analysis of WNT genes in non-syndromic cleft lip with or
without cleft palate, *Eur J Oral Sci* 120 (2012) 1-8.

- [124] M. Mukhopadhyay, Shtrom, S., Rodriguez-Esteban, C., Chen, L., Tsukui, T., Gomer, L., Dorward, D.W., Glinka, A., Grinberg, A., Huang, S.P., Niehrs, C., Izpisua Belmonte, J.C., Westphal, H., Dickkopf1 is required for embryonic head induction and limb morphogenesis in the mouse., *Dev Cell* 3 (2001) 423-434.
- [125] D. Devenport, E. Fuchs, Planar polarization in embryonic epidermis orchestrates global asymmetric morphogenesis of hair follicles, *Nat Cell Biol* 10 (2008) 1257-1268.
- [126] H.K. Matthews, L. Marchant, C. Carmona-Fontaine, S. Kuriyama, J. Larrain, M.R. Holt, M. Parsons, R. Mayor, Directional migration of neural crest cells in vivo is regulated by Syndecan-4/Rac1 and non-canonical Wnt signaling/RhoA, *Development* 135 (2008) 1771-1780.
- [127] Y. Chai, R.E. Maxson, Jr., Recent advances in craniofacial morphogenesis, *Dev Dyn* 235 (2006) 2353-2375.
- [128] D.R. Cordero, S. Brugmann, Y. Chu, R. Bajpai, M. Jame, J.A. Helms, Cranial neural crest cells on the move: their roles in craniofacial development, *Am J Med Genet A* 155A (2011) 270-279.
- [129] J. Dixon, N.C. Jones, L.L. Sandell, S.M. Jayasinghe, J. Crane, J.P. Rey, M.J. Dixon, P.A. Trainor, Tcof1/Treacle is required for neural crest cell formation and proliferation deficiencies that cause craniofacial abnormalities, *Proc Natl Acad Sci U S A* 103 (2006) 13403-13408.
- [130] V. Garg, C. Yamagishi, T. Hu, I.S. Kathiriya, H. Yamagishi, D. Srivastava, Tbx1, a DiGeorge syndrome candidate gene, is regulated by sonic hedgehog during pharyngeal arch development, *Dev Biol* 235 (2001) 62-73.

- [131] N.C. Jones, M.L. Lynn, K. Gaudenz, D. Sakai, K. Aoto, J.P. Rey, E.F. Glynn, L. Ellington, C. Du, J. Dixon, M.J. Dixon, P.A. Trainor, Prevention of the neurocristopathy Treacher Collins syndrome through inhibition of p53 function, *Nat Med* 14 (2008) 125-133.
- [132] F. Vitelli, M. Morishima, I. Taddei, E.A. Lindsay, A. Baldini, Tbx1 mutation causes multiple cardiovascular defects and disrupts neural crest and cranial nerve migratory pathways, *Hum Mol Genet* 11 (2002) 915-922.

VITA

Nevena Cvjetkovic was born in Bosnia and Herzegovina on January 29th, 1988, the first daughter of Nenad and Ljubenka Cvjetkovic. She immigrated to Florida with her parents and sister Jovana in 1998 at the age of 10. After graduation from Lakewood High School in St. Petersburg, FL she entered the University of Florida in Gainesville. She received her Bachelor of Science degree with a major in Microbiology and Cell Science and minor in Classical Studies in May of 2010. In August of 2010 she entered the Genetic Counseling program at The University of Health Science Center Graduate School of Biomedical Sciences at Houston.

Permanent Address:

5939 32nd St. North
St. Petersburg, FL, 33714